



KULYAVETC' L.V.¹, MULYAVA O.M.²

ON THE GROWTH OF A KLASSS OF DIRICHLET SERIES ABSOLUTELY CONVERGENT IN HALF-PLANE

In terms of generalized orders it is investigated a relation between the growth of a Dirichlet series $F(s) = \sum_{n=1}^{\infty} a_n \exp\{s\lambda_n\}$ with the abscissa of absolute convergence $A \in (-\infty, +\infty)$ and the growth of Dirichlet series $F_j(s) = \sum_{n=1}^{\infty} a_{n,j} \exp\{s\lambda_n\}$, $1 \leq j \leq 2$, with the same abscissa of absolute convergence if the coefficients a_n are connected with the coefficients $a_{n,j}$ by correlation

$$\beta \left(\frac{\lambda_n}{\ln(|a_n|e^{A\lambda_n})} \right) = (1 + o(1)) \prod_{j=1}^m \beta \left(\frac{\lambda_n}{\ln(|a_{n,j}|e^{A\lambda_n})} \right)^{\omega_j}, \quad n \rightarrow \infty,$$

where $\omega_j > 0$, $1 \leq j \leq m$, $\sum_{j=1}^m \omega_j = 1$.

Key words and phrases: Dirichlet series, generalized order.

¹ Ivan Franko National University, 1 Universytetska str., 79000, Lviv, Ukraine

² National University of Food Technologies, 68 Volodymyrska str., 01601, Kyiv, Ukraine

E-mail: lubov.kulyavets@gmail.com (Kulyavetc' L.V.), info@nuft.edu.ua (Mulyava O.M.)

INTRODUCTION

For an entire function $f(z) = \sum_{n=0}^{\infty} a_n z^n$ let $\rho[f]$ be its order and $\sigma[f]$ be its type. Using Hadamard's formulas for the finding of these characteristics, E.G. Calys [1] proved the following theorems.

Theorem A. Suppose that entire functions $f_1(z) = \sum_{n=0}^{\infty} a_{n,1} z^n$ and $f_2(z) = \sum_{n=0}^{\infty} a_{n,2} z^n$ have finite orders and regular growth (in sense of the equality of order $\rho[f]$ and lower order $\lambda[f]$) and the sequences $(|a_{n,1}/a_{n+1,1}|)$ and $(|a_{n,2}/a_{n+1,2}|)$ are nondecreasing for $n \geq n_0$. If

$$\ln(1/|a_n|) = (1 + o(1)) \sqrt{\ln(1/|a_{n,1}|) \ln(1/|a_{n,2}|)}$$

as $n \rightarrow \infty$, then the function f has regular growth and $\rho[f] = \sqrt{\rho[f_1]\rho[f_2]}$.

Theorem B. Suppose that functions f_1 and f_2 from Theorem A have the same order $\rho[f_1] = \rho[f_2] = \rho \in (0, +\infty)$ and the types $\sigma[f_1] = \sigma_1$, $\sigma[f_2] = \sigma_2$. Also suppose that $a_{n,1} \neq 0$ and $|a_{n,2}| \geq |a_{n,1}|/l(1/|a_{n,1}|)$ for all $n \geq n_0$, where l is slowly varying function. If

$$|a_n| = (1 + o(1)) \sqrt{|a_{n,1}| |a_{n,2}|}$$

as $n \rightarrow \infty$, then the function f has the order $\rho[f] = \rho$ and the type $\sigma[f] \leq \sqrt{\sigma_1 \sigma_2}$.

УДК 517.537.72

2010 Mathematics Subject Classification: 30B50.

In [2] Theorems A and B are generalized on the case of entire Dirichlet series of finite generalized orders by Sheremeta, moreover instead two functions f_1 and f_2 were considered $n \geq 2$ entire Dirichlet series.

Here we will obtain analogues results for Dirichlet series absolutely convergent in a half-plane.

Let $\Lambda = (\lambda_n)$ be an increasing to $+\infty$ sequence of nonnegative numbers and $S(\Lambda, A)$ be a class of Dirichlet series

$$F(s) = \sum_{n=1}^{\infty} a_n \exp\{s\lambda_n\}, \quad s = \sigma + it \quad (1)$$

with a given sequence (λ_n) of exponents and an abscissa of absolute convergence $\sigma_a = A \in (-\infty, +\infty)$ and $M(\sigma, F) = \sup\{|F(\sigma + it)| : t \in \mathbb{R}\}$ for $\sigma \in (-\infty, A)$.

By L we denote a class of positive continuous functions α on $(-\infty, +\infty)$ such that $\alpha(x) = \alpha(x_0)$ for $x \leq x_0$ and $0 < \alpha(x) \uparrow +\infty$ as $x_0 \leq x \uparrow +\infty$. We say that $\alpha \in L^0$ if $\alpha \in L$ and $\alpha((1 + o(1))x) = (1 + o(1))\alpha(x)$ as $x \rightarrow +\infty$. Finally, $\alpha \in L_{si}$, if $\alpha \in L$ and $\alpha(cx) = (1 + o(1))\alpha(x)$ as $x \rightarrow +\infty$ for each $c \in (0, +\infty)$, i. e. α is slowly increasing function. Clearly, $L_{si} \subset L^0$.

For $\alpha \in L$ and $\beta \in L$ the values

$$\rho_{\alpha, \beta}^A[F] = \overline{\lim}_{\sigma \uparrow A} \frac{\alpha(\ln M(\sigma, F))}{\beta(1/(A - \sigma))}, \quad \lambda_{\alpha, \beta}^A[F] = \underline{\lim}_{\sigma \uparrow A} \frac{\alpha(\ln M(\sigma, F))}{\beta(1/(A - \sigma))}$$

are called [3] generalized order and lower order correspondly of Dirichlet series (1) from the class $S(\Lambda, A)$.

1 ANALOGUES OF THEOREM A.

We need the following lemmas from [3].

Lemma 1.1. *Let $\alpha \in L_{si}$, $\beta \in L_{si}$ and*

$$\frac{x}{\beta^{-1}(c\alpha(x))} \uparrow +\infty, \quad \alpha\left(\frac{x}{\beta^{-1}(c\alpha(x))}\right) = (1 + o(1))\alpha(x) \quad (2)$$

as $x_0 \leq x \rightarrow +\infty$ for each $c \in (0, +\infty)$.

If $\alpha(\lambda_n) = o(\beta(\lambda_n / \ln n))$ as $n \rightarrow \infty$, then

$$\rho_{\alpha, \beta}^A[F] = k_{\alpha, \beta}^A[F] =: \overline{\lim}_{n \rightarrow \infty} \frac{\alpha(\lambda_n)}{\beta(\lambda_n / \ln(|a_n|e^{A\lambda_n}))},$$

and if, moreover, $\alpha(\lambda_{n+1}) = (1 + o(1))\alpha(\lambda_n)$ and $\frac{\ln|a_{n+1}| - \ln|a_n|}{\lambda_{n+1} - \lambda_n} \nearrow A$ as $n_0 \leq n \rightarrow \infty$, then

$$\lambda_{\alpha, \beta}^A[F] = \varkappa_{\alpha, \beta}^A[F] =: \underline{\lim}_{n \rightarrow \infty} \frac{\alpha(\lambda_n)}{\beta(\lambda_n / \ln(|a_n|e^{A\lambda_n}))}.$$

Remark 1.1 ([3]). *In order that $\lambda_{\alpha, \beta}^A[F] \geq \varkappa_{\alpha, \beta}^A[F]$, it is sufficient that $\alpha(\lambda_{n+1}) = (1 + o(1))\alpha(\lambda_n)$ as $n \rightarrow \infty$.*

Lemma 1.2. Let $\alpha \in L_{si}$, $\beta \in L_{si}$ and

$$\frac{x}{\alpha^{-1}(c\beta(x))} \uparrow +\infty, \quad \beta\left(\frac{x}{\alpha^{-1}(c\alpha(x))}\right) = (1+o(1))\beta(x) \quad (3)$$

as $x_0 \leq x \rightarrow +\infty$ for each $c \in (0, +\infty)$.

If $\alpha(\ln n) = o(\beta(\lambda_n))$ as $n \rightarrow \infty$, then

$$\varrho_{\alpha,\beta}^A[F] = k_{\alpha,\beta}^{A*}[F] =: \overline{\lim}_{n \rightarrow \infty} \frac{\alpha(\ln(|a_n|e^{A\lambda_n}))}{\beta(\lambda_n)},$$

and if, moreover, $\beta(\lambda_{n+1}) = (1+o(1))\beta(\lambda_n)$ and $\frac{\ln|a_{n+1}| - \ln|a_n|}{\lambda_{n+1} - \lambda_n} \nearrow A$ as $n_0 \leq n \rightarrow \infty$, then

$$\lambda_{\alpha,\beta}^A[F] = \varkappa_{\alpha,\beta}^{A*}[F] =: \underline{\lim}_{n \rightarrow \infty} \frac{\alpha(\ln(|a_n|e^{A\lambda_n}))}{\beta(\lambda_n)}.$$

Remark 1.2 ([3]). In order that $\lambda_{\alpha,\beta}^A[F] \geq \varkappa_{\alpha,\beta}^{A*}[F]$, it is sufficient that $\beta(\lambda_{n+1}) = (1+o(1))\beta(\lambda_n)$ as $n \rightarrow \infty$.

Suppose that $F_j \in S(\Lambda, A)$, $1 \leq j \leq m$, and

$$F_j(s) = \sum_{n=1}^{\infty} a_{n,j} \exp\{s\lambda_n\}. \quad (4)$$

Using Lemma 1.1, at first we prove the following analog of Theorem A.

Theorem 1. Let functions $\alpha \in L_{si}$ and $\beta \in L_{si}$ satisfy conditions (2), $\alpha(\lambda_n) = o(\beta(\lambda_n/\ln n))$ and $\alpha(\lambda_{n+1}) = (1+o(1))\alpha(\lambda_n)$ as $n \rightarrow \infty$. Suppose that all functions (4) have regular $\alpha\beta$ -growth (i.e. $\lambda_{\alpha,\beta}^A[F_j] = \varrho_{\alpha,\beta}^A[F_j] < +\infty$) and $\frac{\ln|a_{n+1,j}| - \ln|a_{n,j}|}{\lambda_{n+1} - \lambda_n} \nearrow A$ as $n_0 \leq n \rightarrow \infty$.

If $\omega_j > 0$, $1 \leq j \leq m$, $\sum_{j=1}^m \omega_j = 1$ and

$$\beta\left(\frac{\lambda_n}{\ln(|a_n|e^{A\lambda_n})}\right) = (1+o(1)) \prod_{j=1}^m \beta\left(\frac{\lambda_n}{\ln(|a_{n,j}|e^{A\lambda_n})}\right)^{\omega_j}, \quad n \rightarrow \infty, \quad (5)$$

then function (1) has regular $\alpha\beta$ -growth and $\varrho_{\alpha,\beta}^A[F] = \prod_{j=1}^m (\varrho_{\alpha,\beta}^A[F_j])^{\omega_j}$.

Proof. Since $\lambda_{\alpha,\beta}^A[F_j] = \varrho_{\alpha,\beta}^A[F_j] = \varrho_j < +\infty$, by Lemma 1.1 we have

$$\lim_{n \rightarrow \infty} \frac{\alpha(\lambda_n)}{\beta(\lambda_n/\ln(|a_{n,j}|e^{A\lambda_n}))} = \varrho_j.$$

Therefore, from (5) we obtain

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{\alpha(\lambda_n)} \beta\left(\frac{\lambda_n}{\ln(|a_n|e^{A\lambda_n})}\right) &= \lim_{n \rightarrow \infty} \frac{1}{\alpha(\lambda_n)} \prod_{j=1}^m \beta\left(\frac{\lambda_n}{\ln(|a_{n,j}|e^{A\lambda_n})}\right)^{\omega_j} \\ &= \lim_{n \rightarrow \infty} \prod_{j=1}^m \left(\frac{1}{\alpha(\lambda_n)} \beta\left(\frac{\lambda_n}{\ln(|a_{n,j}|e^{A\lambda_n})}\right) \right)^{\omega_j} = \prod_{j=1}^m \lim_{n \rightarrow \infty} \left(\frac{1}{\alpha(\lambda_n)} \beta\left(\frac{\lambda_n}{\ln(|a_{n,j}|e^{A\lambda_n})}\right) \right)^{\omega_j} \\ &= \prod_{j=1}^m (1/\varrho_j)^{\omega_j}, \end{aligned}$$

that is,

$$\lim_{n \rightarrow \infty} \frac{\alpha(\lambda_n)}{\beta(\lambda_n / \ln(|a_n|e^{A\lambda_n}))} = \prod_{j=1}^m \varrho_j^{\omega_j}.$$

Using Lemma 1.1 and the Remark 1.1, hence we get $\prod_{j=1}^m \varrho_j^{\omega_j} \leq \lambda_{\alpha,\beta}^A[F] \leq \varrho_{\alpha,\beta}^A[F] = \prod_{j=1}^m \varrho_j^{\omega_j}$, that is the function F has regular $\alpha\beta$ -growth and $\varrho_{\alpha,\beta}^A[F] = \prod_{j=1}^m (\varrho_{\alpha,\beta}^A[F])^{\omega_j}$. Theorem 1 is proved. \square

From (2) it follows that the function α increases less rapidly than the function β . It is easy to verify that the functions $\alpha(x) = \ln \ln x$ and $\beta(x) = \ln x$ for $x \geq x_0$ satisfy (2) and the condition $\alpha(\lambda_n) = o(\beta(\lambda_n / \ln n))$ holds as $n \rightarrow \infty$, provided $\overline{\lim}_{n \rightarrow \infty} (\ln \ln n) / \ln \lambda_n < 1$. Therefore, Theorem 1 implies the following statement.

Corollary 1.1. *Let $\overline{\lim}_{n \rightarrow \infty} (\ln \ln n) / \ln \lambda_n < 1$, $\ln \ln \lambda_{n+1} = (1 + o(1)) \ln \ln \lambda_n$ as $n \rightarrow \infty$. Suppose that $\lim_{\sigma \uparrow A} \frac{\ln \ln \ln M(\sigma, F_j)}{\ln(1/(A - \sigma))} = \varrho_j$ and $\frac{\ln |a_{n+1,j}| - \ln |a_{n,j}|}{\lambda_{n+1} - \lambda_n} \nearrow A$ as $n_0 \leq n \rightarrow \infty$ for all $1 \leq j \leq m$. If*

$$\ln \left(\frac{\lambda_n}{\ln(|a_n|e^{A\lambda_n})} \right) = (1 + o(1)) \prod_{j=1}^m \ln^{\omega_j} \left(\frac{\lambda_n}{\ln(|a_{n,j}|e^{A\lambda_n})} \right), \quad \sum_{j=1}^m \omega_j = 1,$$

as $n \rightarrow \infty$ then $\lim_{\sigma \uparrow A} \frac{\ln \ln \ln M(\sigma, F)}{\ln(1/(A - \sigma))} = \prod_{j=1}^m \varrho_j^{\omega_j}$.

For the proof of the following theorem we will use Lemma 1.2.

Theorem 2. *Let the functions $\alpha \in L_{si}$ and $\beta \in L_{si}$ satisfy the condition (3), $\alpha(\ln n) = o(\beta(\lambda_n))$ and $\beta(\lambda_{n+1}) = (1 + o(1))\beta(\lambda_n)$ as $n \rightarrow \infty$. Suppose that all functions (4) have regular $\alpha\beta$ -growth and $\frac{\ln |a_{n+1,j}| - \ln |a_{n,j}|}{\lambda_{n+1} - \lambda_n} \nearrow A$ as $n_0 \leq n \rightarrow \infty$.*

If $\omega_j > 0$, $1 \leq j \leq m$, $\sum_{j=1}^m \omega_j = 1$ and

$$\alpha \left(\ln(|a_n|e^{A\lambda_n}) \right) = (1 + o(1)) \prod_{j=1}^m \alpha^{\omega_j} \left(\ln(|a_{n,j}|e^{A\lambda_n}) \right), \quad n \rightarrow \infty, \quad (6)$$

then function (1) has regular $\alpha\beta$ -growth and $\varrho_{\alpha,\beta}^A[F] = \prod_{j=1}^m (\varrho_{\alpha,\beta}^A[F_j])^{\omega_j}$.

Proof. Since $\lambda_{\alpha,\beta}^A[F_j] = \varrho_{\alpha,\beta}^A[F_j] = \varrho_j < +\infty$, by Lemma 1.2 we have

$$\lim_{n \rightarrow \infty} \frac{\alpha \left(\ln(|a_{n,j}|e^{A\lambda_n}) \right)}{\beta(\lambda_n)} = \varrho_j.$$

Therefore, from (6), as in the proof of Theorem 1,

$$\lim_{n \rightarrow \infty} \frac{\alpha \left(\ln(|a_n|e^{A\lambda_n}) \right)}{\beta(\lambda_n)} = \prod_{j=1}^m \lim_{n \rightarrow \infty} \left(\frac{\alpha \left(\ln(|a_{n,j}|e^{A\lambda_n}) \right)}{\beta(\lambda_n)} \right)^{\omega_j} = \prod_{j=1}^m \varrho_j^{\omega_j},$$

whence, as above, we obtain the regular $\alpha\beta$ -growth of the function f and the equality $\varrho_{\alpha,\beta}^A[F] = \prod_{j=1}^m (\varrho_{\alpha,\beta}^A[F_j])^{\omega_j}$. Theorem 2 is proved. \square

From (3) it follows that the function β increases less rapidly than the function α . It is easy to verify that the functions $\alpha(x) = \ln x$ and $\beta(x) = \ln \ln x$ for $x \geq x_0$ satisfy (3). Therefore, Theorem 2 implies the following statement.

Corollary 1.2. *Let $\ln \ln n = o(\ln \ln \lambda_n)$ and $\ln \ln \lambda_{n+1} = (1 + o(1)) \ln \ln \lambda_n$ as $n \rightarrow \infty$. Suppose that $\lim_{\sigma \uparrow A} \frac{\ln \ln M(\sigma, F_j)}{\ln \ln (1/(A - \sigma))} = q_j$ and $\frac{\ln |a_{n+1,j}| - \ln |a_{n,j}|}{\lambda_{n+1} - \lambda_n} \nearrow A$ as $n_0 \leq n \rightarrow \infty$ for all $1 \leq j \leq m$. If*

$$\ln \ln \left(|a_n| e^{A\lambda_n} \right) = (1 + o(1)) \prod_{j=1}^m \ln^{\omega_j} \ln \left(|a_{n,j}| e^{A\lambda_n} \right), \quad \sum_{j=1}^m \omega_j = 1,$$

as $n \rightarrow \infty$ then $\lim_{\sigma \uparrow A} \frac{\ln \ln M(\sigma, F)}{\ln \ln (1/(A - \sigma))} = \prod_{j=1}^m q_j^{\omega_j}$.

2 ANALOGUES OF THEOREM B.

Suppose, as above, that $\alpha \in L_{si}$ and $\beta \in L_{si}$. In order to get the analogues of Theorem B, except the generalized order $\varrho_{\alpha, \beta}^A[F] \in (0, +\infty)$, it is needed to enter a (generalized) type. A definition of the type depends on what from the functions α or β grows slower.

Suppose at first that the function β increases less rapidly than the function α and define a type by the formula

$$T_{\alpha, \beta}^{A*}[F] = \overline{\lim}_{\sigma \uparrow A} \frac{\ln M(\sigma, F)}{\alpha^{-1}(\varrho_{\alpha, \beta}^A[F] \beta(1/(A - \sigma)))}.$$

Since $T_{\alpha, \beta}^{A*}[F] = \varrho_{\alpha_1, \beta_1}^A[F]$, where $\alpha_1(x) = x \notin L_{si}$ and $\beta_1(x) = \alpha^{-1}(\varrho_{\alpha, \beta}^A[F] \beta(x))$ for $x \geq x_0$, we can apply none from the lemmas indicated above. However the following statement is true [3].

Lemma 2.1. *Let $\alpha_1(x) = x$ for $x \geq x_0$, $\beta_1 \in L_{si}$ and*

$$\frac{x}{\beta_1(x)} \uparrow +\infty, \quad \beta_1 \left(\frac{x}{\beta_1(x)} \right) = (1 + o(1)) \beta_1(x), \quad x_0 \leq x \rightarrow +\infty.$$

If $\ln n = o(\beta_1(\lambda_n))$ as $n \rightarrow \infty$ then $\varrho_{\alpha_1, \beta_1}^A[F] = \overline{\lim}_{n \rightarrow \infty} \frac{\ln (|a_n| e^{A\lambda_n})}{\beta_1(\lambda_n)}$.

Since $\beta_1(x) = \alpha^{-1}(\varrho_{\alpha, \beta}^A[F] \beta(x))$ for $x \geq x_0$ then Lemma 2.1 implies the following statement.

Lemma 2.2. *Let $\alpha \in L_{si}$ and $\beta \in L_{si}$ be such that $\alpha^{-1}(c\beta(x)) \in L_{si}$ for each $c \in (0, +\infty)$ and*

$$\frac{x}{\alpha^{-1}(c\beta(x))} \uparrow +\infty, \quad \alpha^{-1} \left(c\beta \left(\frac{x}{\alpha^{-1}(c\beta(x))} \right) \right) = (1 + o(1)) \alpha^{-1}(c\beta(x)) \quad (7)$$

as $x_0 \leq x \rightarrow +\infty$ for each $c \in (0, +\infty)$. If $\ln n = o(\alpha^{-1}(c\beta(\lambda_n)))$ as $n \rightarrow \infty$ for each $c \in (0, +\infty)$, then

$$T_{\alpha, \beta}^{A*}[F] = \overline{\lim}_{n \rightarrow \infty} \frac{\ln (|a_n| e^{A\lambda_n})}{\alpha^{-1}(\varrho_{\alpha, \beta}^A[F] \beta(\lambda_n))}.$$

The following theorem generalizes Theorem B.

Theorem 3. Let $\beta \in L_{si}$, $\alpha(e^x) \in L^0$, $\alpha^{-1}(c\beta(x)) \in L_{si}$, conditions (7) hold and $\ln n = o(\alpha^{-1}(c\beta(\lambda_n)))$ as $n \rightarrow \infty$ for each $c \in (0, +\infty)$. Suppose that all Dirichlet series (4) have the same generalised order $\varrho_{\alpha,\beta}^A[F_j] = \varrho \in (0, +\infty)$ and the types $T_{\alpha,\beta}^{A*}[F_j] \in (0, +\infty)$. Suppose also that $a_{n,1} \neq 0$ for all $n \geq n_0$ and for all $2 \leq j \leq m$

$$\ln \ln \left(|a_{n,j}| e^{A\lambda_n} \right) \geq (1 + o(1)) \ln \ln \left(|a_{n,1}| e^{A\lambda_n} \right), \quad n \rightarrow \infty. \quad (8)$$

If $\omega_j > 0$, $1 \leq j \leq m$, $\sum_{j=1}^m \omega_j = 1$ and

$$\ln \left(|a_n| e^{A\lambda_n} \right) = (1 + o(1)) \prod_{j=1}^m \left(\ln \left(|a_{n,j}| e^{A\lambda_n} \right) \right)^{\omega_j}, \quad n \rightarrow \infty, \quad (9)$$

then Dirichlet series (1) has the generalized order $\varrho_{\alpha,\beta}^A[F] = \varrho$ and the type

$$T_{\alpha,\beta}^{A*}[F] \leq \prod_{j=1}^m T_{\alpha,\beta}^{A*}[F_j]^{\omega_j}.$$

Proof. Since $\alpha(e^x) \in L^0$, then for each $c \in (0, +\infty)$ we have

$$\alpha(cx) = \alpha(e^{\ln x + \ln c}) = \alpha(e^{(1+o(1)) \ln x}) = (1 + o(1))\alpha(e^{\ln x}) = (1 + o(1))\alpha(x)$$

as $x \rightarrow +\infty$, that is $\alpha \in L_{si}$. Hence it follows that $\alpha^{-1}((1 - \eta)x) = o(\alpha^{-1}(x))$ as $x \rightarrow +\infty$ for each $\eta \in (0, 1)$, because if $\alpha^{-1}((1 - \eta)x_k) \geq h\alpha^{-1}(x_k)$ for some number $h > 0$ and an increasing to $+\infty$ sequence (x_k) then $(1 - \eta)x_k \geq \alpha(h\alpha^{-1}(x_k)) = (1 + o(1))x_k$ as $k \rightarrow \infty$, that is impossible.

Therefore, conditions (7) imply the conditions (3). Indeed, if for some $c \in (0, +\infty)$, $\eta \in (0, 1)$ and an increasing to $+\infty$ sequence (x_k) the inequality

$$\beta \left(x_k / \alpha^{-1}(c\beta(x_k)) \right) \leq (1 - \eta)\beta(x_k)$$

is true then $\alpha^{-1}(c\beta(x_k / \alpha^{-1}(c\beta(x_k)))) \leq \alpha^{-1}(c(1 - \eta)\beta(x_k)) = o(\alpha^{-1}(c\beta(x_k)))$ as $k \rightarrow \infty$, that is impossible in view of (7).

Finally, from the condition $\ln n = o(\alpha^{-1}(c\beta(\lambda_n)))$ as $n \rightarrow \infty$ for each $c \in (0, +\infty)$ we have $\ln n \leq \alpha^{-1}(c\beta(\lambda_n))$ for $n \geq n_0$ and each $c \in (0, +\infty)$, that is $\alpha(\ln n) \leq c\beta(\lambda_n)$ and in view of the arbitrariness of c we obtain $\alpha(\ln n) = o(\beta(\lambda_n))$ as $n \rightarrow \infty$.

Thus, from the conditions on the functions α and β and the sequence (λ_n) in Theorem 3 the conditions of Lemma 1.2 follows.

Since all Dirichlet series (4) have the same generalized order $\varrho_{\alpha,\beta}^A[F_j] = \varrho \in (0, +\infty)$, then by Lemma 1.2 for every $\varrho_1 > \varrho$ and all $n \geq n_0(\varrho_1)$ we have $\ln \left(|a_{n,j}| e^{A\lambda_n} \right) \leq \alpha^{-1}(\varrho_1\beta(\lambda_n))$. Therefore, from (9) we obtain

$$\begin{aligned} \varrho_{\alpha,\beta}^A[F] &= \overline{\lim}_{n \rightarrow \infty} \frac{\alpha \left(\ln \left(|a_n| e^{A\lambda_n} \right) \right)}{\beta(\lambda_n)} = \overline{\lim}_{n \rightarrow \infty} \frac{1}{\beta(\lambda_n)} \alpha \left(\prod_{j=1}^m \left(\ln \left(|a_{n,j}| e^{A\lambda_n} \right) \right)^{\omega_j} \right) \\ &= \overline{\lim}_{n \rightarrow \infty} \frac{1}{\beta(\lambda_n)} \alpha \left(\exp \left\{ \sum_{j=1}^m \omega_j \ln \ln \left(|a_{n,j}| e^{A\lambda_n} \right) \right\} \right) \\ &\leq \overline{\lim}_{n \rightarrow \infty} \frac{1}{\beta(\lambda_n)} \alpha \left(\exp \left\{ \sum_{j=1}^m \omega_j \ln \alpha^{-1}(\varrho_1\beta(\lambda_n)) \right\} \right) = \varrho_1, \end{aligned}$$

that is in view of the arbitrariness of ϱ_1 we obtain the inequality $\varrho_{\alpha,\beta}^A[F] \leq \varrho$.

On the other hand, in view of the conditions (8) and $\alpha(e^x) \in L^0$ we have

$$\begin{aligned} \varrho_{\alpha,\beta}^A[F] &= \overline{\lim}_{n \rightarrow \infty} \frac{1}{\beta(\lambda_n)} \alpha \left(\exp \left\{ \sum_{j=1}^m \omega_j \ln \ln \left(|a_{n,j}| e^{A\lambda_n} \right) \right\} \right) \\ &= \overline{\lim}_{n \rightarrow \infty} \frac{1}{\beta(\lambda_n)} \alpha \left(\exp \left\{ \omega_1 \ln \ln \left(|a_{n,1}| e^{A\lambda_n} \right) + \sum_{j=2}^m \omega_j \ln \ln \left(|a_{n,j}| e^{A\lambda_n} \right) \right\} \right) \\ &\geq \overline{\lim}_{n \rightarrow \infty} \frac{1}{\beta(\lambda_n)} \alpha \left(\exp \left\{ \omega_1 \ln \ln \left(|a_{n,1}| e^{A\lambda_n} \right) + \sum_{j=2}^m \omega_j (1 + o(1)) \ln \ln \left(|a_{n,1}| e^{A\lambda_n} \right) \right\} \right) \\ &= \overline{\lim}_{n \rightarrow \infty} \frac{1}{\beta(\lambda_n)} \alpha \left(\exp \left\{ (1 + o(1)) \sum_{j=1}^m \omega_j \ln \ln \left(|a_{n,1}| e^{A\lambda_n} \right) \right\} \right) \\ &= \overline{\lim}_{n \rightarrow \infty} \frac{(1 + o(1))}{\beta(\lambda_n)} \alpha \left(\exp \left\{ \sum_{j=1}^m \omega_j \ln \ln \left(|a_{n,1}| e^{A\lambda_n} \right) \right\} \right) \\ &= \overline{\lim}_{n \rightarrow \infty} \frac{\alpha \left(\ln \left(|a_{n,1}| e^{A\lambda_n} \right) \right)}{\beta(\lambda_n)} = \varrho. \end{aligned}$$

Thus, $\varrho_{\alpha,\beta}^A[F] = \varrho$ and for $T_{\alpha,\beta}^{A*}[F]$ by Lemma 2.2 from (9) we obtain

$$\begin{aligned} T_{\alpha,\beta}^{A*}[F] &= \overline{\lim}_{n \rightarrow \infty} \frac{\ln \left(|a_n| e^{A\lambda_n} \right)}{\alpha^{-1}(\varrho_{\alpha,\beta}^A[F] \beta(\lambda_n))} = \overline{\lim}_{n \rightarrow \infty} \frac{1}{\alpha^{-1}(\varrho_{\alpha,\beta}^A[F] \beta(\lambda_n))} \prod_{j=1}^m \left(\ln \left(|a_{n,j}| e^{A\lambda_n} \right) \right)^{\omega_j} \\ &= \overline{\lim}_{n \rightarrow \infty} \prod_{j=1}^m \left(\frac{\ln \left(|a_{n,j}| e^{A\lambda_n} \right)}{\alpha^{-1}(\varrho_{\alpha,\beta}^A[F] \beta(\lambda_n))} \right)^{\omega_j} \leq \prod_{j=1}^m \overline{\lim}_{n \rightarrow \infty} \left(\frac{\ln \left(|a_{n,j}| e^{A\lambda_n} \right)}{\alpha^{-1}(\varrho_{\alpha,\beta}^A[F] \beta(\lambda_n))} \right)^{\omega_j} = \prod_{j=1}^m T_{\alpha,\beta}^{A*}[F_j]^{\omega_j}. \end{aligned}$$

The proof of Theorem 3 is complete. □

It is easy to verify that the functions $\alpha(x) = \ln x$ and $\beta(x) = \ln \ln x$ for $x \geq x_0$ satisfy the conditions of Theorem 3. Therefore, the following statement is true.

Corollary 2.1. *Let Diriclet series (4) be such that for all $1 \leq j \leq m$*

$$\lim_{\sigma \uparrow A} \frac{\ln \ln M(\sigma, F_j)}{\ln \ln (1/(A - \sigma))} = \varrho, \quad \lim_{\sigma \uparrow A} \frac{\ln M(\sigma, F_j)}{\ln^\varrho (1/(A - \sigma))} = T_j,$$

and $\ln n = O(\ln \ln \lambda_n)$ as $n \rightarrow \infty$. Then the conditions (8) and (9) imply

$$\lim_{\sigma \uparrow A} \frac{\ln \ln M(\sigma, F)}{\ln \ln (1/(A - \sigma))} = \varrho, \quad \lim_{\sigma \uparrow A} \frac{\ln M(\sigma, F)}{\ln^\varrho (1/(A - \sigma))} \leq \prod_{j=1}^m T_j^{\omega_j}.$$

Since $\varrho_{\alpha,\beta}^A[F] = \overline{\lim}_{\sigma \uparrow A} \frac{\ln \exp\{\alpha(\ln M(\sigma, F))\}}{\ln \exp\{\beta(1/(A - \sigma))\}}$, we define the type also by the formula

$$T_{\alpha,\beta}^A[F] = \overline{\lim}_{\sigma \uparrow A} \frac{\exp\{\alpha(\ln M(\sigma, F))\}}{\exp\{\varrho_{\alpha,\beta}^A[F] \beta(1/(A - \sigma))\}},$$

and for the finding by the coefficients we use Lemma 1.1. We obtain the following statement.

Lemma 2.3. Suppose that the function $e^{\alpha(x)}$ and $e^{\beta(x)}$ belongs to L_{si} and

$$\frac{x}{\beta^{-1}(\ln c + \alpha(x))} \uparrow +\infty, \quad \exp \left\{ \alpha \left(\frac{x}{\beta^{-1}(\ln c + \alpha(x))} \right) \right\} = (1 + o(1))e^{\alpha(x)} \quad (10)$$

as $x \rightarrow +\infty$ for each $c \in (0, +\infty)$. If $\exp\{\alpha(\lambda_n)\} = o(\exp\{\beta(\lambda_n/\ln n)\})$ as $n \rightarrow \infty$ then

$$T_{\alpha,\beta}^A[F] = \overline{\lim}_{n \rightarrow \infty} \frac{\exp\{\alpha(\lambda_n)\}}{\exp \left\{ \varrho_{\alpha,\beta}^A[F] \beta \left(\frac{\lambda_n}{\ln(|a_n|e^{A\lambda_n})} \right) \right\}}.$$

Theorem 4. Let the function $e^{\alpha(x)}$ and $e^{\beta(x)}$ belongs to L_{si} , the conditions (2) and (10) hold and $\alpha(\lambda_n) = o(\beta(\lambda_n/\ln n))$ as $n \rightarrow \infty$. Suppose that all Dirichlet series (4) have the same generalized order $\varrho_{\alpha,\beta}^A[F_j] = \varrho \in (0, +\infty)$ and the types $T_{\alpha,\beta}^A[F_j] \in (0, +\infty)$. Suppose also that $a_{n,1} \neq 0$ for all $n \geq n_0$ and for all $2 \leq j \leq m$

$$\beta \left(\frac{\lambda_n}{\ln(|a_{n,j}|e^{A\lambda_n})} \right) \leq (1 + o(1))\beta \left(\frac{\lambda_n}{\ln(|a_{n,1}|e^{A\lambda_n})} \right), \quad n \rightarrow \infty. \quad (11)$$

If $\omega_j > 0, 1 \leq j \leq m, \sum_{j=1}^m \omega_j = 1$ and

$$\exp \left\{ \beta \left(\frac{\lambda_n}{\ln(|a_n|e^{A\lambda_n})} \right) \right\} = (1 + o(1)) \prod_{j=1}^m \exp \left\{ \omega_j \beta \left(\frac{\lambda_n}{\ln(|a_{n,j}|e^{A\lambda_n})} \right) \right\} \quad (12)$$

as $n \rightarrow \infty$ then Dirichlet series (1) has the generalized order $\varrho_{\alpha,\beta}^A[F] = \varrho$ and type

$$T_{\alpha,\beta}^A[F] \leq \prod_{j=1}^m T_{\alpha,\beta}^A[F_j]^{\omega_j}.$$

Proof. From (12) we have

$$\beta \left(\frac{\lambda_n}{\ln(|a_n|e^{A\lambda_n})} \right) = \sum_{j=1}^m \omega_j \beta \left(\frac{\lambda_n}{\ln(|a_{n,j}|e^{A\lambda_n})} \right) + o(1) \quad (13)$$

as $n \rightarrow \infty$. Therefore, by Lemma 1.1

$$\frac{1}{\varrho_{\alpha,\beta}^A[F]} = \lim_{n \rightarrow \infty} \frac{1}{\alpha(\lambda_n)} \beta \left(\frac{\lambda_n}{\ln(|a_n|e^{A\lambda_n})} \right) \geq \sum_{j=1}^m \lim_{n \rightarrow \infty} \frac{\omega_j}{\alpha(\omega_n)} \beta \left(\frac{\lambda_n}{\ln(|a_{n,j}|e^{A\lambda_n})} \right) = \frac{1}{\varrho}.$$

On the other hand, in view of (11) from (13) we obtain

$$\frac{1}{\varrho_{\alpha,\beta}^A[F]} \leq \lim_{n \rightarrow \infty} \sum_{j=1}^m \frac{\omega_j}{\alpha(\lambda_n)} \beta \left(\frac{\lambda_n}{\ln(|a_{n,1}|e^{A\lambda_n})} \right) = \frac{1}{\varrho},$$

that is $\varrho_{\alpha,\beta}^A[F] = \varrho$. From (12) and Lemma 2.3 also it follows that

$$\begin{aligned} \frac{1}{T_{\alpha,\beta}^A[F]} &= \lim_{n \rightarrow \infty} \frac{1}{\exp\{\alpha(\lambda_n)\}} \exp \left\{ \varrho \beta \left(\frac{\lambda_n}{\ln(|a_n|e^{A\lambda_n})} \right) \right\} \\ &= \lim_{n \rightarrow \infty} \frac{1}{\exp\{\alpha(\lambda_n)\}} \prod_{j=1}^m \exp \left\{ \varrho \omega_j \beta \left(\frac{\lambda_n}{\ln(|a_{n,j}|e^{A\lambda_n})} \right) \right\} \\ &\geq \prod_{j=1}^m \lim_{n \rightarrow \infty} \left(\frac{\exp \left\{ \varrho \beta \left(\frac{\lambda_n}{\ln(|a_{n,j}|e^{A\lambda_n})} \right) \right\}}{\exp\{\alpha(\lambda_n)\}} \right)^{\omega_j} = \prod_{j=1}^m \left(\frac{1}{T_{\alpha,\beta}^A[F_j]} \right)^{\omega_j}. \end{aligned}$$

Theorem 4 is proved. \square

It is easy to verify that the functions $\alpha(x) = \ln \ln x$ and $\beta(x) = \ln \ln x$ for $x \geq x_0$ satisfy the conditions (2) and (10). The condition $\alpha(\lambda_n) = o(\beta(\lambda_n / \ln n))$ as $n \rightarrow \infty$ holds, provided $\overline{\lim}_{n \rightarrow \infty} (\ln \ln n) / \ln \lambda_n < 1$. Therefore, Theorem 4 implies the following statement.

Corollary 2.2. Let $\overline{\lim}_{n \rightarrow \infty} (\ln \ln n) / \ln \lambda_n < 1$ and for all $1 \leq j \leq m$

$$\overline{\lim}_{\sigma \uparrow A} \frac{\ln \ln \ln \ln M(\sigma, F_j)}{\ln \ln (1/(A - \sigma))} = \varrho, \quad \overline{\lim}_{\sigma \uparrow A} \frac{\ln \ln \ln M(\sigma, F_j)}{\ln^\varrho (1/(A - \sigma))} = T_j \in (0, +\infty).$$

Suppose that $a_{n,1} \neq 0$ for all $n \geq n_0$ and for all $2 \leq j \leq m$

$$\ln \ln \frac{\lambda_n}{\ln (|a_{n,j}|e^{A\lambda_n})} \leq (1 + o(1)) \ln \ln \frac{\lambda_n}{\ln (|a_{n,1}|e^{A\lambda_n})}, \quad n \rightarrow \infty.$$

If $\omega_j > 0, 1 \leq j \leq m, \sum_{j=1}^m \omega_j = 1$ and

$$\ln \frac{\lambda_n}{\ln (|a_n|e^{A\lambda_n})} = (1 + o(1)) \prod_{j=1}^m \left(\ln \frac{\lambda_n}{\ln (|a_{n,j}|e^{A\lambda_n})} \right)^{\omega_j}$$

as $n \rightarrow \infty$ then

$$\overline{\lim}_{\sigma \uparrow A} \frac{\ln \ln \ln \ln M(\sigma, F)}{\ln \ln (1/(A - \sigma))} = \varrho, \quad \overline{\lim}_{\sigma \uparrow A} \frac{\ln \ln \ln M(\sigma, F)}{\ln^\varrho (1/(A - \sigma))} \leq \prod_{j=1}^m T_j^{\omega_j}.$$

REFERENCES

- [1] Calys E.G. *A note on the order and type of integral functions*. Riv. Mat. Univer. Parma 1964, 5 (2), 133–137.
- [2] Kulyavec' L.V., Mulyava O.M. *On the growth of a class of entire Dirichlet series*. Carpathian Math. Publ. 2014, 6 (2), 300–309. doi:10.15330/cmp.6.2.300-309 (in Ukrainian)
- [3] Gal' Yu.M. *On the growth of analytic functions, represented by Dirichlet series absolutely convergent in half-plane*. Drohobych, 1980. (in Russian)

Received 17.04.2015

Revised 01.06.2017

Кулявечь Л.В., Мулява О.М. Про зростання одного класу абсолютно збіжних у півплощині рядів Діріхле // Карпатські матем. публ. — 2017. — Т.9, №1. — С. 63–71.

У термінах узагальнених порядків досліджено зв'язок між зростанням ряду Діріхле $F(s) = \sum_{n=1}^{\infty} a_n \exp\{s\lambda_n\}$ з абсцисою абсолютної збіжності $A \in (-\infty, +\infty)$ і зростанням рядів Діріхле $F_j(s) = \sum_{n=1}^{\infty} a_{n,j} \exp\{s\lambda_n\}, 1 \leq j \leq 2$, з такою ж абсцисою абсолютної збіжності, якщо, наприклад, коефіцієнти a_n пов'язані з коефіцієнтами $a_{n,j}$ співвідношенням

$$\beta \left(\frac{\lambda_n}{\ln (|a_n|e^{A\lambda_n})} \right) = (1 + o(1)) \prod_{j=1}^m \beta \left(\frac{\lambda_n}{\ln (|a_{n,j}|e^{A\lambda_n})} \right)^{\omega_j}, \quad n \rightarrow \infty,$$

де $\omega_j > 0, 1 \leq j \leq m, \sum_{j=1}^m \omega_j = 1$.

Ключові слова і фрази: ряд Діріхле, узагальнений порядок.