

UDC: 618.39-021.3-084:612.826.33
DOI: 10.24061/2413-4260.XIII.4.50.2023.17

FUNCTIONAL STATE OF THE PINEAL GLAND
IN WOMEN WITH THREATENED ABORTION
IN THE FIRST TRIMESTER – FURTHER
PROSPECTS OF PREGNANCY

A. V. Pushkashu

Bukovinian State Medical University
(Chernivtsi, Ukraine)

Summary

Melatonin is an important hormone for normal pregnancy development, but the influence of serotonin remains poorly understood. Low melatonin production can lead to spontaneous abortion by stimulating progesterone secretion and inhibiting prostaglandin synthesis. Melatonin has an effect on the induction and blocking of apoptosis of the villous cytotrophoblast and promotes the development of the syncytiotrophoblast. Analysis of the frequency of positive pregnancy showed a significant impact on the course of pregnancy and childbirth, compliance with sleep and rest also contribute to a positive effect, but there is no general concept, which makes the subject relevant.

The aim of the study. To evaluate the functional state of the pineal gland, in particular, the amount of melatonin and serotonin in the blood of pregnant women at 10-12 weeks of gestation with a threat of abortion, which was manifested by bloody discharge or formation of retrochorionic hematoma, and the consequences of pregnancy.

Materials and methods. Two groups were formed: I main group – 40 pregnant women with a threat of abortion and formation of retrochorionic hematoma, II control group – 44 healthy pregnant women. Obstetric examination, enzyme-linked immunosorbent assay for determination of melatonin, serotonin, chorionic gonadotropin, and ultrasound were performed.

Informed consent of pregnant women was obtained with adequate explanation of the purpose, objectives, methods and scope of laboratory and instrumental research methods.

The study protocol was approved by the Commission on Biomedical Ethics of the Bukovinian State Medical University, 2015 (minutes of the Commission meeting No 2 from the 19.10.2023 year). The study was conducted in accordance with the basic provisions of GCP (1996), the Convention of the Council of Europe on Human Rights and Biomedicine (April 4, 1997), the Declaration of Helsinki of the World Medical Association for the Ethical Principles of Scientific Medical Research Involving Human Subjects (1964-2008), the Order of the Ministry of Health of Ukraine 690 of September 23, 2009 (as amended by the Order of the Ministry of Health of Ukraine 523 of July 12, 2012).

Statistical analysis was performed using generally accepted methods of variance statistics. Reliability was assessed by Student's *t*-test. Differences were considered significant if the significance level was $p \leq 0.05$. We also performed correlation analysis in Microsoft Excel, using the method of squares (Pearson's method) to determine the correlation coefficient.

The work is a fragment of the research work of the Department of Obstetrics and Gynecology, UDC 618.1-053.34 + 618.1-0536 / 8]. –07-084-08 «Prevention, diagnosis and treatment of disorders of the perinatal period and reproductive system of women and adolescent girls» (No. 201110H, state registration number 0111U006499. The term of implementation is 02.2011-12.2015).

Results of the study.

According to the obtained results, retrochorionic hematoma up to 1 cm is detected three times more often than over 1 cm, while the level of melatonin is not significantly lower; but with retrochorionic hematoma over 1 cm a significant decrease is observed in cases of further normal pregnancy and childbirth. With further complications of pregnancy in retrochoroidal hematoma up to 1 cm, there is a significant significant decrease in melatonin ($p < 0.001$), in contrast to retrochoroidal hematoma over 1 cm, where no significant difference was found in the complicated course of pregnancy ($p > 0.05$).

Serotonin levels were elevated in all subgroups regardless of hematoma size and subsequent pregnancy outcome.

Conclusions. In case of threatened abortion in the first trimester with formation of retrochorionic hematoma, the risk of complicated pregnancy remains in 60.87 % of cases. Determination of pineal hormones revealed a significant increase in serotonin with a simultaneous significant decrease in melatonin, while maintaining a positive relationship between the hormones, indicating a disruption of melatonin synthesis in the central nervous system. This is confirmed by the normal level of chorionic gonadotropin and the preservation of embryonic viability. Since the disruption of hormone formation in the pineal gland has been established, adherence to the recommendations for adequate rest may positively affect the further outcome of pregnancy and childbirth.

Key words: Melatonin; Serotonin; Retrochorionic Hematoma; Threatened Abortion

Introduction

The pineal gland (epiphysis) belongs to the midbrain and functions as an endocrine gland whose main functions are to regulate circadian (daily) biological rhythms, normalize immune responses, endocrine connections, metabolism, neutralize the effects of stress hormones, and adapt the body to changing light conditions that determine its activity. The main hormones of the pineal gland (PG) are melatonin (MT) and serotonin (ST).

It is believed that the cells of the pineal gland secrete ST during the day and begin to synthesize MT in the dark, while information about the level of external illumination is transmitted from the retina to the pineal gland via sympathetic fibers, which is important for the normal functioning of the gland.

MT, also known as 5-methoxy-N-acetyltryptamine, is a neuroendocrine hormone produced mainly in the pineal gland that is regulated by light levels and follows a circadian

rhythm [1, 2]. It is an endogenously produced indolamine containing two functional groups: a 5-methoxy group and an N-acetyl side chain, which determine its specificity and amphiphilicity. MT biosynthesis starts with its precursor tryptophan and consists of four enzymatic steps. In the biosynthesis of MT, serotonin N-acetyltransferase is the rate-limiting enzyme and ST is an important intermediate [3, 4, 5]. The half-life of MT in the circulation is usually short, ranging from 30 to 60 minutes [6]. It has amphiphilic properties that allow MT to easily diffuse and cross all membranes. Approximately 70 % of MT binds to albumin, and the remaining 30 % diffuses into surrounding tissues after release into the bloodstream [7]. Due to its multiple actions, MT acts as a circadian regulator, natural antioxidant, anti-inflammatory and anti-neoplastic agent, among many other functions [8, 9]. Regarding the anti-inflammatory and circadian regulatory functions, a study by Hobson et al. 2018 shows a correlation between the circadian clock and the immune system. It is emphasized that MT is powerful in terms of circadian regulation of lymphocyte proliferation, enhancing phagocytosis and stimulating cytokine production, which is essential for the normal development of pregnancy. Thus, in the context of obstetrics and fetal growth, MT is important for the prevention of preterm labor, gestational diabetes, and pre-eclampsia [10, 11, 12].

The majority of endogenous MT production occurs in the pineal gland, which is regulated by light levels. However, studies by McCarthy R, 2019 using human placenta and ovarian tissue report melatonin-synthesizing enzymes capable of converting ST to MT. Such non-pineal sources are not regulated by light and therefore do not show circadian fluctuations. However, no systematic evaluation of the metabolism of endogenous circulating MT during the course of pregnancy and identification of the placenta as a source of MT during pregnancy has been reported [26].

Low production of MT in the pineal gland may lead to spontaneous abortion in cases where uterine abnormalities and/or chromosomal abnormalities have been excluded. Recurrent pregnancy loss is thought to be associated with oxidative damage and immunologic imbalance; in this context, MT acts as a direct potent free radical scavenger with immunomodulatory effects. In addition, MT stimulates progesterone secretion and inhibits the synthesis of prostaglandins, which are potent inducers of miscarriage and preterm labor [13, 14, 15].

MT has an effect on inducing and blocking apoptosis of the villous cytotrophoblast, promotes the development of syncytiotrophoblast through paracrine, autocrine and/or intracrine action of MT1 and MT2 receptors in the placenta [16, 17, 18], which are used by the placenta to maintain the balance between villous cytotrophoblast and syncytiotrophoblast. This is how the placenta develops normally. The fusion of villous cytotrophoblasts forms a syncytiotrophoblast through a regulated MT process [19].

In addition, placental MT interacts with MT1 and MT2 receptors and reactive oxygen species to reduce oxidative damage to the placenta. Because MT protects the placenta from antioxidant effects, it has been recognized as a regulator of placental homeostasis [20]. In the first

trimester of pregnancy, the MT1 receptor is more important in promoting synchronization of the villous cytotrophoblast by protecting trophoblastic cells from oxidative damage and promoting apoptosis in altered cells [21].

The involvement of ST in pregnancy is not well understood. ST is a phylogenetically ancient neurotransmitter that is widely distributed in key brain regions and influences emotional state, impulsivity, learning and memory, attention, sleep, aggression, and neurovegetative control [22]. It is also found in enterochromaffin cells of the intestine – apodocytes, a set of which is a special endocrine organ belonging to the APUD (Amine Precursor Uptake and Decarboxylation) system, blood platelets, mast cells of connective tissue, bronchial epithelium, ovaries, thyroid cells, which also belong to the APUD system [23]. It is believed that the threat of miscarriage increases the number of ST, which may lead to the induction of habitual abortions [24].

The evidence base for the effectiveness of MT and ST is currently being formed, however, an analysis of the frequency of positive pregnancy in patients showed a significant impact on the course of pregnancy and childbirth, the presence of complications in the case of synthesis disorders [25, 26]. Compliance with sleep and rest regimens also contributes to a positive effect.

The aim of this study was to assess the functional state of the gonads, in particular the amount of MT and ST in the blood of pregnant women at 10-12 weeks' gestation with threatened abortion, manifested by bloody discharge or formation of retrochorionic hematoma (RCH), and the outcome of pregnancy.

Materials and methods. To achieve this goal, we formed two groups: I Main Group (MG) – 40 pregnant women with signs of threatened abortion, especially the presence of bloody discharge or RHG, II Control Group (CG) – 44 pregnant women without complications in the first trimester of pregnancy. As known from the studies of De Almeida Chuffa et al. 2019, the main functions of MT, such as decidualization and implantation, are important, which makes this period crucial for the normal development of pregnancy, so the results can be used to predict the further course of pregnancy and birth outcomes [7]. The two groups are representative in terms of age, lifestyle, and obstetric and gynecologic history.

In the course of the study, we used a general clinical examination (clarification of complaints, anamnesis, general somatic medical examination), obstetric examination, special research methods to determine the number of MT, ST, ultrasound to determine the condition of the fetus, the size of the RCH. Chorionic gonadotropin (hCG) measurement to determine the functional state of the chorion.

Preparation of patients to determine the amount of MT, ST, hCG according to the general rules. Time of sampling: 8.00-9.00 a.m., material: blood serum.

Melatonin and serotonin ELISA kits are designed for determination of MT and ST levels in blood of pregnant women. The principle of analysis is based on enzyme-linked immunosorbent assay.

Determination of free beta-hCG is part of prenatal screening in the first trimester at 10-12 weeks of gestation, calculated by PRISCA or ASTRAIA programs. It is used for early confirmation of pregnancy and dynamic monitoring of its course, assessment of the risk of fetal anomalies and complicated pregnancy.

Considering that the determination of hCG, MT and ST in CG was performed at 10-12 weeks of pregnancy during the first prenatal screening, the results of the study were compared for data reliability with similar indicators of 23 pregnant women in MG who presented with complaints at 10-12 weeks of pregnancy.

Statistical analysis was performed using generally accepted methods of variance statistics. Reliability was assessed using Student's t-test. Differences were considered significant at a significance level of $p \leq 0.05$. Correlation analysis was also performed in Microsoft Excel, using the method of squares (Pearson's method) to determine the correlation coefficient.

Informed consent of pregnant women was obtained with adequate explanation of the purpose, objectives, methods and scope of laboratory and instrumental research methods.

The study protocol was approved by the Commission on Biomedical Ethics of the Bukovinian State Medical University, 2015 (minutes of the Commission meeting No 2 from the 19.10.2023 year). The study was conducted in accordance with the basic provisions of GCP (1996), the Convention of the Council of Europe on Human Rights and Biomedicine (April 4, 1997), the Declaration of

Helsinki of the World Medical Association for the Ethical Principles of Scientific Medical Research Involving Human Subjects (1964-2008), the Order of the Ministry of Health of Ukraine 690 of September 23, 2009 (as amended by the Order of the Ministry of Health of Ukraine 523 of July 12, 2012). The work is a fragment of the research work «Prevention, diagnostics and treatment of perinatal and reproductive disorders of women and adolescent girls» (No. 201110H, state registration number 0111U006499).

Results of the study and their discussion. Determination of MT, ST was performed in pregnant from MG and CG. To determine the functional state of chorion, hCG was evaluated.

According to the studies of Ejaz, 2020 [9], placental synthesis of MT was established, so it is obvious that in case of impaired blood supply to the chorion, the level of hormones formed in it should decrease. However, the results obtained were somewhat different, indicating the possibility of compensatory extraplacental synthesis of MT.

The results of pregnancy and childbirth in both groups are not encouraging: we considered the CG as a group of healthy pregnant women whose first trimester of pregnancy was uneventful, but the results of further pregnancy and childbirth did not differ significantly between groups. In this regard, we formed subgroups according to the subsequent normal course of pregnancy and childbirth and the complicated course of pregnancy and childbirth. The complications are shown in Table 1.

Table 1

The results of pregnancy and childbirth

| | Main Group (n=23) | Control Group (n=44) |
|---|-------------------|--|
| Normal delivery | 9 (39,13 %) | 21 (47,72 %) |
| Premature rupture of the membranes | 3 (13,04 %) | 4 (9,09 %) |
| Abnormalities of labor activity | 2 (8,7 %) | 7 (15,91 %) |
| Polyhydramnios | 1 (4,35 %) | 1 (2,27 %) |
| Preterm labor | 1 (4,35 %) | 3 (6,82 %) |
| Premature detachment of the normal placenta | 1 (4,35 %) | 2 (4,55 %) (MT level was significantly reduced in one case) |
| Placental dysfunction | 3 (13,04 %) | - |
| Fetal growth retardation syndrome | 1 (4,35 %) | - |
| Childbirth Distress | 1 (4,35 %) | 4 (9,09 %) |
| Gestosis | 1 (4,35 %) | 1 (2,27 %) |

From the results obtained, it can be concluded that there is a slight difference in the number of complications between the two groups, except for placental dysfunction and fetal growth retardation syndrome, which were found only in MG, and gestosis, which was found twice as often in MG. However, obstructed labor and abnormal uterine activity were twice as common in CG. Thus, complications related to placental function were observed in MG, confirming the results of our study on the effect of MT on chorionic and placental development.

Pregnancy and delivery were completed by cesarean section in 12 cases of CG (27.3 %), half of which were emergency cesarean sections (50 %). The situation is similar in MG: 12 cases of cesarean section (30 %), but one case of

emergency cesarean section (8.3 %). The main indications for cesarean section in both groups were uterine scarring, abnormal uterine contractions that could not be corrected by medication, and breech presentation. In both groups, cesarean section was performed with premature abruption of the normally positioned placenta. There were no significant differences in the number or outcome of operative deliveries.

As a result of the study, it was found that the level of MT and ST depends on the size of the RCH, so we calculated the PG hormones depending on the size of the RCH – up to 1 cm and more than 1 cm (Table 2).

To determine the reliability of the indicators, the results were compared with the CG, where pregnancy and delivery were normal.

According to the results obtained, the presence of RCH up to 1 cm was found to be three times more frequent, while the level of MT was not significantly lower compared to CG ($p > 0.05$), but in the case of RCH over 1 cm and dangerous for the embryo, a significant decrease was observed in the

case of further normal pregnancy and delivery. With further pregnancy complications, a significant decrease in MT was observed for RCH up to 1 cm ($p < 0.001$), in contrast to RCH over 1 cm, where no significant difference was found for complicated pregnancies ($p > 0.05$).

Table 2

Comparative characteristics of melatonin (pg/ml) and serotonin (ng/ml) in the main and control groups with different sizes of retrochorionic hematoma in normal and complicated pregnancy and delivery

| | CG (n=44: -21 with normal course, -23 with complicated course) | MG RCH up to 1 cm (n=17: -6 with normal course, -11 with complicated course) | MG RCH over 1 cm (n=6: -3 with normal course, -3 with complicated course) |
|-----------|--|--|---|
| Melatonin | -124,9 ± 15,84 -123,7 ± 14,29 | -82,99 ± 27,52 -68,7 ± 7,1** | -86,35 ± 7,55* -101,6 ± 33,32 |
| Serotonin | -354,48 ± 61,57 -654 ± 138,57* | -537,13 ± 168,76 -690,03 ± 105,2** | -783,89 ± 160,93** -622,17 ± 147,65* |

Note^ * $p < 0,05$ ** $p < 0,001$

In his study, McCarthy, 2019, argues that just as placental MT can cross into maternal blood, maternal plasma MT can cross into the placenta unchanged. In the placenta, MT is thought to protect the mononuclear villous cytotrophoblasts from apoptosis, allowing them to continuously regenerate and fuse with the syncytiotrophoblast layer, keeping it healthy. This layer is in direct contact with the mother's blood and ensures the exchange of gases, nutrients and wastes. One possible explanation for the initial decrease in serum melatonin levels in the first trimester is that placental mitochondria and polymorphonuclear leukocytes generate large amounts of superoxide free radicals, which may lead to a temporary decrease in maternal serum melatonin levels to protect developing tissues from oxidative stress [22]. The results of our study partially confirmed the data of

the study – in MG with RCH up to 1 cm with subsequent complicated pregnancy and delivery, the level of MT is significantly reduced, so the next task of our study was to determine ST, which is an intermediate hormone for the synthesis of MT.

The amount of ST, in contrast to melatonin, is increased in all subgroups, despite the fact that ST is an intermediate hormone in the synthesis of MT, so we determined the correlation in the subgroups. An almost twofold increase in ST in MG compared to CG, where pregnancy and labor were uncomplicated, was observed both at RCH over 1 cm and at RCH less than 1 cm. An increase in ST was also observed in CG where pregnancy and labor were complicated. The results of the correlation analysis are presented in Table 3.

Table 3

Results of correlation analysis of melatonin (pg/ml) and serotonin (ng/ml) in the main and control groups with different sizes of retrochorionic hematoma in normal and complicated pregnancy and delivery

| | CG | MG RCH up to 1 cm | MG RCH over 1 cm |
|--------------------------------------|-----------|-------------------|------------------|
| With normal course of pregnancy | R = -0,14 | R = -0,05 | R = 0,97 |
| With complicated course of pregnancy | R = 0,39 | R = -0,3 | R = 0,77 |

Positive correlations were found in CG with complicated course of second half of pregnancy, delivery and strong positive correlations in case of RCH over 1 cm, regardless of further course of pregnancy. Therefore, we can conclude that MT synthesis in the CNS is impaired. Our opinion is confirmed by the results of hCG, which in pregnant women from MG was 53.63 ± 2.56 ng/ml and in CG – 45.85 ± 3.71 ng/ml. Thus, no significant difference was found ($p=0.05$) despite the presence of RCH. Also, no significant difference was found in the subsequent normal and complicated course of pregnancy.

Normal hCG levels may be evidence of chorionic function and pregnancy maintenance despite a significant decrease in MT. A significant increase in the level of ST and an insufficient level of MT may indicate a disturbance in the synthesis of MT of central genesis, while an increase in MT at an RCH over 1 cm may be

due to an increase in synthesis in the CNS as a stress response to compensate and normalize the fetal state in extreme conditions of blood supply disruption for further pregnancy development. The results obtained are important for the development of recommendations for the observance of sleep and rest regimens for the full formation of MT in the CNS.

Conclusions.

1. In the case of threatened abortion in the first trimester with the formation of retrochorionic hematoma, the risk of complicated pregnancy remains in 60.87 % of cases.

2. Determination of pineal hormones revealed a significant increase in serotonin with a simultaneous significant decrease in melatonin with a positive relationship between the hormones, indicating a disorder

of melatonin synthesis in the central nervous system. This is confirmed by the normal level of chorionic gonadotropin and the preservation of embryonic viability.

3. Since the formation of hormones in the pineal gland has been found to be impaired, adherence to the recommendations for adequate rest can positively influence the subsequent outcome of pregnancy and childbirth.

References:

- Hobson SR, Gurusinge S, Lim R, Alers NO, Miller SL, Kingdom JC, et al. Melatonin improves endothelial function in vitro and prolongs pregnancy in women with early-onset preeclampsia. *J Pineal Res* [Internet]. 2018[cited 2023 Nov 11];65(3): e12508. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/jpi.12508> doi: 10.1111/jpi.12508
- Dou Y, Lin B, Cheng H, Wang C, Zhao M, Zhang J, et al. The reduction of melatonin levels is associated with the development of preeclampsia: a meta-analysis. *Hypertens Pregnancy*. 2019;38(2):65-72. doi: 10.1080/10641955.2019.1581215
- Chuffa LGA, Lupi LA, Cuciolo MS, Silveira HS, Reiter RJ, Seiva FRF. Melatonin Promotes Uterine and Placental Health: Potential Molecular Mechanisms. *Int J Mol Sci* [Internet]. 2019[cited 2023 Nov 11];21(1):300. Available from: <https://www.mdpi.com/1422-0067/21/1/300> doi: 10.3390/ijms21010300
- Huang B, Wang YK, Qin LY, Wei Q, Liu N, Jiang M, et al. A functional polymorphism rs10830963 in melatonin receptor 1B associated with the risk of gestational diabetes mellitus. *Biosci Rep* [Internet]. 2019[cited 2023 Nov 11];39(12): BSR20190744. Available from: <https://portlandpress.com/bioscirep/article-pdf/39/12/BSR20190744/864294/bsr-2019-0744.pdf> doi: 10.1042/BSR20190744
- Lim AJR, Indran IR, Kramer MS, Yong EL. Phenotypic spectrum of polycystic ovary syndrome and their relationship to the circadian biomarkers, melatonin and cortisol. *Endocrinol Diabetes Metab* [Internet]. 2019[cited 2023 Nov 11];2(3): e00047. Available from: <https://onlinelibrary.wiley.com/doi/epdf/10.1002/edm2.47> doi: 10.1002/edm2.47
- Cox B, Tsamou M, Vrijens K, Neven KY, Winckelmans E, de Kok TM, et al. A Co-expression Analysis of the Placental Transcriptome in Association With Maternal Pre-pregnancy BMI and Newborn Birth Weight. *Front Genet* [Internet]. 2019[cited 2023 Nov 11];10:354. Available from: <https://www.frontiersin.org/articles/10.3389/fgene.2019.00354/full> doi: 10.3389/fgene.2019.00354
- Bae H, Yang C, Lee JY, Park S, Bazer FW, Song G, Lim W. Melatonin improves uterine-conceptus interaction via regulation of SIRT1 during early pregnancy. *J Pineal Res* [Internet]. 2020[cited 2023 Nov 11];69(2): e12670. Available from: <https://onlinelibrary.wiley.com/doi/epdf/10.1111/jpi.12670> doi: 10.1111/jpi.12670
- Gomes PRL, Motta-Teixeira LC, Gallo CC, Carmo Buonfiglio DD, Camargo LS, Quintela T, et al. Maternal pineal melatonin in gestation and lactation physiology, and in fetal development and programming. *Gen Comp Endocrinol* [Internet]. 2021[cited 2023 Nov 11];300:113633. Available from: <https://www.sciencedirect.com/science/article/abs/pii/S0016648020303865?via%3Dihub> doi: 10.1016/j.ygcen.2020.113633
- Ejaz H, Figaro JK, Woolner AMF, Thottakam BMV, Galley HF. Maternal Serum Melatonin Increases During Pregnancy and Falls Immediately After Delivery Implicating the Placenta as a Major Source of Melatonin. *Front Endocrinol (Lausanne)* [Internet]. 2021[cited 2023 Nov 11];11:623038. Available from: <https://www.frontiersin.org/articles/10.3389/fendo.2020.623038/full> doi: 10.3389/fendo.2020.623038
- Wilkinson D, Shepherd E, Wallace EM. Melatonin for women in pregnancy for neuroprotection of the fetus. *Cochrane Database Syst Rev* [Internet]. 2016[cited 2023 Nov 11];3(3): CD010527. Available from: <https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD010527.pub2/full> doi: 10.1002/14651858.CD010527.pub2
- Langston-Cox A, Marshall SA, Lu D, Palmer KR, Wallace EM. Melatonin for the Management of Preeclampsia: A Review. *Antioxidants (Basel)* [Internet]. 2021[cited 2023 Nov 11];10(3):376. Available from: <https://www.mdpi.com/2076-3921/10/3/376> doi: 10.3390/antiox10030376
- Lee JY, Li S, Shin NE, Na Q, Dong J, Jia B, et al. Melatonin for prevention of placental malperfusion and fetal compromise associated with intrauterine inflammation-induced oxidative stress in a mouse model. *J Pineal Res* [Internet]. 2019[cited 2023 Nov 11];67(3): e12591. Available from: <https://onlinelibrary.wiley.com/doi/epdf/10.1111/jpi.12591> doi: 10.1111/jpi.12591
- Ma Q, Reiter RJ, Chen Y. Role of melatonin in controlling angiogenesis under physiological and pathological conditions. *Angiogenesis*. 2020;23(2):91-104. doi: 10.1007/s10456-019-09689-7
- Nagasawa Y, Nanami M, Kuragano T, Ishihara M. Melatonin and gestational hypertension. *Hypertens Res*. 2021;44(11):1540-2. doi: 10.1038/s41440-021-00722-1
- Sun Y, Wang C, Zhang N, Liu F. Melatonin ameliorates hypertension in hypertensive pregnant mice and suppresses the hypertension-induced decrease in Ca²⁺-activated K⁺ channels in uterine arteries. *Hypertens Res*. 2021;44(9):1079-86. doi: 10.1038/s41440-021-00675-5
- Thakor AS, Herrera EA, Serón-Ferré M, Giussani DA. Melatonin and vitamin C increase umbilical blood flow via nitric oxide-dependent mechanisms. *J Pineal Res*. 2010;49(4):399-406. doi: 10.1111/j.1600-079X.2010.00813.x
- Tordjman S, Chokron S, Delorme R, Charrier A, Bellissant E, Jaafari N, et al. Melatonin: Pharmacology, Functions and Therapeutic Benefits. *Curr Neuropharmacol*. 2017;15(3):434-43. doi: 10.2174/1570159X14666161228122115
- Mokhtari F, Akbari Asbagh F, Azmoodeh O, Bakhtiyari M, Almasi-Hashiani A. Effects of Melatonin Administration on Chemical Pregnancy Rates of Polycystic Ovary Syndrome Patients Undergoing Intrauterine Insemination: A Randomized Clinical Trial. *Int J Fertil Steril*. 2019;13(3):225-9. doi: 10.22074/ijfs.2019.5717
- Magnus MC, Wilcox AJ, Morken NH, Weinberg CR, Håberg SE. Role of maternal age and pregnancy history in risk of miscarriage: prospective register based study. *BMJ* [Internet]. 2019[cited 2023 Nov 11];364: 1869. Available from: <https://www.bmj.com/content/364/bmj.1869.long> doi: 10.1136/bmj.1869
- Fernando S, Wallace EM, Vollenhoven B, Lolatgis N, Hope N, Wong M, et al. Melatonin in Assisted Reproductive Technology: A Pilot Double-Blind Randomized Placebo-Controlled Clinical Trial. *Front Endocrinol (Lausanne)* [Internet]. 2018[cited 2023 Nov 11];9:545. Available from: <https://www.frontiersin.org/articles/10.3389/fendo.2018.00545/full> doi: 10.3389/fendo.2018.00545
- Hsu CN, Huang LT, Tain YL. Perinatal Use of Melatonin for Offspring Health: Focus on Cardiovascular and Neurological Diseases. *Int J Mol Sci* [Internet]. 2019[cited 2023 Nov 11];20(22):5681. Available from: <https://www.mdpi.com/1422-0067/20/22/5681> doi: 10.3390/ijms20225681

Prospects for further research. Translating research into practice.

Conflicts of interest. The authors declare no competing interests.

Funding sources. The article was published without financial support.

22. McCarthy R, Jungheim ES, Fay JC, Bates K, Herzog ED, England SK. Riding the Rhythm of Melatonin Through Pregnancy to Deliver on Time. *Front Endocrinol (Lausanne)* [Internet]. 2019[cited 2023 Nov 11];10:616. Available from: <https://www.frontiersin.org/articles/10.3389/fendo.2019.00616/full> doi: 10.3389/fendo.2019.00616
23. Mathiesen L, Bay-Richter C, Wegener G, Liebenberg N, Knudsen LE. Maternal stress and placental function; ex vivo placental perfusion studying cortisol, cortisone, tryptophan and serotonin. *PLoS One* [Internet]. 2020[cited 2023 Nov 11];15(6): e0233979. Available from: <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0233979> doi: 10.1371/journal.pone.0233979
24. Khan HL, Bhatti S, Abbas S, Kaloglu C, Qurat-Ul-Ain Zahra S, Khan YL, et al. Melatonin levels and microRNA (miRNA) relative expression profile in the follicular ambient microenvironment in patients undergoing in vitro fertilization process. *J Assist Reprod Genet.* 2021;38(2):443-59. doi: 10.1007/s10815-020-02010-2
25. Lee JY, Song H, Dash O, Park M, Shin NE, McLane MW, et al. Administration of melatonin for prevention of preterm birth and fetal brain injury associated with premature birth in a mouse model. *Am J Reprod Immunol* [Internet]. 2019[cited 2023 Nov 11];82(3): e13151. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/aji.13151> doi: 10.1111/aji.13151
26. Laste G, Silva AAD, Gheno BR, Rychcik PM. Relationship between melatonin and high-risk pregnancy: A review of investigations published between the years 2010 and 2020. *Chronobiol Int.* 2021;38(2):168-81. doi: 10.1080/07420528.2020.1863975

ФУНКЦІОНАЛЬНИЙ СТАН ШИШКОВИДНОГО ТІЛА У ЖІНОК ІЗ ЗАГРОЗЛИВИМ АБОРТОМ В ПЕРШОМУ ТРИМЕСТРІ – ПОДАЛЬШІ ПЕРСПЕКТИВИ ВАГІТНОСТІ

А. В. Пушкашу

Буковинський державний медичний університет
(м. Чернівці, Україна)

Резюме.

Мелатонін є важливим гормоном, який забезпечує нормальний розвиток вагітності, однак вплив серотоніну при цьому залишається недостатньо вивченим. Низька продукція мелатоніну може спричинити розвиток спонтанного абортів, так як він стимулює секрецію прогестерону та пригнічує синтез простагландинів. Мелатонін має вплив на індукцію та блокування апоптозу ворсинчастого цитотрофобласта, сприяє розвитку синцитіотрофобласту. Аналіз частоти позитивного перебігу вагітності показав суттєве значення впливу на перебіг вагітності та пологів, дотримання режиму сну та відпочинку також сприяють позитивному ефекту, проте загальної концепції, не існує, що робить тему актуальною.

Мета дослідження. Оцінити функціональний стан шишковидного тіла, зокрема кількість мелатоніну та серотоніну у крові вагітних у терміні 10-12 тижнів вагітності із загрозою переривання вагітності, яка проявлялася кров'янистими виділеннями чи формуванням ретрохоріальної гематоми та наслідки вагітності.

Матеріали та методи. Сформовано дві групи: I основна – 40 вагітних з загрозою переривання вагітності та формуванням ретрохоріальної гематоми, II контрольна – 44 здорових вагітних. Проведено акушерське дослідження, імуоферментне для визначення мелатоніну, серотоніну, хоріонічного гонадотропіну, УЗД.

Передбачено інформаційну згоду вагітних при відповідному роз'ясненні мети, завдань, методів та обсягу лабораторних та інструментальних методів дослідження.

Протокол дослідження схвалено Комісією з питань біомедичної етики Буковинського державного медичного університету, 2015 (протокол засідання комісії № 2 від 19.10.2023 року).

Дослідження виконувалися із дотриманням основних положень GCP (1996 рік), Конвенції Ради Європи про права людини та біомедицину (від 4 квітня 1997 р.), Гельсінської декларації Всесвітньої медичної асоціації про етичні принципи проведення наукових медичних досліджень за участю людини (1964-2008 рр.), наказу МОЗ України № 690 від 23.09.2009 р. (із змінами, внесеними згідно з Наказом Міністерства охорони здоров'я України № 523 від 12.07.2012).

Статистичний аналіз виконували за загальноприйнятими методами варіаційної статистики. Достовірність оцінювали за t-критерієм Стьюдента. Відмінності визнавали істотними при рівні значимості $p \leq 0,05$. Нами проведено також кореляційний аналіз системи Microsoft Excel, для визначення коефіцієнта кореляції застосували метод квадратів (метод Пірсона).

Робота є фрагментом НДР кафедри акушерства та гінекології, УДК 618.1-053.34 + 618.1-0536/.8]–07-084-08 «Профілактика, діагностика та лікування розладів перинатального періоду та репродуктивної системи жінок і дівчат-підлітків» (№ 201110Н, номер державної реєстрації 0111U006499. Термін виконання 02.2011-12.2015).

Результати дослідження. Згідно отриманих результатів ретрохоріальна гематома до 1 см виявляється втричі частіше, ніж більше 1 см, при цьому рівень мелатоніну не є достовірно нижчий, однак при ретрохоріальній гематомі, яка більше 1 см спостерігається достовірно зниження у випадках подальшого нормального перебігу вагітності та пологів. При подальшому ускладненому перебігу вагітності при ретрохоріальній гематомі до 1 см спостерігається значно виражене достовірно зниження мелатоніну ($p < 0,001$), на відміну від ретрохоріальної гематоми більше 1 см, де при ускладненому перебігу вагітності не виявлено достовірної різниці ($p > 0,05$). Кількість серотоніну є підвищеною у всіх підгрупах, незалежно від розмірів гематоми та подальшого перебігу вагітності.

Висновки. У випадку загрози переривання вагітності у першому триместрі з утворенням ретрохоріальної гематоми у подальшому зберігаються ризики ускладненого перебігу вагітності у 60,87 % випадків. При визначенні гормонів шишковидного тіла встановлено наявність достовірного збільшення серотоніну з одночасним достовірним зниженням мелатоніну зі збереженням позитивного взаємозв'язку між гормонами, що свідчить про порушення синтезу мелатоніну у ЦНС. Підтвердженням є нормальний рівень хоріонічного гонадотропіну та збереження життєздатності ембріонів. Оскільки встановлено порушення утворення гормонів у шишковидному тілі, дотримання рекомендацій щодо повноцінного відпочинку може позитивно вплинути на подальші результати вагітності та пологів.

Ключові слова: мелатонін; серотонін; ретрохоріальна гематома; загроза переривання вагітності.

Contact Information:

Anastasiia Pushkashu –postgraduate student of the Department of Obstetrics and Gynecology, Bukovinian State Medical University (Chernivtsi, Ukraine).

Contact Address: Teatralna Square, 2, Chernivtsi, 58000, Ukraine.

Contact Phone: +380509148333

e-mail: anastasiia.pushkashu@gmail.com

Відомості про автора:

Пушкашу Анастасія Василівна – аспірант кафедри акушерства та гінекології, Буковинський державний медичний університет (м. Чернівці, Україна).

Контактна адреса: площа Театральна, 2, м. Чернівці, 58002, Україна.

Контактний телефон: +380509148333

e-mail: anastasiia.pushkashu@gmail.com



Received for editorial office on 23/07/2023

Signed for printing on 10/10/2023
