

## On modular representations of semigroups $S_p \times T_p$

Vitaliy M. Bondarenko, Elina M. Kostyshyn

Communicated by V. V. Kirichenko

**ABSTRACT.** Let  $p$  be simple, and let  $S_p$  and  $T_p$  be the symmetric group and the symmetric semigroup of degree  $p$ , respectively. The theorem of this paper says that the direct product  $S_p \times T_p$  are of wild representation type over any field of characteristic  $p$ . The main case is  $p = 2$ .

Let  $k$  be a field. A semigroup is called *of tame representation type* (resp. *of wild representation type*) over  $k$  if so is the problem of classifying its representations (see precise general definitions in [1]).

We give the precise definition of semigroups of wild representation type in matrix language.

For a semigroup  $S$  and a  $k$ -algebra  $\Lambda$ , we denote by  $R_\Lambda(S)$  the set of all matrix representations of  $S$  over  $\Lambda$ ;  $R_k(\Lambda)$  denotes the category of matrix representations of  $\Lambda$  over  $k$ .

A semigroup  $S$  is called of wild representation type (or simply wild) over  $k$  if there exists a matrix representation  $M$  of  $S$  over  $\Lambda = K_2 = k \langle x, y \rangle$  such that the following conditions hold:

- 1) the matrix representation  $M \otimes X$  (of  $S$  over  $k$ ) with  $X \in R_k(\Lambda)$  is indecomposable if so is  $X$ ;
- 2) the matrix representations  $M \otimes X$  and  $M \otimes X'$  are nonequivalent if so are  $X$  and  $X'$ .

---

**2010 MSC:** 16G, 20M30.

**Key words and phrases:** matrix, wild, transformation, symmetric semigroup, modular representations.

Here  $K_2 = k \langle x, y \rangle$  denotes the free associative  $k$ -algebra in two noncommuting variables  $x$  and  $y$ .

We call such an  $M$  a *perfect representation of  $S$  over  $\Lambda$* .

In practice, to simplify the proofs of wildness (not only semigroup but also other objects) one can replace  $K_2$  by any wild  $k$ -algebra.

The main result of this paper is the following theorem.

**Theorem.** *Let  $k$  be a field of characteristic  $p \neq 0$  and let  $S_p$  and  $T_p$  be the symmetric group and the symmetric semigroup of degree  $p$ , respectively. Then the semigroup  $S_p \times T_p$  is wild over  $k$ .*

Here  $\times$  denotes, as usual, the sign of the direct product.

Note that  $T_p$  and  $S_p \times T_p$  are monoids.

Since the factor semigroup of  $T_p$  by its only maximal two-sided ideal (generated by all the non-invertible elements) is isomorphic to  $S_p$ , the semigroup  $S_p \times T_p$  is wild for  $p \neq 2$  by the criterion of tameness and wildness of finite groups [2]. In case  $p = 2$  we will indicate a perfect representation of  $S_p \times T_p$  over the  $k$ -algebra  $\Lambda = k\Gamma$  of paths of the quiver  $\Gamma$  with two vertices  $p_1, p_2$  and two arrows  $x : p_1 \rightarrow p_1, y : p_1 \rightarrow p_2$  (this quiver is wild [3, 4]).

The monoid  $T_2$  of transformations of the set  $\{1, 2\}$  is generated by the elements  $a, b$ , where  $a(1) = 2, a(2) = 1, b(1) = 2, b(2) = 2$ , with defining relations  $a^2 = 1, b^2 = b, ab = b$  [5]. Obviously that the monoid  $S_2 \times T_2$  is generated by the elements  $g, a, b$  with the additional relations  $g^2 = 1, ga = ag, gb = bg$  ( $g$  denotes the non-identity element of  $S_2$ ).

Consider the next matrix representation  $\gamma$  of  $S_2 \times T_2$  over the algebra  $\Lambda = k\Gamma$ :

$$\gamma(g) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & y & 1 & x \\ 0 & 0 & 0 & 1 \end{pmatrix}, \gamma(a) = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \gamma(b) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

( $\gamma(1)$  is equal to the identity matrix).

We will prove that  $\gamma$  is a perfect representation.

Let  $\varphi, \varphi'$  be representations of  $\Lambda$  over  $k$  having the same dimension  $s$  and let  $G = (\gamma \otimes \varphi)(g), A = (\gamma \otimes \varphi)(a), B = (\gamma \otimes \varphi)(b), G' = (\gamma \otimes \varphi')(g), A' = (\gamma \otimes \varphi')(a), B' = (\gamma \otimes \varphi')(b)$ . Consider the matrix equalities (in the variable  $X$ )

$$GX = XG', \quad AX = XA', \quad BX = XB', \quad (*)$$

viewing all their matrices as  $s \times s$  block ones.

The equalities (of the  $s \times s$   $ij$ -blocks)

$$(GX)_{ij} = (XG')_{ij}, \quad (AX)_{ij} = (XA')_{ij}, \quad (BX)_{ij} = (XB')_{ij},$$

$i, j \in \{1, 2, 3, 4\}$  are denoted by  $(1; ij)$ ,  $(2; ij)$ ,  $(3; ij)$ , respectively.

We first write down all equalities of the forms  $(2; ij)$  and  $(3; ij)$  besides the trivial identities  $0 = 0$  and  $X_{ii} = X_{ii}$ :

$$\begin{aligned} (2; 1, 1) : X_{21} &= 0, & (2; 1, 2) : X_{22} &= X_{11}, & (2; 1, 3) : X_{23} &= 0, \\ (2; 1, 4) : X_{24} &= X_{13}, & (2; 2, 2) : 0 &= X_{21}, & (2; 2, 4) : 0 &= X_{23}, \\ (2; 3, 1) : X_{41} &= 0, & (2; 3, 2) : X_{42} &= X_{31}, & (2; 3, 3) : X_{43} &= 0, \\ (2; 3, 4) : X_{44} &= X_{33}, & (2; 4, 2) : 0 &= X_{41}, & (2; 4, 4) : 0 &= X_{43}, \\ (3; 1, 2) : X_{12} &= 0, & (3; 1, 3) : X_{13} &= 0, & (3; 1, 4) : X_{14} &= 0, \\ (3; 2, 1) : 0 &= X_{21}, & (3; 3, 1) : 0 &= X_{31}, & (3; 4, 1) : 0 &= X_{41}. \end{aligned}$$

From these equalities it follows that

$$X = \begin{pmatrix} X_{11} & 0 & 0 & 0 \\ 0 & X_{11} & 0 & 0 \\ 0 & X_{32} & X_{33} & X_{34} \\ 0 & 0 & 0 & X_{33} \end{pmatrix}.$$

Then from the equalities

$$(1; 3, 2) : \varphi(y)X_{11} = X_{33}\varphi'(y), \quad (1; 3, 4) : \varphi(x)X_{33} = X_{33}\varphi'(x) \quad (**)$$

(the only two nontrivial equalities of the form  $(1; ij)$  modulo the equalities  $(2; ij)$  and  $(3; ij)$ ) we have that the matrix  $k$ -representations  $\varphi$  and  $\varphi'$  of  $\Lambda = k\Gamma$  are equivalent if so are the matrix  $k$ -representations  $\gamma \otimes \varphi$  and  $\gamma \otimes \varphi'$  of  $S_2 \times T_2$  (because  $X_{11}$  and  $X_{33}$  are invertible if so is  $X$ ).

Thus, for the representation  $\gamma$  condition 2) of the definition of wild semigroups holds.

From the form of the matrix  $X$  it follows that the endomorphism algebra of  $\gamma \otimes \varphi$  is local if and only if so is the endomorphism algebra of  $\varphi$  (these algebras are defined, respectively, by  $(*)$  and  $(**)$  with  $\varphi = \varphi'$ ). Therefore  $\gamma \otimes \varphi$  is indecomposable if  $\varphi$  is indecomposable, and consequently  $\gamma$  satisfies condition 1) of the mentioned definition too.

The theorem is proved.

Because as a perfect matrix representation of the quiver  $\Gamma$  over the algebra  $K'_2 = k \langle x', y' \rangle$  one can take the representation

$$x \rightarrow \begin{pmatrix} 0 & x' \\ 1 & y' \end{pmatrix}, \quad y \rightarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix},$$

it follows from the proof of our theorem that the following representation  $\lambda$  of the semigroup  $S_2 \times T_2$  over  $K'_2$  is perfect:

$$\lambda(g) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & x' \\ 0 & 0 & 0 & 1 & 1 & y' \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad \lambda(a) = \begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$\lambda(b) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

### References

- [1] Yu. A. Drozd, *Tame and wild matrix problems*, Lecture Notes in Math. **832** (1980), pp. 242-258.
- [2] V. M. Bondarenko, Ju. A. Drozd, *Representation type of finite groups*, Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) **71** (1977), pp. 24-42 (in Russian); English trans. in J. Soviet Math. **20** (1982), pp. 2515-2528.
- [3] L. A. Nazarova, *Representations of quivers of infinite type*, Izv. Akad. Nauk SSSR Ser. Mat. **37** (1973), pp. 752-791 (in Russian); English trans. in Math. USSR-Izv. **7** (1973), pp. 749-792.
- [4] P. Donovan, M.-R. Freislich, *The representation theory of finite graphs and associated algebras*, Carleton Math. Lecture Notes, No. 5. Carleton University, Ottawa, Ont., 1973, 83 pp.
- [5] V. M. Bondarenko, E. M. Kostyshyn, *Modular representations of the semigroup  $T_2$* , Nauk. Visn. Uzhgorod. Univ., Ser. Mat. Inform. **22** (2011), pp. 26-34 (in Ukrainian).

### CONTACT INFORMATION

**V. M. Bondarenko** Institute of Mathematics, NAS, Kyiv, Ukraine  
*E-Mail:* vitalij.bond@gmail.com

**E. M. Kostyshyn** Department of Mechanics and Mathematics,  
 Kyiv National Taras Shevchenko Univ.,  
 Volodymyrska str., 64, 01033 Kyiv, Ukraine  
*E-Mail:* elina.kostyshyn@mail.ru

Received by the editors: 17.07.2013  
 and in final form 17.07.2013.