

615.453.2:661.183.1:638.17

- ... M
- ... yAHM



[8, 9].

[7].

) [3, 4].

[6, 7].

[10].

35%

20%

10%,
s ulus §4.

[2, 5, 9].

s ulus §3

§4

- 10%

[6].

37 °. 30 6 , 12

in vitro , 24 .

in vitro, -

[1, 4, 10].

in vitro . 0,2% (

4 : 10 % () .

0,5-1 10% 3



8 , 45°

0,2 , 1, . 1.

10% 1

	0,5	1	1,5	2	2,5			5		12
1	2	5	12	17	20	25		25		25
2			14	18	22	2		28		28

*



)

)

. I.

:)

)

1.

s ulus (

2.

s ulus:

—

—

10%

(), 1%

—

—

-

;

—

—

-

—

-

—

-

—

-

—

-

—

-

—

-

—

-

—

-

—

-

—

-

—

-

—

-

—

-

—

-

(), - 10% - 10% -

3.) -

§4, 10%

1. /

" - I- : , 2000. - 556 .

2. B.A. //PM . - 2002. - . 10. - 26.

3. -M.: M

- 2000. - 976 .

4. AM., B.A., M . .

Aesculus. // - 1999. - 15. - . 24-28.

5. H.A. // - 2003. - 22. - 16-18.

6. B. / . B. .P . - M.: . . , 1967. - 373 .

7. A.B., A.I. // - 1999. - 12. - . 63-68.

. Bassler D, Okpanyi S, Schrodter A, et al. Bioavailability of beta-aescin from horse chestnut seed extract: comparative clinical studies of two Galenic formulations. //Adv Ther. - 2003, Sep-Oct. - . 20, 5 - . 295-304.

9. Bielanski , Piotrowski ZH. Horse-chestnut seed extract for chronic venous insufficiency. //J Farm Pract. - 1999 Mar. - . 48, 3 - . 171-172.

10. Potselueva L.A., and Baranova A.A. Biofarmaceutical method of study of fermental preparations in experiments in vitro /- Hersteller: Agens - Werk ejeuer + Reisser, 1051 Wien // a. - . 63, 3 - 2001. - 255 p.

615.453.2:661.183.1:638.17(001.8)

in vitro

A s ulus.

10%

§4

. . M

in vitro
A s ulus.

10%

§4

r v , V.A. Soboleva

**BIOPHARMACEUTICAL RESEARCHES AND QUALITATIVE ANALYSIS OF A HOMEOPATHIC DRUG OF HORSE-
CHESNUT.**

Key words: Horse-chestnut, homeopathic drugs, biopharmaceutical researches, phenolic bonds

The lead biopharmaceutical researches of rate and completeness release of reacting materials in experiences in vitro with use of a method of agar plates. Fixed, that the addition aeroforce for rising flowability of a drug does not variate parameters of release of medicinal materials. The quantification of contents of phenolic bonds of Propolisum in a drug on pharmaceutical procedure with the help of tool methods of analysis is lead.

« : / . . . - : , 2009. - 480 .»