

УДК 517.5

KHATS' R.V.

ASYMPTOTIC BEHAVIOR OF ENTIRE FUNCTIONS OF IMPROVED REGULAR GROWTH IN THE METRIC OF $L^p[0, 2\pi]$

Khats' R.V. *Asymptotic behavior of entire functions of improved regular growth in the metric of $L^p[0, 2\pi]$* . Carpathian Mathematical Publications 2013, 5 (2), 341–344.

We describe an asymptotic behavior of entire functions of improved regular growth with zeros on a finite system of rays in the metric of $L^p[0, 2\pi]$.

Key words and phrases: entire function of improved regular growth, finite system of rays, Fourier coefficients, Hausdorff-Young theorem.

Drohobych Ivan Franko State Pedagogical University, Drohobych, Ukraine

E-mail: khats@ukr.net

1 INTRODUCTION AND MAIN RESULT

The asymptotic behavior of entire and meromorphic functions of positive order of completely regular growth (for details, see [2, 3, 5, 11]) in the metric of $L^p[0, 2\pi]$ was described in [6, 11, 12, 13]. Similar results for entire functions of zero order whose zero-counting functions are slowly increasing were obtained in [1]. In particular, from [11, Theorem 7.2, p. 78] it follows the following statement.

Theorem A. *If an entire function f of order $\rho \in (0, +\infty)$ with the indicator $h(\theta, f)$ is of completely regular growth, then for any $p \in [1, +\infty)$*

$$\lim_{r \rightarrow +\infty} \left\{ \frac{1}{2\pi} \int_0^{2\pi} \left| \frac{\log |f(re^{i\theta})|}{r^\rho} - h(\theta, f) \right|^p d\theta \right\}^{1/p} = 0.$$

Conversely, if for some entire function f , $f(0) = 1$, there exist $p \in [1, +\infty)$ and $\tilde{h} \in L^p[0, 2\pi]$ such that

$$\lim_{r \rightarrow +\infty} \left\{ \frac{1}{2\pi} \int_0^{2\pi} \left| \frac{\log |f(re^{i\theta})|}{r^\rho} - \tilde{h}(\theta) \right|^p d\theta \right\}^{1/p} = 0,$$

then f is of completely regular growth and $\tilde{h}(\theta) = h(\theta, f)$ for almost all $\theta \in [0, 2\pi]$.

The aim of the present paper is obtain an analog of Theorem A for entire functions of improved regular growth with zeros on a finite system of rays (see [4, 7, 8, 9, 10, 14]).

An entire function f is called a function of *improved regular growth* (see [7, 8, 9, 10, 14]) if for some $\rho \in (0, +\infty)$ and $\rho_1 \in (0, \rho)$, and a 2π -periodic ρ -trigonometrically convex function $h(\varphi) \not\equiv -\infty$ there exists the set $U \subset \mathbb{C}$ contained in the union of disks with finite sum of radii and such that $\log |f(z)| = |z|^\rho h(\varphi) + o(|z|^{\rho_1})$, $U \ni z = re^{i\varphi} \rightarrow \infty$.

If an entire function f is a function of improved regular growth, then it has the order ρ and indicator h [14]. Our main result is the following theorem.

2010 *Mathematics Subject Classification:* 30D15.

Theorem 1. *An entire function f of order $\rho \in (0, +\infty)$ with zeros on a finite system of rays $\{z : \arg z = \psi_j\}, j \in \{1, \dots, m\}, 0 \leq \psi_1 < \psi_2 < \dots < \psi_m < 2\pi$, is a function of improved regular growth if and only if for some $\rho_2 \in (0, \rho)$ and any $p \in [1, +\infty)$, one has*

$$\left\{ \frac{1}{2\pi} \int_0^{2\pi} \left| \frac{\log |f(re^{i\varphi})|}{r^\rho} - h(\varphi) \right|^p d\varphi \right\}^{1/p} = o(r^{\rho_2 - \rho}), \quad r \rightarrow +\infty. \tag{1}$$

2 PRELIMINARIES

Let f be an entire function with $f(0) = 1$, let $(\lambda_n)_{n \in \mathbb{N}}$ be the sequence of its zeros, let p be the smallest integer for which $\sum_{n=1}^\infty |\lambda_n|^{-p-1} < +\infty$, let Q_ρ be the coefficient of z^ρ in the exponential factor in the Hadamard-Borel representation [5, p. 38] of an entire function of finite order, and let $c_k(r, \log |f|)$ be the Fourier coefficients of $\log |f|$, i.e.

$$c_k(r, \log |f|) := \frac{1}{2\pi} \int_0^{2\pi} e^{-ik\varphi} \log |f(re^{i\varphi})| d\varphi, \quad k \in \mathbb{Z}, \quad r > 0.$$

Further, let f be an entire function of order $\rho \in (0, +\infty)$ with zeros on a finite system of rays $\{z : \arg z = \psi_j\}, j \in \{1, \dots, m\}, 0 \leq \psi_1 < \psi_2 < \dots < \psi_m < 2\pi$. Furthermore, if ρ is noninteger and f is a function of improved regular growth, then an indicator h of f has the form ([8, 10, 14])

$$h(\varphi) = \sum_{j=1}^m h_j(\varphi),$$

where $h_j(\varphi)$ is a 2π -periodic function such that on $[\psi_j, \psi_j + 2\pi)$

$$h_j(\varphi) = \frac{\pi \Delta_j}{\sin \pi \rho} \cos \rho(\varphi - \psi_j - \pi), \quad \Delta_j \in [0, +\infty).$$

In the case $\rho \in \mathbb{N}$, the indicator h is defined by the formula ([8, 9, 10])

$$h(\varphi) = \begin{cases} \tau_f \cos(\rho\varphi + \theta_f) + \sum_{j=1}^m h_j(\varphi), & p = \rho, \\ Q_\rho \cos \rho\varphi, & p = \rho - 1, \end{cases}$$

where $\delta_f \in \mathbb{C}, \tau_f = |\delta_f/\rho + Q_\rho|, \theta_f = \arg(\delta_f/\rho + Q_\rho)$ and $h_j(\varphi)$ is a 2π -periodic function such that on $[\psi_j, \psi_j + 2\pi)$ we have

$$h_j(\varphi) = \Delta_j(\pi - \varphi + \psi_j) \sin \rho(\varphi - \psi_j) - \frac{\Delta_j}{\rho} \cos \rho(\varphi - \psi_j).$$

Lemma 1 ([8]). *If an entire function f of order $\rho \in (0, +\infty)$ with zeros on a finite system of rays $\{z : \arg z = \psi_j\}, j \in \{1, \dots, m\}, 0 \leq \psi_1 < \psi_2 < \dots < \psi_m < 2\pi$, is of improved regular growth, then for some $\rho_3 \in (0, \rho)$*

$$c_k(r, \log |f|) = c_k r^\rho + \frac{o(r^{\rho_3})}{k^2 + 1}, \quad r \rightarrow +\infty, \tag{2}$$

holds uniformly in $k \in \mathbb{Z}$, where

$$c_k := \frac{1}{2\pi} \int_0^{2\pi} e^{-ik\varphi} h(\varphi) d\varphi = \frac{\rho}{\rho^2 - k^2} \sum_{j=1}^m \Delta_j e^{-ik\psi_j}, \quad \Delta_j \in [0, +\infty),$$

for a noninteger ρ , and

$$c_k = \begin{cases} \frac{\rho}{\rho^2 - k^2} \sum_{j=1}^m \Delta_j e^{-ik\psi_j}, & |k| \neq \rho = p, \\ \frac{\tau_f e^{i\theta_f}}{2} - \frac{1}{4\rho} \sum_{j=1}^m \Delta_j e^{-i\rho\psi_j}, & k = \rho = p, \\ 0, & |k| \neq \rho = p + 1, \\ \frac{Q_\rho}{2}, & k = \rho = p + 1, \end{cases}$$

if ρ is an integer.

Remark that, using Lemma 1 and the Riesz-Fischer theorem [11, p. 5], we get that there exists an indicator $h \in L^2[0, 2\pi]$ defined by the equality $h(\varphi) := \sum_{k \in \mathbb{Z}} c_k e^{ik\varphi}$ (see [11, Definition 7.2, p. 77]).

3 PROOF OF THEOREM 1

Necessity. If an entire function f of order $\rho \in (0, +\infty)$ with zeros on a finite system of rays $\{z : \arg z = \psi_j\}$, $j \in \{1, \dots, m\}$, $0 \leq \psi_1 < \psi_2 < \dots < \psi_m < 2\pi$, is of improved regular growth, then by (2), we have

$$\left| \frac{c_k(r, \log |f|)}{r^\rho} - c_k \right| \leq \frac{C}{k^2 + 1}, \quad k \in \mathbb{Z}, \quad (3)$$

for some constant $C > 0$ and all $r > 0$. In view of this, the sequence $(r^{-\rho} c_k(r, \log |f|) - c_k)_{k \in \mathbb{Z}}$ belongs to the space l_q for all $q > 1$ and $r > 0$. Moreover, applying the Hausdorff-Young theorem [11, p. 5], for $p \geq 2$, $p^{-1} + q^{-1} = 1$, we get

$$\left\{ \frac{1}{2\pi} \int_0^{2\pi} \left| \frac{\log |f(re^{i\varphi})|}{r^\rho} - h(\varphi) \right|^p d\varphi \right\}^{1/p} \leq \left\{ \sum_{k \in \mathbb{Z}} \left| \frac{c_k(r, \log |f|)}{r^\rho} - c_k \right|^q \right\}^{1/q}.$$

According to (3), the last series is uniformly convergent with respect to r . Therefore, using Lemma 1, we obtain (1) for $p \geq 2$. From this and Hölder's inequality it follows that (1) holds for $1 \leq p < 2$.

Sufficiency. Let (1) holds. Then for some $\rho_2 \in (0, \rho)$ and each $k \in \mathbb{Z}$

$$\begin{aligned} \left| \frac{c_k(r, \log |f|)}{r^\rho} - c_k \right| &\leq \frac{1}{2\pi} \int_0^{2\pi} \left| \frac{\log |f(re^{i\varphi})|}{r^\rho} - h(\varphi) \right| d\varphi \\ &\leq \left\{ \frac{1}{2\pi} \int_0^{2\pi} \left| \frac{\log |f(re^{i\varphi})|}{r^\rho} - h(\varphi) \right|^p d\varphi \right\}^{1/p} = o(r^{\rho_2 - \rho}), \quad r \rightarrow +\infty. \end{aligned}$$

Hence, for some $\rho_2 \in (0, \rho)$ and each $k \in \mathbb{Z}$ we get $c_k(r, \log |f|) = c_k r^\rho + o(r^{\rho_2})$, $r \rightarrow +\infty$. Then by [10, Theorem 1, p. 1718] an entire function f is a function of improved regular growth. This concludes the proof of the theorem.

REFERENCES

- [1] Bodnar O.V., Zabolots'kyi M.V. *Criteria for the regularity of growth of the logarithm of modulus and the argument of an entire function*. Ukrainian Math. J. 2010, **62** (7), 1028–1039. doi:10.1007/s11253-010-0411-x (translation of Ukr. Mat. Zh. 2010, **62** (7), 885–893. (in Ukrainian))
- [2] Gol'dberg A.A. *B.Ya. Levin, the founder of the theory of entire functions of completely regular growth*. Mat. Fiz., Anal., Geom. 1994, **1** (2), 186–192. (in Russian)
- [3] Gol'dberg A.A., Levin B.Ya., Ostrovskii I.V. *Entire and meromorphic functions*. VINITI 1991, **85**, 5–186. (in Russian)
- [4] Hirnyk M.O. *Subharmonic functions of improved regular growth*. Dopov. Nats. Akad. Nauk Ukr. 2009, **4**, 13–18. (in Ukrainian)
- [5] Levin B.Ya. *Distribution of zeros of entire functions*. Gostekhizdat, Moscow, 1956. (in Russian)
- [6] Kalynets' R.Z., Kondratyuk A.A. *On the regularity of the growth of the modulus and argument of an entire function in the metric of $L^p[0, 2\pi]$* . Ukrainian Math. J. 1998, **50** (7), 1009–1018. doi: 10.1007/BF02528830 (translation of Ukr. Mat. Zh. 1998, **50** (7), 889–896. (in Ukrainian))
- [7] Khats' R.V. *Asymptotic behavior of averaging of entire functions of improved regular growth*. Carpathian Math. Publ. 2013, **5** (1), 129–133.
- [8] Khats' R.V. *Averaging of entire functions of improved regular growth with zeros on a finite system of rays*. Visn. Nats. Univ. L'viv. Politekh., Fiz.-Mat. Nauky. 2011, **718** (718), 10–14.
- [9] Khats' R.V. *On entire functions of improved regular growth of integer order with zeros on a finite system of rays*. Mat. Stud. 2006, **26** (1), 17–24.
- [10] Khats' R.V. *Regularity of growth of Fourier coefficients of entire functions of improved regular growth*. Ukrainian Math. J. 2012, **63** (12), 1953–1960. doi:10.1007/s11253-012-0624-2 (translation of Ukr. Mat. Zh. 2011, **63** (12), 1717–1723. (in Ukrainian))
- [11] Kondratyuk A.A. *Fourier series and meromorphic functions*. Vyshcha shkola, Kyiv, 1988. (in Russian)
- [12] Vasyl'kiv Ya.V. *Asymptotic behavior of logarithmic derivatives and logarithms of meromorphic functions of completely regular growth in the metric of $L^p[0, 2\pi]$. I*. Mat. Stud. 1999, **12** (1), 37–58. (in Ukrainian)
- [13] Vasyl'kiv Ya.V. *Asymptotic behavior of logarithmic derivatives and logarithms of meromorphic functions of completely regular growth in the metric of $L^p[0, 2\pi]$. II*. Mat. Stud. 1999, **12** (2), 135–144. (in Ukrainian)
- [14] Vynnyts'kyi B.V., Khats' R.V. *On growth regularity of entire function of noninteger order with zeros on a finite system of rays*. Mat. Stud. 2005, **24** (1), 31–38. (in Ukrainian)

Received 21.08.2013

Хаць Р.В. Асимптотична поведінка цілих функцій покращеного регулярного зростання в $L^p[0, 2\pi]$ -метриці // Карпатські математичні публікації. — 2013. — Т.5, №2. — С. 341–344.

Описано асимптотичну поведінку цілих функцій покращеного регулярного зростання з нулями на скінченній системі променів в $L^p[0, 2\pi]$ -метриці.

Ключові слова і фрази: ціла функція покращеного регулярного зростання, скінченна система променів, коефіцієнти Фур'є, теорема Гаусдорфа-Юнга.

Хаць Р.В. Асимптотическое поведение целых функций улучшенного регулярного роста в $L^p[0, 2\pi]$ -метрике // Карпатские математические публикации. — 2013. — Т.5, №2. — С. 341–344.

Описано асимптотическое поведение целых функций улучшенного регулярного роста с нулями на конечной системе лучей в $L^p[0, 2\pi]$ -метрике.

Ключевые слова и фразы: целая функция улучшенного регулярного роста, конечная система лучей, коэффициенты Фурье, теорема Гаусдорфа-Юнга.