



## CONDITIONS OF SEMISCALAR EQUIVALENCE OF POLYNOMIAL 2-BY-2 MATRICES

B.Z. SHAVAROVSKII<sup>1</sup>

<sup>1</sup>*Pidstryhach Institute for Applied Problems of Mechanics and Mathematics, 3-b,  
Naukova Str., L'viv, Ukraine.*

---

B.Z. Shavarovskii, *Conditions of semiscalar equivalence of polynomial 2-by-2 matrices*, Math. Bull. Shevchenko Sci. Soc. **14** (2017) 7–28.

We consider the problem of determining whether two polynomial matrices are semiscalarly equivalent. Large difficulties in this problem arise already for matrices of second order. In this connection under certain restrictions the necessary and sufficient conditions of semiscalar equivalence of 2-by-2 polynomial matrices are found.

Б.З. Шаваровський. *Умови напівскалярної еквівалентності поліноміальних  $2 \times 2$ -матриць* // Мат. вісник НТШ, **14** (2017) 7–28.

Досліджується проблема визначення, коли дві поліноміальні матриці є напівскалярно еквівалентними. Значні труднощі у цій проблемі виникають вже для матриць другого порядку. У зв'язку з цим при певних обмеженнях вказані необхідні та достатні умови напівскалярної еквівалентності  $2 \times 2$  поліноміальних матриць.

---

### Introduction

The notion of semiscalar equivalence of matrices was introduced to algebra by P.S. Kazimirskii and V.M. Petrychkovych in 1977. By definition [1], two polynomial matrices are called semiscalarly equivalent if they can be transformed one to the other by multiplying from the left by a nonsingular numerical matrix and from the right by

---

2010 *Mathematics Subject Classification*: 15A21

УДК: 512.64

*Key words and phrases*: semiscalar equivalence, polynomial matrix, invariant.

*E-mail*: bshavarovskii@gmail.com

an invertible polynomial matrix. The notion of *PS*-equivalence of matrices was introduced by J.A. Dias da Silva and T.J. Laffey in 1999. By definition [2], polynomial matrices  $A(x)$  and  $B(x)$  are called *PS*-equivalent provided  $B(x) = P(x)A(x)Q$  for some invertible matrices  $Q$  – independent of  $x$  and  $P(x)$  – dependent on  $x$ . We consider the problem of determining whether two matrices are semiscalarly equivalent. It contains the classical linear algebra problem of reducing a pair of numerical matrices to a canonical form by simultaneous similarity transformations. This problem has not been extensively studied in the literature. Large difficulties in this problem arise already for matrices of second order.

In this paper the semiscalar equivalence for one class of polynomial matrices of second order is investigated. In particular, necessary and sufficient conditions are found under which two 2-by-2 matrices over the ring  $\mathbb{C}[x]$  are semiscalarly equivalent.

We consider the ring  $M(2, \mathbb{C}[x])$  of order two polynomial matrices over the field of complex numbers  $\mathbb{C}$ . Let  $A(x) \in M(2, \mathbb{C}[x])$ . Suppose that the rank of matrix  $A(x)$  is full. According to [1] (see also [3], Section IV, § 1 or [4]) the matrix  $A(x)$  is semiscalarly equivalent to a matrix of lower triangular form with invariant multipliers on the main diagonal. Without loss of generality, we can assume that the first invariant multiplier of the considered matrix is identity. Therefore, this matrix can be considered in the form

$$A(x) = \left\| \begin{array}{cc} 1 & 0 \\ a(x) & \delta(x) \end{array} \right\|, \quad 0 < \deg a(x) < \deg \delta(x) \quad (1)$$

since the case with  $a(x) \equiv 0$  is trivial.

In this paper we use the standard notations. In particular,  $a^{(t)}(\alpha)$  is the value at  $x = \alpha$  of the  $t$ -th derivative of the polynomial  $a(x) \in \mathbb{C}[x]$  for  $t \in \mathbb{N}$ . Further, by the symbol  $\bar{0}$  we denote the zero column of arbitrary height. The transpose operation is denoted by the symbol  $T$ . Let  $M(2, \mathbb{C})$ ,  $M(2, \mathbb{C}[x])$  denote the rings of  $2 \times 2$  matrices over  $\mathbb{C}$ ,  $\mathbb{C}[x]$ , respectively, and let  $GL(2, \mathbb{C})$ ,  $GL(2, \mathbb{C}[x])$  be their groups of units, respectively.

The determinant  $|A(x)|$  is called the characteristic polynomial of  $A(x)$  and its roots are called the characteristic roots of matrix  $A(x)$ . In this article the characteristic roots of a matrix can be of arbitrary multiplicity. The question of finding a complete set of invariants for *PS*-equivalence in the case 2-by-2 matrices without multiple characteristic roots is discussed in [2]. A different case of semiscalar equivalence of polynomial matrices of second order is considered in [5].

## 1. Main results

Let us denote by  $M$  the set of characteristic roots of a matrix  $A(x)$  of the form (1). It is clear that the set  $M$  is an invariant of the matrix  $A(x)$  with respect to semiscalar equivalence.

On the set  $M$  consider the equivalence relation

$$E = \{(\alpha, \beta) \in M \times M : a(\alpha) = a(\beta)\},$$

which determined a partition

$$M = \{E(\alpha) : \alpha \in M\} \quad (2)$$

of  $M$  into the equivalence classes  $E(\alpha) := \{\beta \in M : (\alpha, \beta) \in E\}$ .

The following two assertions yield some invariants of the matrix  $A(x)$  with respect to semiscalar equivalence.

**Proposition 1.1.** *The partition (2) of the set  $M$  of characteristic roots of matrix  $A(x)$  of the form (1) is invariant for the class  $\{CA(x)Q(x)\}$  of semiscalarly equivalent matrices.*

*Proof.* Let the matrices  $A(x)$  (1) and

$$B(x) = \begin{vmatrix} 1 & 0 \\ b(x) & \Delta(x) \end{vmatrix}, \quad \deg b(x) < \deg \Delta(x), \quad (3)$$

be semiscalarly equivalent, i.e.

$$SA(x)R(x) = B(x), \quad (4)$$

where  $S \in GL(2, \mathbb{C})$  and  $R(x) \in GL(2, \mathbb{C}[x])$ . Introducing the notations  $S = \|s_{ij}\|_1^2$ ,  $R^{-1}(x) = \|r_{ij}(x)\|_1^2$ , we deduce from (4) the relations

$$s_{11} + s_{12}a(x) = r_{11}(x), \quad (5)$$

$$s_{12}\delta(x) = r_{12}(x), \quad (6)$$

$$s_{21} + s_{22}a(x) = b(x)r_{11}(x) + \delta(x)r_{21}(x). \quad (7)$$

Setting  $x = \alpha$  and  $x = \beta$ ,  $\alpha \neq \beta$ ,  $\alpha, \beta \in M$ , we obtain the equalities

$$s_{21} + s_{22}a(\alpha) = b(\alpha)r_{11}(\alpha) \quad \text{and} \quad s_{21} + s_{22}a(\beta) = b(\beta)r_{11}(\beta).$$

If  $a(\alpha) = a(\beta)$ , then left sides of the resulting relations are equal. Therefore, from the equality of right sides, taking into account  $r_{11}(\alpha) = r_{11}(\beta) \neq 0$  (see (5), (6)), we have  $b(\alpha) = b(\beta)$ . The notion of semiscalar equivalence is a symmetric relation. Then from  $b(\alpha) = b(\beta)$  a similar argument yields  $a(\alpha) = a(\beta)$ . This completes the proof.  $\square$

**Proposition 1.2.** *Let  $\alpha$  be a characteristic root of multiplicity  $n$  of a matrix  $A(x)$  of form (1). Let also  $m$  be the lowest (non-zero) order of the non-zero derivative  $a^{(m)}(\alpha) \neq 0$  of the entry  $a(x)$  of this matrix at  $x = \alpha$ . Then the number  $m$  (and  $n$ ) is invariant for the class  $\{CA(x)Q(x)\}$  of semiscalarly equivalent matrices, if  $m < n$ .*

*Proof.* Let a matrix  $A(x)$  (1) be semiscalarly equivalent to a matrix  $B(x)$  (3) and  $m'$  be the lowest (non-zero) order of the non-zero derivative  $b^{(m')}(\alpha) \neq 0$  of the entry  $b(x)$  of the matrix  $B(x)$  at  $x = \alpha$ . Suppose that  $m < m'$ . From relations (5) and (7) we obtain

$$s_{21} + s_{22}a(x) - s_{11}b(x) - s_{12}a(x)b(x) = \delta(x)r_{21}(x). \quad (8)$$

Substituting  $x = \alpha$  into (8), we find

$$s_{21} + s_{22}a(\alpha) - s_{11}b(\alpha) - s_{12}a(\alpha)b(\alpha) = 0. \quad (9)$$

Differentiating both sides of equality (8)  $m$  times at  $x = \alpha$ , we obtain

$$s_{22}a^{(m)}(\alpha) - s_{12}a^{(m)}(\alpha)b(\alpha) = 0.$$

The division of both sides of obtained equality by  $a^{(m)}(\alpha) \neq 0$  and the substitution in (9) yields

$$\begin{cases} s_{22} - s_{12}b(\alpha) = 0, \\ s_{21} - s_{11}b(\alpha) = 0. \end{cases}$$

It is impossible, since the matrix  $\|s_{ij}\|_1^2$  is nonsingular. Therefore,  $m \geq m'$ . Taking into account that the semiscalar equivalence is a symmetric relation, we conclude that  $m = m'$ .  $\square$

In this paper, we shall restrict our attention to considering the case in which the partition (2) of the set  $M$  has only two equivalence classes:

$$M = M_1 \cup M_2. \quad (10)$$

Recall that the invariance of the partition (2) was established in Proposition 1.

Since the matrix  $A(x)$  has full rank, the entry  $a(x)$  of the matrix  $A(x)$  (1) satisfies the inequality  $a(\alpha) \neq 0$  for some root  $\alpha \in M$ .

**Theorem 1.3.** *Let the partition (2) of the set  $M$  of characteristic roots of matrices  $A(x)$  (1) and  $B(x)$  (3) be of the form (10). Let also  $n_j$  be the multiplicity of an arbitrary root  $\alpha_j \in M$ ;  $m_j$  be the lowest (non-zero) order of the non-zero derivatives*

$a^{(m_j)}(\alpha_j) \neq 0, b^{(m_j)}(\alpha_j) \neq 0$  of the entries  $a(x), b(x)$  of the matrices  $A(x), B(x)$  at  $x = \alpha_j$ ;

$$a(x) = a_j + \sum_{t=0}^{s-m_j-1} a_{jt}(x - \alpha_j)^{m_j+t}, \quad a_j = a(\alpha_j), \quad a_{j0} \neq 0, \quad (11)$$

$$b(x) = b_j + \sum_{t=0}^{s-m_j-1} b_{jt}(x - \alpha_j)^{m_j+t}, \quad b_j = b(\alpha_j), \quad b_{j0} \neq 0, \quad (12)$$

are binomial decomposition of the entries  $a(x), b(x)$  into degrees of  $x - \alpha_j$  if  $m_j < n_j$ ;  $s = \deg \delta(x)$ ;  $a_i = b_i = 0$  for some  $\alpha_i \in M$ . Matrices  $A(x), B(x)$  are semiscalarly equivalent if and only if there exists a number  $c \neq 0$  satisfying the following conditions:

1)  $a_{i0} = cb_{i0}$  and

$$\begin{vmatrix} a_{i1} & a_{i2} & \dots & a_{i,s_i-1} & a_{i,s_i} \\ a_{i0} & a_{i1} & \dots & a_{i,s_i-2} & a_{i,s_i-1} \\ & \ddots & \ddots & \vdots & \vdots \\ & & \ddots & a_{i1} & a_{i2} \\ 0 & & & a_{i0} & a_{i1} \end{vmatrix} = c^{s_i} \begin{vmatrix} b_{i1} & b_{i2} & \dots & b_{i,s_i-1} & b_{i,s_i} \\ b_{i0} & b_{i1} & \dots & b_{i,s_i-2} & b_{i,s_i-1} \\ & \ddots & \ddots & \vdots & \vdots \\ & & \ddots & b_{i1} & b_{i2} \\ 0 & & & b_{i0} & b_{i1} \end{vmatrix}, \quad (13)$$

$s_i = 1, \dots, m_i - 1, m_i + 1, \dots, n_i - m_i - 1$ , for every root  $\alpha_i \in M$  such that  $a_i = b_i = 0, m_i < n_i$ ;

2)

$$c \left( \frac{a_{i,m_i}}{a_{i0}^2} - \frac{a_{l,m_l}}{a_{l0}^2} \right) = \frac{b_{i,m_i}}{b_{i0}^2} - \frac{b_{l,m_l}}{b_{l0}^2} \quad (14)$$

for every pair of roots  $\alpha_i, \alpha_l \in M$  such that  $a_i = b_i = a_l = b_l = 0, 2m_i < n_i, 2m_l < n_l$ ;

3)

$$a_{i,m_i} + \frac{a_{i0}^2}{a_p} = c \left( b_{i,m_i} + \frac{b_{i0}^2}{b_p} \right) \quad (15)$$

for every root  $\alpha_i \in M$  such that  $a_i = b_i = 0, 2m_i < n_i$  and for some root  $\alpha_p \in M$  such that  $a_p, b_p \neq 0$ ;

$$\begin{aligned}
& 4) \\
& c \begin{vmatrix} 0 & \dots & 0 & a_{p0} & a_{p1} & \dots & a_{ps_p} \\ a_p & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ & \ddots & \ddots & \ddots & \ddots & \ddots & a_{p1} \\ & & \ddots & \ddots & \ddots & \ddots & a_{p0} \\ & & & \ddots & \ddots & \ddots & 0 \\ & & & & \ddots & \ddots & \vdots \\ 0 & & & & & a_p & 0 \end{vmatrix} = \frac{a_p^{m_p+s_p+1}}{b_p^{m_p+s_p+1}} \begin{vmatrix} 0 & \dots & 0 & b_{p0} & b_{p1} & \dots & b_{ps_p} \\ b_p & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ & \ddots & \ddots & \ddots & \ddots & \ddots & b_{p1} \\ & & \ddots & \ddots & \ddots & \ddots & b_{p0} \\ & & & \ddots & \ddots & \ddots & 0 \\ & & & & \ddots & \ddots & \vdots \\ 0 & & & & & b_p & 0 \end{vmatrix}, \\
& \hspace{15em} (16)
\end{aligned}$$

$s_p = 0, 1, \dots, n_p - m_p - 1$ , for every root  $\alpha_p \in M$  such that  $a_p, b_p \neq 0$ ,  $m_p < n_p$ .

## 2. Auxiliary statements

We first obtain two lemmas, which will be used in the proof of Theorem 1.3.

**Lemma 2.1.** *For the existence of the non-zero solution of the equation*

$$\begin{vmatrix} a_0 & b_0 & 0 \\ a_1 & b_1 & 0 \\ \vdots & \vdots & \vdots \\ a_{m-1} & b_{m-1} & 0 \\ a_m & b_m & a_0 b_0 \\ a_{m+1} & b_{m+1} & a_0 b_1 + a_1 b_0 \\ \vdots & \vdots & \vdots \\ a_k & b_k & \sum_{t=0}^{k-m} a_t b_{k-m-t} \end{vmatrix} \cdot \begin{vmatrix} x_1 \\ x_2 \\ x_3 \end{vmatrix} = \bar{0}, \quad (17)$$

over  $\mathbb{C}$ , where  $a_0, b_0 \neq 0$ , it is necessary and sufficient that the following conditions hold:

$$\begin{vmatrix} a_1 & a_2 & \dots & a_{v-1} & a_v \\ a_0 & a_1 & \dots & a_{v-2} & a_{v-1} \\ & \ddots & \ddots & \vdots & \vdots \\ & & \ddots & a_1 & a_2 \\ 0 & & & a_0 & a_1 \end{vmatrix} = c^v \begin{vmatrix} b_1 & b_2 & \dots & b_{v-1} & b_v \\ b_0 & b_1 & \dots & b_{v-2} & b_{v-1} \\ & \ddots & \ddots & \vdots & \vdots \\ & & \ddots & b_1 & b_2 \\ 0 & & & b_0 & b_1 \end{vmatrix}, \quad (18)$$

$v = 1, \dots, m-1, m+1, \dots, k, c = a_0/b_0$ .

Under the conditions of Lemma every non-zero solution  $\|x_{10} \ x_{20} \ x_{30}\|^T$  of the equation (17) has  $x_{10}, x_{20} \neq 0$ .

*Proof. Necessity.* Let the equation (17) have a non-zero solution  $\|x_{10} \ x_{20} \ x_{30}\|^T$ . Then

$$\begin{cases} a_0x_{10} + b_0x_{20} = 0, \\ a_1x_{10} + b_1x_{20} = 0, \\ \dots \dots \dots \\ a_{m-1}x_{10} + b_{m-1}x_{20} = 0, \end{cases} \tag{19}$$

and

$$\begin{cases} a_mx_{10} + b_mx_{20} + a_0b_0x_{30} = 0, \\ a_{m+1}x_{10} + b_{m+1}x_{20} + (a_0b_1 + a_1b_1)x_{30} = 0, \\ \dots \dots \dots \\ a_kx_{10} + b_kx_{20} + x_{30} \sum_{t=0}^{k-m} a_tb_{k-m-t} = 0. \end{cases} \tag{20}$$

We assume that  $x_{10} = 0$  or  $x_{20} = 0$ . Then, from first equalities (19) and (20) we obtain  $x_{10} = x_{20} = x_{30} = 0$ . This contradicts the assumption that the solution  $\|x_{10} \ x_{20} \ x_{30}\|^T$  is non-zero. So, we have  $x_{10}, x_{20} \neq 0$ . From equalities (19) we obtain the conditions (18), where  $c = -x_{20}/x_{10} = a_0/b_0$ , for  $v = 1, \dots, m-1$ .

By (20), if  $a_m = cb_m$ , then  $x_{30} = 0$  and  $a_h = cb_h$  for  $h \in \{m+1, \dots, k\}$ . For this reason in what follows  $a_m \neq cb_m$ , where  $c = -x_{20}/x_{10} = a_0/b_0$ . From the first and second equalities (20) by excluding  $x_{30}$  we obtain

$$a_0a_{m+1} - a_m(cb_1 + a_1) = c^2b_0b_{m+1} - cb_m(cb_1 + a_1). \tag{21}$$

If  $m = 1$ , then  $a_0a_2 - a_1^2 = c^2(b_0b_2 - b_1^2)$ . This means that conditions (20) are fulfilled for  $v = m+1 = 2$ . If  $m > 1$ , then  $a_1 = cb_1$  and from (21) by multiplication of the both sides by  $a_0^{m-1} = c^{m-1}b_0^{m-1}$  we find

$$a_0^m a_{m+1} - 2a_0^{m-1} a_1 a_m = c(b_0^m b_{m+1} - 2b_0^{m-1} b_1 b_m). \tag{22}$$

Denote by  $A_{uw}, B_{uw}$  the submatrices obtained, respectively, from matrices

$$\left\| \begin{array}{cccccc} a_1 & a_2 & \dots & a_m & a_{m+1} & \\ a_0 & a_1 & \dots & a_{m-1} & a_m & \\ & \ddots & \ddots & \vdots & \vdots & \\ & & \ddots & a_1 & a_2 & \\ 0 & & & a_0 & a_1 & \end{array} \right\| \text{ and } \left\| \begin{array}{cccccc} b_1 & b_2 & \dots & b_m & b_{m+1} & \\ b_0 & b_1 & \dots & b_{m-1} & b_m & \\ & \ddots & \ddots & \vdots & \vdots & \\ & & \ddots & b_1 & b_2 & \\ 0 & & & b_0 & b_1 & \end{array} \right\| \tag{23}$$

by obliterating of two last columns and  $u$ -th and  $w$ -th rows. Denote also by  $\Delta_{m+1}(A), \Delta_{m+1}(B)$  the determinants of matrices (23), respectively. Decompose them for minors of order two that are contained in the last two columns. Since  $|A_{uw}| = |B_{uw}| =$

0 for  $u \neq m + 1$ , we have

$$\Delta_{m+1}(A) = (-1)^{m+1} \left( \begin{vmatrix} a_m & a_{m+1} \\ a_0 & a_1 \end{vmatrix} \cdot |A_{1, m+1}| - \begin{vmatrix} a_{m-1} & a_m \\ a_0 & a_1 \end{vmatrix} \cdot |A_{2, m+1}| + \dots + \begin{vmatrix} a_1 & a_2 \\ a_0 & a_1 \end{vmatrix} \cdot |A_{m, m+1}| \right),$$

and

$$\Delta_{m+1}(B) = (-1)^{m+1} \left( \begin{vmatrix} b_m & b_{m+1} \\ b_0 & b_1 \end{vmatrix} \cdot |B_{1, m+1}| - \begin{vmatrix} b_{m-1} & b_m \\ b_0 & b_1 \end{vmatrix} \cdot |B_{2, m+1}| + \dots + \begin{vmatrix} b_1 & b_2 \\ b_0 & b_1 \end{vmatrix} \cdot |B_{m, m+1}| \right).$$

Since  $\|a_0 a_1 \dots a_{m-1}\| = c \|b_0 b_1 \dots b_{m-1}\|$ , each summand of expression in parenthesis for  $\Delta_{m+1}(A)$ , possibly except the first two, differs from the corresponding summand for  $\Delta_{m+1}(B)$  by the multiplier  $c^{m+1}$ . From this fact and from the equality (22) the equality (18) follows for  $v = m + 1$ .

Denote by  $\Delta_v(A)$  and  $\Delta_v(B)$  the determinants in the left and right sides of equality (18), respectively. Suppose by induction  $\Delta_r(A) = c^r \Delta_r(B)$  for all  $r$  such that  $m < r < k$ . Accept for the sake of determinacy  $r > 2m$ . In the case where  $r \leq 2m$  the proof is analogous. From the first  $r$  equalities (20) by excluding  $x_{30}$  and by sufficiently evident transformations we arrive at the system

$$\left\{ \begin{array}{l} (a_{m+1} - (a_0 b_0)^{-1} a_m \sum_{u=0}^1 a_u b_{1-u}) (-a_0)^m \Delta_{r-m}(A) = \\ \quad = c^{r+1} (b_{m+1} - (a_0 b_0)^{-1} b_m \sum_{u=0}^1 a_u b_{1-u}) (-b_0)^m \Delta_{r-m}(B), \\ (a_{m+2} - (a_0 b_0)^{-1} a_m \sum_{u=0}^2 a_u b_{2-u}) (-a_0)^{m+1} \Delta_{r-m-1}(A) = \\ \quad = c^{r+1} (b_{m+2} - (a_0 b_0)^{-1} b_m \sum_{u=0}^2 a_u b_{2-u}) (-b_0)^{m+1} \Delta_{r-m-1}(B), \\ \dots \\ (a_{r-m+1} - (a_0 b_0)^{-1} a_m \sum_{u=0}^{r-2m+1} a_u b_{r-2m-u+1} - a_{r-m+1} + \\ \quad + (a_0 b_0)^{-1} a_m \sum_{u=0}^{r-2m+1} a_u b_{r-2m-u+1}) (-a_0)^{r-m} \Delta_m(A) = \\ \quad = c^{r+1} (b_{r-m+1} - (a_0 b_0)^{-1} b_m \sum_{u=0}^{r-2m+1} a_u b_{r-2m-u+1} - b_{r-m+1} + \\ \quad \quad + (a_0 b_0)^{-1} b_m \sum_{u=0}^{r-2m+1} a_u b_{r-2m-u+1}) (-b_0)^{r-m} \Delta_m(B), \\ \dots \\ (a_r - (a_0 b_0)^{-1} a_m \sum_{u=0}^{r-m} a_u b_{r-m-u}) (-a_0)^{r-1} \Delta_1(A) = \\ \quad = c^{r+1} (b_r - (a_0 b_0)^{-1} b_m \sum_{u=0}^{r-m} a_u b_{r-m-u}) (-b_0)^{r-1} \Delta_1(B), \\ (a_{r+1} - (a_0 b_0)^{-1} a_m \sum_{u=0}^{r-m+1} a_u b_{r-m-u+1}) (-a_0)^r = \\ \quad = c^{r+1} (b_{r+1} - (a_0 b_0)^{-1} b_m \sum_{u=0}^{r-m+1} a_u b_{r-m-u+1}) (-b_0)^r, \end{array} \right. \quad (24)$$

where  $c = a_0/b_0$ .



Adding the left sides of equality (24) and separately the right sides, we obtain the equality:

$$\begin{aligned}
 & (-a_0)^r a_{r+1} + (-a_0)^{r-1} a_r \Delta_1(A) + \dots + (-a_0)^{r-m} a_{r-m+1} \Delta_m(A) + \dots \\
 & + (-a_0)^m a_{m+1} \Delta_{r-m}(A) + (-a_0)^{m-1} a_m \Delta_{r-m+1}(A) - \\
 & - (a_0 b_0)^{-1} a_m (b_1 \Delta_{r-m}(A) (-a_0)^{m+1} + b_2 \Delta_{r-m-1}(A) (-a_0)^{m+2} + \dots \\
 & + b_{j, r-m_j} \delta_{j1}(A) (-a_{j0})^r) + (a_0 b_0)^{-1} a_m (b_1 \Delta_{r-m}(A) (-a_0)^{m+1} + \\
 & + b_2 \Delta_{r-m-1}(A) (-a_0)^{m+2} + \dots + b_{r-m} \Delta_1(A) (-a_0)^r + b_{r-m+1} (-a_0)^{r+1}) - \\
 & - (a_{r-m+1} - (a_0 b_0)^{-1} a_m \sum_{u=0}^{r-2m+1} a_u b_{r-2m-u+1}) (-a_0)^{r-m} \Delta_m(A) = \\
 & = c^{r+1} ((-b_0)^r b_{r+1} + (-b_0)^{r-1} b_r \Delta_1(B) + \dots + (-b_0)^{r-m} b_{r-m+1} \Delta_m(B) + \dots \\
 & + (-b_0)^m b_{m+1} \Delta_{r-m}(B) + (-b_0)^{m-1} b_m \Delta_{r-m+1}(B) - \\
 & - (a_0 b_0)^{-1} b_m (a_1 \Delta_{r-m}(B) (-b_0)^{m+1} + a_2 \Delta_{r-m-1}(B) (-b_0)^{m+2} + \dots \\
 & + a_{r-m} \Delta_1(B) (-b_0)^r) + (a_0 b_0)^{-1} b_m (a_1 \Delta_{r-m}(B) (-b_0)^{m+1} + \\
 & + a_2 \Delta_{r-m-1}(B) (-b_0)^{m+2} + \dots + a_{r-m} \Delta_1(B) (-b_0)^r + a_{r-m+1} (-b_0)^{r+1}) - \\
 & - (b_{r-m+1} - (a_0 b_0)^{-1} b_m \sum_{u=0}^{r-2m+1} a_u b_{r-2m-u+1}) (-b_{j0})^{r-m_j} \Delta_m(B). \tag{25}
 \end{aligned}$$

Grouping similar terms in both sides of obtained equality we arrive to

$$\begin{aligned}
 & (-a_0)^r a_{r+1} + (-a_0)^{r-1} a_r \Delta_1(A) + \dots + (-a_0)^{r-m} a_{r-m+1} \Delta_m(A) + \dots + \\
 & + (-a_0)^m a_{m+1} \Delta_{r-m}(A) + (-a_0)^{m-1} a_m \Delta_{r-m+1}(A) + (a_0 b_0)^{-1} a_m b_{r-m+1} (-a_0)^{r+1} - \\
 & - (a_{r-m+1} - (a_0 b_0)^{-1} a_m \sum_{u=1}^{r-2m+1} a_u b_{r-2m-u+1}) (-a_0)^{r-m} \Delta_m(A) = \\
 & = c^{r+1} ((-b_0)^r b_{r+1} + (-b_0)^{r-1} b_r \Delta_1(B) + \dots + (-b_0)^{r-m} b_{r-m+1} \Delta_m(B) + \dots \\
 & + (-b_0)^m b_{m+1} \Delta_{r-m}(B) + (-b_0)^{m-1} b_m \Delta_{r-m+1}(B) + (a_0 b_0)^{-1} b_m a_{r-m+1} (-b_0)^{r+1} - \\
 & - (b_{r-m+1} - (a_0 b_0)^{-1} b_m \sum_{u=1}^{r-2m+1} a_u b_{r-2m-u+1}) (-b_0)^{r-m} \Delta_m(B). \tag{26}
 \end{aligned}$$

It follows from (20), that

$$\begin{aligned}
 a_{r-m+1} + (a_0 b_0)^{-1} a_m \sum_{u=0}^{r-2m+1} a_u b_{r-2m-u+1} & = \\
 & = c (b_{r-m+1} + (a_0 b_0)^{-1} b_m \sum_{u=0}^{r-2m+1} a_u b_{r-2m-u+1}).
 \end{aligned}$$

From this relation it is easy to check that the following equality is true

$$\begin{aligned}
& (a_0 b_0)^{-1} a_m b_{r-m+1} (-a_0)^{r+1} - \\
& - (a_{r-m+1} - (a_0 b_0)^{-1} a_m \sum_{u=0}^{r-2m+1} a_u b_{r-2m-u+1}) (-a_0)^{r-m} \Delta_m(A) = \\
& = c^{r+1} ((a_0 b_0)^{-1} b_m a_{r-m+1} (-b_0)^{r+1} - \\
& - (b_{r-m+1} (a_0 b_0)^{-1} b_m \sum_{u=0}^{r-2m+1} a_u b_{r-2m-u+1}) (-b_0)^{r-m} \Delta_m(B)).
\end{aligned} \tag{27}$$

From (19) and the induction hypothesis we can write

$$\begin{aligned}
& (-a_0)^{m-2} a_{m-1} \Delta_{r-m+2}(A) + \dots + (-a_0) a_2 \Delta_{r-1}(A) + a_1 \Delta_r(A) = \\
& c^{r+1} ((-b_0)^{m-2} b_{m-1} \Delta_{r-m+2}(B) + \dots + (-b_{j_0}) b_{j_2} \delta_{j,r-1}(B) + b_1 \Delta_r(B)).
\end{aligned} \tag{28}$$

Comparing (26), (27) and (28), we obtain equality

$$\begin{aligned}
& (-a_0)^r a_{r+1} + (-a_0)^{r-1} a_r \Delta_1(A) + \dots + (-a_0) a_2 \Delta_{r-1}(A) + a_1 \Delta_r(A) = \\
& c^{r+1} ((-b_0)^r b_{r+1} + (-b_0)^{r-1} b_r \Delta_1(B) + \dots + (-b_0) b_2 \Delta_{r-1}(B) + b_1 \Delta_r(B)),
\end{aligned} \tag{29}$$

i.e.  $\Delta_{r+1}(A) = c^{r+1} \Delta_{r+1}(B)$ , where  $c = a_0/b_0$ . The necessity of conditions of the Lemma is proved.

**Sufficiency.** Consider equalities (19) and (20) as one system of equations in three unknowns  $x_{10}$ ,  $x_{20}$ ,  $x_{30}$ . In conditions (18)  $c = a_0/b_0$ . This means that  $x_{10} = 1$ ,  $x_{20} = -c$  satisfies the first equation of the system (19). From condition with  $v = 1$  it follows that  $x_{10} = 1$ ,  $x_{20} = -c$  satisfies the second equation of (19). Next, third and all following equalities in system (19) for  $x_{10} = 1$ ,  $x_{20} = -c$  can be recurrently obtained from conditions (18) with  $v = 2, \dots, m-1$ . Evidently, the values

$$x_{10} = 1, x_{20} = -c, x_{30} = (cb_m - a_m)(a_0 b_0)^{-1}, \tag{30}$$

satisfy the first equation (20). We compute both determinants in equality (20) with  $v = m+1$ . After annihilation of equal summands on both sides of the obtained equality and after division by  $a_0^m = c^m b_0^m$  with the help of simple transformations we can obtain the following relation

$$a_{m+1} - c b_{m+1} + (cb_m - a_m)(a_0 b_0)^{-1}(a_0 b_1 + a_1 b_1) = 0.$$

This means that (30) satisfies the second equation of system (20).



the following conditions hold:

$$\underbrace{\begin{vmatrix} 0 & \dots & 0 & a_0 & a_1 & \dots & a_v \\ a & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ & \ddots & \ddots & \ddots & \ddots & \ddots & a_1 \\ & & \ddots & \ddots & \ddots & \ddots & a_0 \\ & & & \ddots & \ddots & \ddots & 0 \\ & & & & \ddots & \ddots & \vdots \\ 0 & & & & & a & 0 \end{vmatrix}}_{m+v} = \frac{a^{m+v-1}a_0}{b^{m+v-1}b_0} \underbrace{\begin{vmatrix} 0 & \dots & 0 & b_0 & b_1 & \dots & b_v \\ b & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ & \ddots & \ddots & \ddots & \ddots & \ddots & b_1 \\ & & \ddots & \ddots & \ddots & \ddots & b_0 \\ & & & \ddots & \ddots & \ddots & 0 \\ & & & & \ddots & \ddots & \vdots \\ 0 & & & & & b & 0 \end{vmatrix}}_{m+v} \quad (33)$$

for  $v \in \{0, 1, \dots, k\}$ .

Under the conditions of Lemma in every non-zero solution  $\|x_{10} \ x_{20} \ x_{30}\|^T$  of the equation (17) it is necessary that  $x_{10}, x_{20} \neq 0$ .

*Proof. Necessity.* The equality (33) for  $v = 0$  is trivially true. Assume that the equation (32) has a non-zero solution. Then the matrix of this equation has rank less than 3. The two first rows of the matrix are linearly independent. Then, each other row of the matrix linearly depends on these first two rows. Since

$$\begin{vmatrix} a & b & ab \\ a_0 & b_0 & ab_0 + a_0b \\ a_v & b_v & ab_v + a_vb \end{vmatrix} = 0, \quad v = 1, \dots, m-1,$$

we have  $\frac{a_v}{a_0} = \frac{b_v}{b_0}$ . This implies the equality (33) for  $v = 1, \dots, m-1$ . From the equality

$$\begin{vmatrix} a & b & ab \\ a_0 & b_0 & ab_0 + a_0b \\ a_m & b_m & ab_m + a_0b_0 + a_mb \end{vmatrix} = 0$$

we have

$$ab_0^2a_0 - a_0b_mab + a_mb_0ab - a_0^2b_0b = 0,$$

whence, after division of both sides by  $aba_0b_0$ , we obtain equality (33) for  $v = m$ . Assume by induction that equality (33) holds for all  $v$  with  $m \leq v \leq r < k$ . Consider the equality

$$\begin{vmatrix} a & b & ab \\ a_0 & b_0 & ab_0 + a_0b \\ a_{r+1} & b_{r+1} & ab_{r+1} + \sum_{t=0}^{r+1-m} a_t b_{r+1-m-t} + a_{r+1}b \end{vmatrix} = 0.$$



It is easy to make sure that the following identity is true

$$\begin{aligned}
& b^{-1}(b_0(a^r a_0)^{-1} \Delta_{r-m+1}(a)(-1)^{m-1} + b_1(a^{r-1} a_0)^{-1} \Delta_{r-m}(a)(-1)^m + \dots + \\
& + b_{r-m}(a^m a_0)^{-1} \Delta_1(a)(-1)^{r-1} + b_{r-m+1}(a^{m-1} a_0)^{-1} \Delta_0(a)(-1)^r) + \\
& + a^{-1}(-a_0(b^r b_0)^{-1} \Delta_{r-m+1}(b)(-1)^{m-1} - a_1(b^{r-1} b_0)^{-1} \Delta_{r-m}(b)(-1)^m - \dots - \\
& - a_{r-m}(b^m b_0)^{-1} \Delta_1(b)(-1)^{r-1} - a_{r-m+1}(b^{m-1} b_0)^{-1} \Delta_0(b)(-1)^r) + \quad (39) \\
& + (-1)^{m+r} (a_0 b)^{-1} \sum_{t=0}^{r+1-m} a_t b_{r+1-m-t} - (-1)^{m+r} (a b_0)^{-1} \sum_{t=0}^{r+1-m} a_t b_{r+1-m-t} = 0.
\end{aligned}$$

We add left sides of equalities (35), (36) and multiplied by  $(-1)^{m+r}$  left side of the equality (34) and separately we add right sides of the these equalities. Taking into account the expression (37), (38) for  $(a^{m+r} a_0)^{-1} \Delta_{r+1}(a)$ ,  $(b^{m+r} b_0)^{-1} \Delta_{r+1}(b)$  and the identity (39), we obtain  $(a^{m+r} a_0)^{-1} \Delta_{r+1}(a) - (b^{m+r} b_0)^{-1} \Delta_{r+1}(b) = 0$ . This means that the equality (33) for  $v = r + 1$  is true. The necessity of the conditions of Lemma is proved.

**Sufficiency.** Assume that the conditions (33) are satisfied for  $v = 1, \dots, m - 1$ . Then, the submatrix formed by first  $m + 1$  rows of the matrix from the equation (32) has rank less than 3. In fact, this rank is equal to 2, because, as has been stated above, the first two rows of this matrix are linearly independent. The equality (33) for  $v = m$  implies that  $(m + 2)$ th row of the matrix of equation (32) linearly depends on its first two rows. Our inductive assumption is the following. Let each of the first  $r + 2$ ,  $m \leq r < k$ , rows of the matrix of equation (32) linearly depend on its first two rows. We now reverse the order of arguments, as compared to the proof of the necessity, passing from relation (33) for  $v = r + 1$  to relation (34). This relation implies that the minor of order 3 of the matrix of equation (32) that is contained in 1-th, 2-nd and  $(r + 3)$ th rows is equal to zero. This means that indicated rows are linearly dependent. The above argument inductively proves that the matrix of equation (32) has the 2. This implies that the equation (32) has a non-zero solution.

The remaining part of the lemma will be proved by contradiction. Let  $\|x_{10} \ x_{20} \ x_{30}\|^T$  be a non-zero solution of the equation (32), where  $x_{10} = 0$  or  $x_{20} = 0$ . For this reason we have the equality

$$\left\| \begin{array}{cc} b & ab \\ b_0 & ab_0 + a_0 b \end{array} \right\| \cdot \left\| \begin{array}{c} x_{20} \\ x_{30} \end{array} \right\| = \bar{0} \quad \text{or} \quad \left\| \begin{array}{cc} a & ab \\ a_0 & ab_0 + a_0 b \end{array} \right\| \cdot \left\| \begin{array}{c} x_{10} \\ x_{30} \end{array} \right\| = \bar{0}.$$

Since  $a, b, a_0, b_0 \neq 0$ , the determinants of the  $2 \times 2$  matrices of these equations are different from 0. Hence,  $\|x_{20} \ x_{30}\|^T = \bar{0}$  or  $\|x_{10} \ x_{30}\|^T = \bar{0}$ , i.e., actually,  $\|x_{10} \ x_{20} \ x_{30}\|^T = \bar{0}$ , contrary to the our assumption. Therefore, in the non-zero

solution  $\|x_{10} \ x_{20} \ x_{30}\|^T$  of the equation (32) we necessarily have  $x_{10}, \ x_{20} \neq 0$ . Lemma is proved.  $\square$

### 3. Proof of the Theorem

**Necessity.** Assume that matrices  $A(x)$  and  $B(x)$  of the forms (1) and (3) are semiscalarly equivalent. Then, the entries  $a(x)$  and  $b(x)$  of these matrices satisfy the congruence

$$s_{22}a(x) - s_{11}b(x) - s_{12}a(x)b(x) \equiv 0 \pmod{\delta(x)}, \quad (40)$$

where  $s_{11}, \ s_{22} \neq 0$ . Construct the decompositions of the entries  $a(x), \ b(x)$  into degrees of binomial  $x - \alpha_i$  (see (11), (12)) for  $\alpha_i \in M$  such that  $a_i = b_i = 0, \ m_i < n_i$ , and compare the coefficients of equal degrees of binomial  $x - \alpha_i$  on both sides of the congruence (40). Then, we obtain

$$s_{22}a_{it} - s_{11}b_{it} = 0 \text{ for } t \in \{0, 1, \dots, l_i - m_i - 1\} \text{ where } l_i := \min(2m_i, n_i).$$

This implies  $s_{11}/s_{22} = a_{i0}/b_{i0}$  and equality (13), where  $c = s_{11}/s_{22} = a_{i0}/b_{i0}$ , for all  $\alpha_i \in M$  such that  $2m_i \geq n_i$ .

Taking into account the roots  $\alpha_i \in M$  such that  $2m_i < n_i$  and comparing the coefficients of equal degrees of binomial  $x - \alpha_i$  on both sides of the congruence (40), we obtain the system of the equalities, written in matrix form:

$$\left\| \begin{array}{ccc} a_{i0} & b_{i0} & 0 \\ a_{i1} & b_{i1} & 0 \\ \vdots & \vdots & \vdots \\ a_{i, m_i-1} & b_{i, m_i-1} & 0 \\ a_{im_i} & b_{im_i} & a_{i0}b_{i0} \\ a_{i, m_i+1} & b_{i, m_i+1} & a_{i0}b_{i1} + a_{i1}b_{i0} \\ \vdots & \vdots & \vdots \\ a_{i, n_i-m_i-1} & b_{i, n_i-m_i-1} & \sum_{t=0}^{n_i-2m_i-1} a_{it}b_{i, n_i-2m_i-1-t} \end{array} \right\| \cdot \left\| \begin{array}{c} s_{22} \\ -s_{11} \\ -s_{12} \end{array} \right\| = \bar{0}.$$

Since  $s_{11}, s_{22} \neq 0$ , this equality implies the first condition of Theorem 1.3 (according to Lemma 2.1).

Let  $\alpha_i, \ \alpha_l \in M$  be the arbitrary pair of the entries such that  $a_i = b_i = 0, \ 2m_i < n_i, \ a_l = b_l = 0, \ 2m_l < n_l$ . For the coefficients  $a_{im_i}, \ a_{lm_l}, \ b_{im_i}, \ b_{lm_l}, \ a_{i0} \neq 0, \ a_{l0} \neq 0, \ b_{i0} \neq 0, \ b_{l0} \neq 0$  of the decomposition of the forms (11), (12) into degrees of binomials  $x - \alpha_i, \ x - \alpha_l$ , the congruence (40) implies the relations

$$s_{22}a_{im_i} - s_{11}b_{im_i} - s_{12}a_{i0}b_{i0} = 0, \quad s_{22}a_{lm_l} - s_{11}b_{lm_l} - s_{12}a_{l0}b_{l0} = 0.$$

Excluding  $s_{12}$  from these equalities and considering that  $a_{i0} = cb_{i0}$ ,  $a_{l0} = cb_{l0}$ ,  $c = s_{11}/s_{22}$ , we obtain the equality (14) from the second condition of the Theorem.

For the coefficients  $a_{i0}$ ,  $a_{im_i}$ ,  $b_{i0}$ ,  $b_{im_i}$  and for the free terms of the decompositions of the form (11), (12) into degrees of  $x - \alpha_i$ ,  $x - \alpha_p$  for roots  $\alpha_i$ ,  $\alpha_p \in M$  such that  $a_i = b_i = 0$ ,  $2m_i < n_i$ ,  $a_p$ ,  $b_p \neq 0$ , the congruence (40) yields the system

$$\begin{cases} s_{22}a_{i0} - s_{11}b_{i0} = 0, \\ s_{22}a_{im_i} - s_{11}b_{im_i} - s_{12}a_{i0}b_{i0} = 0, \\ s_{22}a_p - s_{11}b_p - s_{12}a_p b_p = 0. \end{cases}$$

Excluding  $s_{12}$  from two last equalities of these system and putting  $a_{i0} = cb_{i0}$  for  $c = s_{11}/s_{22}$ , we obtain the equality (15) from the third condition of Theorem 1.3.

For the coefficients of the decompositions (11), (12) into degrees of binomial  $x - \alpha_p$  for roots  $\alpha_p \in M$  such that  $a_p$ ,  $b_p \neq 0$ ,  $m_p < n_p$ , the congruence (40) yields the system

$$\begin{cases} s_{22}a_p - s_{11}b_p - s_{12}a_p b_p = 0, \\ s_{22}a_{pt} - s_{11}b_{pt} - s_{12}(a_{pt}b_p + b_{pt}a_p) = 0, \end{cases}$$

for  $t \in \{0, 1, \dots, l_p - m_p - 1\}$  where  $l_p = \min(2m_p, n_p)$ . After excluding  $s_{12}$  from this system we obtain

$$c \frac{a_{pt}}{a_p^2} = \frac{b_{pt}}{b_p^2} \quad (41)$$

for  $t \in \{0, 1, \dots, l_p - m_p - 1\}$ , where  $c = s_{11}/s_{22}$ . This implies the equality (16) for the roots  $\alpha_p \in M$  such that  $a_p$ ,  $b_p \neq 0$ ,  $m_p < n_p$ ,  $2m_p \geq n_p$ . If  $2m_p < n_p$ , then the congruence (40) implies the system of the equalities, written in the matrix form as

$$\|F \ G \ H\| \cdot \|s_{22} \ -s_{11} \ -s_{12}\|^T = \bar{0},$$

where

$$F = \begin{pmatrix} a_p \\ a_{p0} \\ a_{p1} \\ \vdots \\ a_{p, m_p-1} \\ a_{pm_p} \\ a_{p, m_p+1} \\ \vdots \\ a_{p, n_p-m_p-1} \end{pmatrix}, \quad G = \begin{pmatrix} b_p \\ b_{p0} \\ b_{p1} \\ \vdots \\ b_{p, m_p-1} \\ b_{pm_p} \\ b_{p, m_p+1} \\ \vdots \\ b_{p, n_p-m_p-1} \end{pmatrix}, \quad (42)$$



$$H = \left\| \begin{array}{c} a_p b_p \\ a_{p0} b_p + a_p b_{p0} \\ a_{p1} b_p + a_p b_{p1} \\ \vdots \\ a_{p, m_p-1} b_p + a_p b_{p, m_p-1} \\ a_{pm_p} b_p + a_{p0} b_{p0} + a_p b_{pm_p} \\ a_{p, m_p+1} b_p + a_{p0} b_{p1} + a_{p1} b_{p0} + a_p b_{p, m_p+1} \\ \vdots \\ a_{p, n_p-m_p-1} b_p + \sum_{t=0}^{n_p-2m_p-1} a_{pt} b_{p, n_p-2m_p-1-t} + a_p b_{p, n_p-m_p-1} \end{array} \right\|. \quad (43)$$

Since  $s_{11}, s_{22} \neq 0$ , by the Lemma 2 from the obtained equality we have

$$\underbrace{\left\| \begin{array}{ccccccc} 0 & \dots & 0 & a_{p0} & a_{p1} & \dots & a_{ps_p} \\ a_p & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & a_{p1} \\ \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & a_{p0} \\ \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & 0 \\ \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ 0 & & & & a_p & & 0 \end{array} \right\|}_{m_p+s_p} = \frac{a_p^{m_p+s_p-1} a_{p0}}{b_p^{m_p+s_p-1} b_{p0}} \underbrace{\left\| \begin{array}{ccccccc} 0 & \dots & 0 & b_{p0} & b_{p1} & \dots & b_{ps_p} \\ b_p & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & b_{p1} \\ \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & b_{p0} \\ \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & 0 \\ \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ 0 & & & & b_p & & 0 \end{array} \right\|}_{m_p+s_p}, \quad (44)$$

for  $s_p \in \{0, 1, \dots, n_p - m_p - 1\}$ . Here for  $t \in \{0, 1, \dots, m_p - 1\}$  the equalities (41) hold. Now multiply the left side of the equality (44) by the left side of the equality (41) at  $t = 0$  and carry out analogous operation with right sides. As a result, we obtain the equality (16). This proves the last condition of Theorem 1.3 and necessity entirely.

**Sufficieny.** Assume that the conditions 1) – 4) of Theorem 1.3 are fulfilled. Consider the matrix equations

$$\left\| \begin{array}{ccc} a_{i0} & b_{i0} & 0 \\ a_{i1} & b_{i1} & 0 \\ \vdots & \vdots & \vdots \\ a_{i, m_i-1} & b_{i, m_i-1} & 0 \end{array} \right\| \cdot \left\| \begin{array}{c} x_1 \\ x_2 \\ x_3 \end{array} \right\| = \bar{0}, \quad (45)$$

$$\left\| \begin{array}{ccc} a_{im_i} & b_{im_i} & a_{i0}b_{i0} \\ a_{i, m_i+1} & s_{i, m_i+1} & a_{i0}b_{i1} + a_{i1}b_{i0} \\ \vdots & \vdots & \vdots \\ a_{i, n_i-m_i-1} & b_{i, n_i-m_i-1} & \sum_{t=0}^{n_i-2m_i-1} a_{it}b_{i, n_i-2m_i-1-t} \end{array} \right\| \cdot \left\| \begin{array}{c} x_1 \\ x_2 \\ x_3 \end{array} \right\| = \bar{0}, \quad (46)$$

in the unknown vector-column  $\|x_1 \ x_2 \ x_3\|^T$  are written using the coefficients of the decompositions (11), (12) for some root  $\alpha_i \in M$ . For an arbitrary fixed root  $\alpha_i \in M$  such that  $a_i = b_i = 0$ ,  $m_i < n_i$ ,  $2m_i \geq n_i$ , by Lemma 2.1 the equation (45) has a non-zero solution. If  $2m_i < n_i$ , then by Lemma 2.1 both these equations (45), (46) have common non-zero solution. Observe that first two rows of the matrix of each of these equations are linearly independent and each other row of this matrix is a linear combination of the first two rows. Taking into account that

$$\left\| \begin{array}{ccc} a_{i0} & b_{i0} & 0 \\ a_{im_i} & b_{im_i} & a_{i0}b_{i0} \end{array} \right\| \cdot \left\| \begin{array}{c} 1 \\ -c \\ (cb_{im_i} - a_{im_i})(a_{i0}b_{i0})^{-1} \end{array} \right\| = \bar{0},$$

we conclude that

$$\|x_1 \ x_2 \ x_3\|^T = \|1 \ -c \ (cb_{im_i} - a_{im_i})(a_{i0}b_{i0})^{-1}\|^T \quad (47)$$

is the common non-zero solution of the equations (45), (46).

From the condition 2) we deduce the relation

$$(cb_{im_i} - a_{im_i})(a_{i0}b_{i0})^{-1} = (cb_{lm_l} - a_{lm_l})(a_{l0}b_{l0})^{-1}$$

for other arbitrary root  $\alpha_l \in M$  such that  $a_l = b_l = 0$ ,  $2m_l < n_l$ , different from  $\alpha_i \in M$ . This means that the solution (47) does not depend on the choice of the root  $\alpha_i \in M$  satisfying the condition 1). So, we can write the congruence

$$a(x) - cb(x) - c_1a(x)b(x) \equiv 0 \pmod{(x - \alpha_i)^{n_i}}, \quad (48)$$

where

$$c_1 = (a_{im_i} - cb_{im_i})(a_{i0}b_{i0})^{-1}, \quad (49)$$

for arbitrary root  $\alpha_i \in M$  such that  $a_i = b_i = 0$  (including the roots  $\alpha_i$  with  $m_i \geq n_i$ , if such roots exist). Thus, one subset of the partition (10) of the set  $M$  of roots  $\alpha_i$  is exhausted.

The fulfillment of the equality (16) for  $s_p = 0$  means that  $\frac{ca_{p0}}{a_p^2} = \frac{b_{p0}}{b_p^2}$ . Taking into account this fact, we see that the relation (44) follows from the equality (16) for every  $s_p \in \{1, \dots, n_p - m_p - 1\}$ . Moreover, by Lemma 2.2 the equation

$$\|F \ G \ H\| \cdot \|x_1 \ x_2 \ x_3\|^T = \bar{0} \quad (50)$$

(with  $F$ ,  $G$  and  $H$  defined as in (42), (43)) has a non-zero solution. Since the first two rows of the matrix of the equation (50) are linearly independent and

$$\left\| \begin{array}{ccc} a_p & b_p & a_p b_p \\ a_{p0} & b_{p0} & a_{p0} b_p + a_p b_{p0} \end{array} \right\| \left\| \begin{array}{c} 1 \\ -c \\ (cb_p - a_p)(a_p b_p)^{-1} \end{array} \right\| = \bar{0},$$

the column  $\|x_1 \ x_3 \ x_3\|^T = \|1 \ -c \ (cb_p - a_p)(a_p b_p)^{-1}\|^T$  is the solution of the equation (50) for arbitrary root  $\alpha_p \in M$  such that  $a_p, b_p \neq 0$ . From the condition 3) we deduce the relation

$$(cb_p - a_p)(a_p b_p)^{-1} = (cb_{im_i} - a_{im_i})(a_{i0} b_{i0})^{-1}.$$

Therefore, a solution of the equation (50) can be written in the form (47). This means that the following congruence is satisfied

$$a(x) - cb(x) - c_1 a(x)b(x) \equiv 0 \pmod{(x - \alpha_p)^{n_p}}, \quad (51)$$

where  $c_1$  is defined as in congruence (48) (see (49)). In addition, the congruence (51) is fulfilled for arbitrary root  $\alpha_p \in M$  such that  $a_p, b_p \neq 0$ . Now, comparing (48) and (51), we obtain the congruence

$$a(x) - cb(x) - c_1 a(x)b(x) \equiv 0 \pmod{\delta(x)}.$$

Let  $r_{21}(x)$  be determined as the result of dividing the left side of the congruence by  $\delta(x)$ . Then, we can write

$$a(x) - cb(x) - c_1 a(x)b(x) = \delta(x)r_{21}(x). \quad (52)$$

Write (52) in the form  $a(x) = b(x)r_{11}(x) + \delta(x)r_{21}(x)$ , where  $r_{11}(x) = c + c_1 a(x)$ , and denote  $r_{12}(x) = c_1 \delta(x)$ ,  $r_{22}(x) = 1 - b(x)r_{12}(x)$ . Now it can be written

$$\left\| \begin{array}{cc} c & c_1 \\ 0 & 1 \end{array} \right\| \left\| \begin{array}{cc} 1 & 0 \\ a(x) & \delta(x) \end{array} \right\| = \left\| \begin{array}{cc} 1 & 0 \\ b(x) & \delta(x) \end{array} \right\| \left\| \begin{array}{cc} r_{11}(x) & r_{12}(x) \\ r_{21}(x) & r_{22}(x) \end{array} \right\|.$$

It is evident that the left factor on the left side of determined equation and the right factor on the right side are invertible matrices. Thus, the matrices  $A(x)$  and  $B(x)$  are semiscalarly equivalent. Theorem 1.3 is proved.

#### 4. Some corollaries of Theorem 1.3

**Corollary 4.1.** *Let the partition (2) of the set  $M$  of characteristic roots of matrices  $A(x)$  be of the form (10). Let also the entry  $a(x)$  of the matrix  $A(x)$  in the notations of Theorem 1.3 satisfy the following conditions:*

- 1)  $m_i < n_i \leq 2m_i, a_i = 0$ , for roots  $\alpha_i \in M, i \in \{1, \dots, r\}$  where  $r \geq 1$ ;
- 2)  $m_p < n_p \leq 2m_p, a_p \neq 0$ , for roots  $\alpha_p \in M, p \in \{r+1, \dots, q\}$  where  $q > r$ ;
- 3)  $m_h \geq n_h$  for all  $\alpha_h \in M$  which are different from  $\alpha_i, \alpha_p$ .

Such a matrix  $A(x)$  in the class  $\{CA(x)Q(x)\}$  of semiscalarly equivalent matrices is determined up to constant factors  $c \neq 0$  and  $1/c$  of the rows

$$\|a_{i0} \ a_{i1} \ \dots \ a_{i, n_i - m_i - 1} \ \dots \ a_{r0} \ a_{r1} \ \dots \ a_{r, n_r - m_r - 1}\|$$

and

$$\frac{1}{a_{r+1}^2} \|a_{r+1,0} \ a_{r+1,1} \ \dots \ a_{r+1, n_{r+1} - m_{r+1} - 1} \ \dots \ a_{q0} \ a_{q1} \ \dots \ a_{q, n_q - m_q - 1}\|,$$

respectively.

*Proof.* Since  $n_i \leq 2m_i$  and  $n_p \leq 2m_p$ , it follows that  $n_i - m_i - 1 \leq m_i - 1$  and  $n_p - m_p - 1 \leq m_p - 1$ . By Theorem 1.3, in the case under consideration matrices  $A(x), B(x)$  are semiscalarly equivalent if and only if the ratio  $c = a_{i0}/b_{i0}$  does not depend on the choice of  $i \in \{1, \dots, r\}$  and the equalities (13), (16) are valid for every  $s_i \in \{1, \dots, n_i - m_i - 1\}$  and  $s_p \in \{0, 1, \dots, n_p - m_p - 1\}$ , respectively. The fulfillment of the equality (13), where  $c = a_{i0}/b_{i0}$ , for  $s_i \in \{1, \dots, n_i - m_i - 1\}$ , is equivalent to the fulfillment of the equality

$$\|a_{i0} \ a_{i1} \ \dots \ a_{i, n_i - m_i - 1}\| = c \|b_{i0} \ b_{i1} \ \dots \ b_{i, n_i - m_i - 1}\|$$

for arbitrary  $i \in \{1, \dots, r\}$ . The equality (16) for  $s_p \in \{0, 1, \dots, n_p - m_p - 1\}$  is valid if and only if when

$$\frac{1}{a_p^2} \|a_{p0} \ a_{p1} \ \dots \ a_{p, n_p - m_p - 1}\| = \frac{1}{c b_p^2} \|b_{p0} \ b_{p1} \ \dots \ b_{p, n_p - m_p - 1}\|$$

for arbitrary  $p \in \{r+1, \dots, q\}$ . □

**Corollary 4.2.** *Let the partition (2) of the set  $M$  of characteristic roots of matrices  $A(x)$  be of the form (10). Let also in the notations of Theorem 1.3  $a_i = 0$  for some root  $\alpha_i \in M$  and  $m_j \geq n_j$  for every root  $\alpha_j \in M$ . Such matrix  $A(x)$  in the class  $\{CA(x)Q(x)\}$  of semiscalarly equivalent matrices is determined up to constant factor of its entry  $a(x)$ .*

*Proof.* Let the entry  $b(x)$  of the matrix  $B(x)$  of the form (3) satisfy the condition  $b(\alpha_i) = b_i = 0$  for the same  $\alpha_i \in M$  for which  $a_i = 0$ . By Proposition 1.1, if matrices  $A(x)$  and  $B(x)$  are semiscalarly equivalent, then the equivalence  $a_i = 0 \Leftrightarrow$

$b_l = 0$  is valid for arbitrary  $\alpha_l \in M$ . Moreover, by (the same) Proposition 1.1, all non-zero values of the polynomial  $b(x)$  (as  $a(x)$ ) at the points of the set  $M$  are equal, i.e. for arbitrary pair  $\alpha_p, \alpha_q \in M$  such that  $a_p, a_q \neq 0$  we have  $a_p = a_q \neq 0$  and  $b_p = b_q \neq 0$ . We take into consideration that by Proposition 1.2,  $a^{(t)}(\alpha_j) = b^{(t)}(\alpha_j) = 0, t \in \{1, \dots, n_j - 1\}$ , for every  $\alpha_j \in M$ . From what has been said it follows that there exists a number  $c \in \mathbb{C}$  such that every characteristic root  $\alpha_j \in M$  of multiplicity  $n_j$  is the root of the polynomial  $\varphi(x) = a(x) - cb(x)$  of the same multiplicity  $n_j$ . Taking into account that  $\deg a(x), \deg b(x) < \deg \delta(x)$ , we conclude that  $\varphi(x) \equiv 0$  and thus  $a(x) = cb(x)$ .  $\square$

**Remark.** The conditions of Corollary 4.2 hold for any matrix  $A(x)$  of the form (1) without multiple characteristic roots, if its entry  $a(x)$  takes on the set of characteristic roots exactly two different values, one of which is zero.

## 5. Examples

**Example 5.1.** Consider the matrices

$$A(x) = \begin{vmatrix} 1 & 0 \\ x^2 - x^3 + 4x^4 - 3x^5 & x^4 - 4x^5 + 6x^6 - 4x^7 + x^8 \end{vmatrix}, \\
 B(x) = \begin{vmatrix} 1 & 0 \\ x^2 - x^3 + \frac{21}{4}x^4 - \frac{45}{4}x^5 + \frac{37}{4}x^6 - \frac{11}{4}x^7 & x^4 - 4x^5 + 6x^6 - 4x^7 + x^8 \end{vmatrix}.$$

Each of the characteristic roots  $\alpha_1 = 0, \alpha_2 = 1$  of these matrices are of multiplicity 4. The decompositions of entries

$$a(x) = x^2 - x^3 + 4x^4 - 3x^5, \quad b(x) = x^2 - x^3 + \frac{21}{4}x^4 - \frac{45}{4}x^5 + \frac{37}{4}x^6 - \frac{11}{4}x^7$$

into degrees of binomial  $x - 1$  are of the form:

$$a(x) = 1 - 8(x - 1)^2 - 15(x - 1)^3 - 11(x - 1)^4 - 3(x - 1)^5, \\
 b(x) = \frac{1}{2} - 2(x - 1)^2 - \frac{15}{4}(x - 1)^3 - \frac{17}{2}(x - 1)^4 - \frac{27}{2}(x - 1)^5 - 10(x - 1)^6 - \frac{35}{2}(x - 1)^7.$$

By Corollary 4.1, the matrices  $A(x), B(x)$  are semiscalarly equivalent. In fact,

$$A(x) = \begin{vmatrix} 1 & -1 \\ 0 & 1 \end{vmatrix} \cdot B(x) \cdot \\
 \cdot \begin{vmatrix} 1 + x^2 - x^3 + 4x^4 - 3x^5 & x^4 - 4x^5 + 6x^6 - 4x^7 + x^8 \\ -\frac{9}{4} - \frac{5}{4}x - x^2 + \frac{23}{4}x^3 - \frac{33}{4}x^4 & 1 - x^2 + x^3 - \frac{21}{4}x^4 + \frac{45}{4}x^5 - \frac{37}{4}x^6 + \frac{11}{4}x^7 \end{vmatrix}.$$

**Example 5.2.** Consider the matrix

$$A(x) = \begin{vmatrix} 1 & 0 \\ 2 - 3x + x^3 & 1 - 2x^2 + x^4 \end{vmatrix}$$

with characteristic roots  $\alpha_1 = 1$ ,  $\alpha_2 = -1$ , each of multiplicity 2. The decompositions of entry  $a(x) = 2 - 3x + x^3$  into degrees of binomials  $x - 1$ ,  $x + 1$  are of the form:

$$a(x) = 3(x - 1)^2 + (x - 1)^3 = 4 - 3(x + 1)^2 + (x + 1)^3.$$

Matrix  $A(x)$  satisfies the conditions of Corollary 4.2. Therefore, for matrices  $A(x)$  and  $B(x) = \begin{vmatrix} 1 & 0 \\ b(x) & 1 - 2x^2 + x^4 \end{vmatrix}$ , where  $b(1) = 0$ , to be semiscalarly equivalent, it is necessary and sufficient that  $a(x) = cb(x)$ , where  $c \in \mathbb{C}$ ,  $c \neq 0$ .

**Remark.** Assume that the matrices  $A(x)$  and  $B(x)$  from Theorem 1.3 are semiscalarly equivalent, i.e.  $SA(x)R(x) = B(x)$ . In this case Theorem 1.3 provides a method for constructing the transforming matrices  $S$ ,  $R(x)$ . Such method was used in Example 5.1. However, this is the subject of another study.

## REFERENCES

1. P.S. Kazimirs'kii, V.M. Petrychkovych, *On the equivalence of polynomials matrices*, Theoretical and Applied Problems in Algebra and Differential Equations. Naukova Dumka, Kyiv, (1977), 61–66 (in Ukrainian).
2. J.A. Dias da Silva, T.J. Laffey, *On simultaneous similarity of matrices and related questions*, Linear Algebra Appl. **291** (1999), 167–184.
3. P.S. Kazimirs'kii, *Factorization of Matrix Polynomials*, Naukova Dumka, Kyiv, (1981) (in Ukrainian).
4. L. Baratchart, *Un theoreme de factorisation et son application a la representation des systemes cyclique causaux*, C. R. Acad. Sci. Paris, Ser. 1, Math., **295**:3 (1982), 223–226.
5. B.Z. Shavarovskii, *On invariants and canonical form of matrices of second order with respect to semiscalar equivalence*, BASM. **80**:3 (2016), 12–23.

---

Received 07.04.2017

Revised 20.12.2017