

Hormonal response to different rest intervals during resistance training with light loads

Payam Mohamad-Panahi¹, Hadi Rohani², Navid Lotfi^{*1}

Islamic Azad University, Ghorveh, Iran¹

Sport Science Research Institute of Iran, Mashhad, Iran²

Annotations:

Purpose: The purpose of the present study was to determine the appropriate rest time between sets during weight training with light load. **Material:** Seventeen cadet wrestlers (age =16.7±0.6 yrs.; height =169.2±8.2 cm; and weight =51.4±7.9 kg) were recruited from wrestling clubs in the Iranian province of Kurdistan and served as subjects in this study. This study was conducted over seven sessions with 48 hours recovery between sessions. In the first session, the characteristic features of subjects were recorded and the one repetition maximum in the bench press test was determined for each subject. On 6 separate occasions, subjects performed a 4 set of bench press at 60% 1RM with a 90 and 240 seconds rest interval until volitional fatigue. The numbers of repetition performed by the subjects, and also, cortisol and testosterone levels and 1RM were recorded. The results showed that there was a significant difference in the sustainability of repetitions during 4 sets bench press with 60 % load between 90 and 240 seconds rest intervals (rest interval effect) ($p<0.05$) as well as with 90% load. **Results:** Additionally, there was a significant difference in the sustainability of repetitions during 4 sets bench press in 90 and 240 seconds rest intervals, both, between light and heavy loads (load effect). Plasma cortisol concentrations significantly increased after all bench press trials. Also, the rest interval effect was statistically significant in both 60 % and 90% load trials. But, the load effect was only statistically significant in 90 seconds rest interval trial ($p<0.05$). In contrast, plasma testosterone concentrations significantly increased after 4 sets bench press only in 90 seconds rest interval with heavy load and 240 seconds rest interval with light load ($p<0.05$). Accordingly, testosterone to cortisol (T:C) ratio were significantly decreased after 4 sets bench press in 90 seconds rest interval with light load and 240 seconds rest interval with heavy load. Also, load effect was statistically significant only in 90 seconds rest interval trials and rest interval effect was statistically significant only in heavy load trials ($p<0.05$). **Conclusions:** Based on these results it can be concluded that when resistance training with light loads, higher resting interval (240 seconds) may be provides a better anabolic environment for muscle growth and keep the training intensity.

Key words:

rest interval, training load, resistance training.

Паям Мохаммад-Панахі, Хаді Рохані, Навід Лотфі. Гормональні реакції на різні інтервали відпочинку під час силових тренувань з невеликими навантаженнями. **Мета:** визначити оптимальний час відпочинку між підходами під час силового тренування з легким навантаженням. **Матеріал:** 17 кадетів борців (вік 16,7 ± 0,6 років; зрост 169,2 ± 8,2 см; вага 51,4 ± 7,9 кг) були набрані з борцівських клубів в іранській провінції Курдистан і служили в якості суб'єктів в даному дослідженні. Дане дослідження було проведено протягом семи сесій з 48 годинним відновленням у період між сесіями. У першій сесії були записані індивідуальні показники спортсменів і був визначений для кожного суб'єкта максимальний показник одного повторення в стенових випробуваннях. У 6 окремих випадках суб'єкти 4 рази не виконали жим лежачи (60 % від 1 максимального повторення, від 90 і до 240 секунд – інтервал відпочинку) до вольової втоми. Число повторень кожного спортсмена, а також рівень кортизолу і тестостерону були записані. **Результати:** результати показали, що існує значна різниця у стійкості повторень протягом 4 підходів з 60 % навантаженням і інтервалом відпочинку від 90 до 240 секунд (інтервал відпочинку ефективний, $p < 0,05$), а також з 90 % навантаженням. Крім того, існує значна різниця у стійкості повторень протягом 4 підходів у жимі лежачи з інтервалом відпочинку від 90 і до 240 секунд між легкими і великими навантаженнями (ефект навантаження). Концентрація кортизолу значно збільшується після всіх прес – стенових випробувань. Крім того, в інших інтервалах ефект був статистично значущим в обох випробуваннях навантаження 60 % і 90 %. Але, ефект навантаження був статистично значущим тільки при 90 секундній перерві ($p < 0,05$). На відміну від цього, концентрація тестостерону значно зросла після 4 підходів у жимі лежачи при 90 секундному інтервалі відпочинку при великих навантаженнях і при 240 секундному інтервалі відпочинку при невеликому навантаженні ($p < 0,05$). Відповідно, коефіцієнт співвідношення тестостерон – кортизол (Т : С) значно скоротився після 4 підходів у жимі лежачи при 90 секундному інтервалі відпочинку при невеликому навантаженні і при 240 секундному інтервалі відпочинку з великим навантаженням. Крім того, ефект навантаження був статистично значущим тільки при 90 секундному відпочинку. Співвідношення інтервал і відпочинку ефект був статистично значущим тільки в тестах з великим навантаженням ($p < 0,05$). **Висновки:** на підставі цих результатів можна зробити висновок, що при тренуваннях з опором у вигляді невеликих навантажень інтервал відпочинку в 240 секунд і вище забезпечує кращу анаболічне середовище для росту м'язів і підтримки інтенсивності тренування.

інтервал відпочинку, тренувальна навантаження, тренувальна протидія.

Паям Мохаммад-Панахи, Хаді Рохані, Навід Лотфі. Гормональные реакции на разные интервалы отдыха во время силовых тренировок с небольшими нагрузками. **Цель:** определить оптимальное время отдыха между подходами во время силовой тренировки с легкой нагрузкой. **Материал:** 17 кадетов борцов (возраст 16,7 ± 0,6 лет; рост 169,2 ± 8,2 см; вес = 51,4±7,9 кг) были набраны из борцовских клубов в иранской провинции Курдистан и служили в качестве субъектов в данном исследовании. Данное исследование было проведено в течение семи сессий с 48 часовым восстановлением в период между сессиями. В первой сессии были записаны индивидуальные показатели спортсменов и был определен для каждого субъекта максимальный показатель одного повторения в стеновых испытаниях. В 6 отдельных случаях субъекты 4 раза не выполнили жим лежа (60% от 1 максимального повторения, от 90 и до 240 секунд – интервал отдыха) до волевой усталости. Число повторений каждого спортсмена, а также уровень кортизола и тестостерона были записаны. **Результаты:** результаты показали, что существует значительное различие в устойчивости повторений в течение 4 подходов с 60%-ной нагрузкой и интервалом отдыха от 90 до 240 секунд (интервал отдыха эффективный, $p < 0,05$), а также с 90%-ной нагрузкой. Кроме того, существует значительная разница в устойчивости повторений в течение 4 подходов в жиме лежа с интервалом отдыха от 90 и до 240 секунд между легкими и большими нагрузками (эффект нагрузки). Концентрация кортизола значительно увеличивается после всех прес- стеновых испытаний. Кроме того, в остальных интервалах эффект был статистически значимым в обоих испытаниях нагрузка 60 % и 90%. Но, эффект нагрузки был статистически значимым только при 90 секундном перерыве ($p < 0,05$). В отличие от этого, концентрация тестостерона значительно возросла после 4 подходов в жиме лежа при 90 секундном интервале отдыха при больших нагрузках и при 240 секундном интервале отдыха при небольшой нагрузке ($p < 0,05$). Соответственно, коэффициент соотношения тестостерон – кортизол (Т : С) значительно сократился после 4 подходов в жиме лежа при 90 секундном интервале отдыха при небольшой нагрузке и при 240 секундном интервале отдыха с большой нагрузкой. Кроме того, эффект нагрузки был статистически значимым только при 90 секундном отдыхе. Соотношение интервал и отдых эффект был статистически значимым только в тестах с большой нагрузкой ($p < 0,05$). **Выводы:** на основании этих результатов можно сделать вывод, что при тренировках с сопротивлением в виде небольших нагрузок интервал отдыха в 240 секунд и выше обеспечивает лучшую анаболическую среду для роста мышц и поддержания интенсивности тренировки.

интервал отдыха, тренировочная нагрузка, тренировочное противодействие.

Introduction

In recent years, one of the important issues that have attracted the attention of many researchers is the optimum recovery time in exercise training and sports competition. Due to the nature of the sports and different needs of fitness, there is a specific training method for obtaining the necessary preparations for any sport [4]. Among the training factors, the rest interval time between workouts is a key element of success in any training program [3].

Recovery or return to the base state is the time between the end of a workout or strenuous activity and the next activity or in other words, the duration between the end of one activity and the start next break [15].

Due to the great diversity of sports, the optimal recovery time varies between different fields. Therefore, athletes should train with different methods based on the nature of their sport and the need for strength, endurance and speed.

Weight training in many sports is important for improving physical fitness factors. Hence, athletes to achieve the desired level of performance and increase overall physiological abilities, should use strength-endurance training. According to many studies in the past two decades, rest interval and a proper recovery time between activities is one of the main factors for the success of any training program [6, 15, 17]. In the absence of adequate rest and recovery, athletes will not benefit from the full effect of exercise.

There has been enough information about the optimal weights, repetitions and sets in weight training programs. Much research was done in the area of weight training based on principles of overload, resistance and features. De Salles et al (2009) studied the rest interval between sets in strength training [6]. They reported that resting 3-5 minutes between sets produced greater increases in absolute strength, due to higher intensities and volumes of training. Furthermore, they reported that higher levels of muscular power were demonstrated over multiple sets with 3 or 5 minutes versus 1 minute of rest between sets. Jensen and Ebben (2003) studied the kinetic analysis of complex training rest interval effect on vertical jump performance [9]. They showed that to optimize jump performance it appears that athletes should not perform jumps immediately following resistance training. It may be possible that beyond 4 minutes of recovery performance could be enhanced. Robinson et al (1995) investigated the effects of different weight training exercise/rest intervals on strength, power, and high-intensity exercise endurance [20]. They showed that except for maximum strength, adaptations, to short-term, high-volume training may not be dependent on the length of rest intervals. Matuszak et al (2003) studied the effect of rest interval length on repeated 1 repetition maximum back squats [13]. They reported that indicate that 1-minute rest intervals are sufficient for recovery between attempted lifts during 1RM testing or training for the free-weight back squat when involving lifters of this caliber. Willardson et al

(2006) investigated the effect of rest interval length on bench press performance with heavy vs. light loads [25]. They showed that when the training goal is the maximal strength development, 3 minutes of rest should be taken between sets to avoid significant declines in repetitions. The ability to sustain repetitions while keeping the intensity constant may result in a higher training volume and consequently greater gains in muscular strength. Rahimi (2005) studied the effect of different rest intervals on the exercise volume completed during squat bouts [17]. He reported that the 5-minute rest condition resulted in the highest volume completed, followed in descending order by the 2- and 1-minute rest conditions. The ability to perform a higher volume of training with a given load may stimulate greater strength adaptations.

In recent years, several studies have examined the effect of exercise loads, number of repetitions and sets discussed. However, there is no general consensus about rest periods between sets. Furthermore, the period of rest between weight training sessions are different in strength and endurance sports. Therefore, the purpose of the present study was to determine the appropriate rest time between sets during weight training with light load.

Methods

Subjects

Seventeen cadet wrestlers were recruited from wrestling clubs in the Iranian province of Kurdistan and served as subjects in this study. They all had at least 3 years training experience and were representative of the top wrestlers of Kurdistan competing in national competitions. Before participating, the subjects' parents were informed of the potential risks and gave their written informed consent for their children to participate in this study, which was consistent with the human subject policy of the University of Guilan Research Center. Subject characteristics were as follows (mean \pm SD): age = 16.7 \pm 0.6 yr; height = 169.2 \pm 8.2 cm; and weight = 51.4 \pm 7.9 kg.

This study was conducted over seven sessions with 48 hours recovery between sessions. In the first session, the characteristic features of subjects were recorded and the one repetition maximum in the bench press test was determined for each subject. On 6 separate occasions, subjects performed a 4 set of bench press at 60% 1RM with a 90 and 240 seconds rest interval until volitional fatigue.

The numbers of repetition performed by the subjects were recorded. 5 ml blood was drawn from the antecubital vein before and 5 minutes after exercise and transferred immediately to the lab for assessing blood testosterone and cortisol levels. Serum levels of testosterone and cortisol were measured by radioimmunoassay (RIA) using the commercial kit (IM1119, IMMUNOTECH) and (IMMUNOTECH, IM1841). The one-repetition maximum (1RM) of subject measured by bench press test and Brzycki formula:

$$1RM = \text{weight (kg)} \times (1.0278 - (0.0278 \times \text{repetition})) \quad [18].$$

Statistical Methods

All descriptive data are expressed as means \pm SD. Data were analyzed using ANOVA with repeated measure

and Bonferroni post hoc test. Statistical analysis was conducted using SPSS 16.0 for Windows.

Results

The results of sustainability of repetition, Cortisol concentrations, Testosterone concentrations, and Testosterone to Cortisol (T: C) ratio during 4 sets bench press in different trails are shown in Fig 1. There was a significant difference in the sustainability of repetitions during 4 sets bench press with 60 % load between 90 and 240 seconds rest intervals ($p < 0.05$; Fig 1-A).

Plasma cortisol concentrations significantly increased after both bench press trials. Also, the rest interval effect was statistically significant in 60 % load trial ($p < 0.05$; Fig 1-B). In contrast, plasma testosterone concentrations significantly increased after 4 sets bench press only in 240 seconds rest interval with light load ($p < 0.05$; Fig 1-C). Accordingly, testosterone to cortisol (T:C) ratio were significantly decreased after 4 sets bench press in 90 seconds rest interval ($p < 0.05$; Fig 1-D).

Discussion

One of the experimental methods to evaluate the physically and mentally state of athletes is the investigation of the anabolic and catabolic hormones changes during exercise. In this context, the role of cortisol as a primary catabolic hormone is important. It has been suggested that cortisol changes in response to exercise and, mental and physical stress [12].

Our results showed that plasma cortisol concentrations increased significantly after 4 sets bench press exercise with 90 and 240 seconds rest intervals between sets with


light and heavy loads. Interestingly, mean changes were significantly higher in 90 seconds rest interval with 60% load in compare to 90% load.

It is reported that changes in core temperature during exercise can affect plasma cortisol levels and there is a direct correlation between core temperature and cortisol levels [5]. Furthermore, other researchers showed that exercise is a powerful stimulant for the central nervous system (CNS) and hypothalamic-pituitary-adrenal axis [1].

Moreover, other authors reported that increase in core temperature during exercise via stimulating hypothalamic-pituitary-adrenal axis increases cortisol values [23]. If you don't get sufficient rest between workouts, this hormone will change in a way that puts your body in a catabolic state. Cortisol is a catabolic and steroid hormone which is largely involved in regulation of metabolism and the body's response to exercise stress. However, long-term elevation will cause the problems that immune system disorders and protein degradation are most important of these [24].

Cortisol changes during exercise training are explained by different mechanisms. In the high intensity exercise, a 2-fold increase in plasma cortisol is observed, which is due to the increased hormone secretion rather to the excretion [10].

The intensity and duration of physical activity are important factors that affect cortisol response. In the exercise with intensity of higher than 60 % maximum oxygen consumption, the levels of this hormone is increased [10].



Although, more increase in cortisol level in 240 seconds in compare to 90 seconds rest interval is unaccountable in heavy load trials, using light load, lower the rest interval resulted in higher cortisol levels in current study. Many factors such as stress, physiological circadian rhythm and body temperature affects on the acute cortisol response to exercise [10].

We observed that mean plasma testosterone concentrations do not change significantly after 4 sets bench press with light load and 90 seconds rest interval, as similar as, with heavy load and 240 seconds rest interval. In contrast, as mentioned above, cortisol level has significantly increased in these trials. Since cortisol is the inhibitor of testosterone secretion and the level of this hormone increased, it is expected that in this rest intervals, testosterone does not change significantly [7].

On the other hand, it is known that skeletal muscle fibers have specific androgen receptors that can remove these hormones from the circulation and cleansing by the increase in the metabolism of these. Therefore, in the present study, may be increased removal of testosterone by muscle result in increased insignificant testosterone [2, 21].

As a result of cortisol and testosterone changes, the testosterone to cortisol ratio, a criteria for balance of anabolic-catabolic processes [14, 22], significantly changed in those trials that testosterone has no significant changes, i.e. light load with 90 s rest interval and heavy load with 240 s rest interval.

Mechanisms underlying testosterone changes following physical activity is not well understood. However, one of the possible mechanisms that proposed for testosterone changes is shifts in blood volume [16]. If blood volume reduced during physical activity is likely that increased blood viscosity and increase hormone was unreal.

Based on the intensity and duration of physical activity, temporary changes occur in the balance of

anabolic-catabolic processes. Repetitive exercises without adequate and appropriate recovery can cause prolonged irregularities in the balance of anabolic-catabolic processes [8, 11]. Furthermore, this ratio decreases with increasing intensity and duration of exercise [8].

The results showed that the testosterone to cortisol ratio has significantly changed after 4 sets of bench press with 90 seconds rest interval. Although, increased the levels of testosterone and cortisol, both, the testosterone cortisol ratio was significantly reduced due to higher levels of cortisol in compare to testosterone.

Finally, the ability to maintain repetitions of bench press is higher in long vs. short rest intervals. It is possible that insufficient time for recovery and reconstruction energy stores and waste disposal in the short rest intervals is the main cause. It can be concluded that in the situations that muscle hypertrophy is a goal; it can be obtained by assigning a long time rest interval and heavy load in resistant exercise training.

Overall, the results indicated that the Sustainability of repetition, Testosterone concentration and testosterone to cortisol ratio in training with light load and rest intervals of 240 seconds was more than 90 seconds. Also, the cortisol concentration in 240 seconds resting interval was less than 90 seconds resting interval. Based on these results it can be concluded that when resistance training with light loads, higher resting interval (240 seconds) may be provides a better anabolic environment for muscle growth and keep the training intensity.

Acknowledgments

The authors wish to thank the Vice Chancellor for Research of Islamic Azad university of Ghorveh branch for their finance assistants and athletes and coaches of Takhti wrestling club who participated in present study. This work was supported by the Islamic Azad university of Ghorveh branch.

References

1. Avois L., N. Robinson et al. Central nervous system stimulants and sport practice. *British journal of sports medicine*. 2006, vol.40, pp. 116-120.
2. Azarbaijani M.A., Nikbakht H., Rasae M.J., Sabeti K.h. Effect of exhaustive incremental exercise session on salivary testosterone and cortisol in wrestlers. *The Journal of Applied Sport Science Research*. 2002, vol.4, pp. 101-114.
3. Baechle T.R., Earle R.W. *Weight training: steps to success*, Human Kinetics. 2011, 124 p.
4. Berger R. Effect of varied weight training programs on strength. *Research Quarterly*. American Association for Health. *Physical Education and Recreation*. 1962, vol.33(2), pp. 168-181.
5. Brenner I., Zamecnik J. The impact of heat exposure and repeated exercise on circulating stress hormones. *European journal of applied physiology and occupational physiology*. 1997, vol.76(5), pp. 445-454.
6. De Salles B.F., Miranda F. Rest interval between sets in strength training. *Sports Medicine*. 2009, vol.39(9), pp. 765-777.
7. Doerr P., Pirke K.M. Cortisol-induced suppression of plasma testosterone in normal adult males. *The Journal of Clinical Endocrinology and Metabolism*. 1976, vol.43(3), pp. 622-629.
8. Eliakim A., Nemet D. Exercise training, physical fitness and the growth hormone-insulin-like growth factor-1 axis and cytokine balance. *Medicine and Sport Science*. 2010, vol.55, pp. 128-40. doi: 10.1159/000321977
9. Jensen R.L., Ebben W.P. Kinetic analysis of complex training rest interval effect on vertical jump performance. *The Journal of Strength & Conditioning Research*. 2003, vol.17(2), pp. 345-349.
10. Kraemer W.J. Endocrine responses to resistance exercise. *Medicine and Sport Science*. 1988, vol.20(5), pp. 152-157.
11. Kraemer W.J., Volek J.S. Hormonal responses to consecutive days of heavy-resistance exercise with or without nutritional supplementation. *Journal of Applied Physiology*. 1998, vol.85(4), pp. 1544-1555.
12. Lovallo W.R., Farag N.H. Cortisol responses to mental stress, exercise, and meals following caffeine intake in men and women. *Pharmacology Biochemistry and Behavior*. 2006, vol.83(3), pp. 441-447.
13. Matuszak M.E., FRY A.C. Effect of rest interval length on repeated 1 repetition maximum back squats. *The Journal of Strength & Conditioning Research*. 2003, vol.17(4), pp. 634-637.
14. Obmiński Z., Stupnicki R. Comparison of the testosterone-to-cortisol ratio values obtained from hormonal assays in saliva and serum. *The Journal of sports medicine and physical fitness*. 1997, vol.37(1), pp. 50-55.
15. Pincivero D.M., Lephart S. M. Effects of intrasession rest interval on strength recovery and reliability during high intensity exercise. *The Journal of Strength & Conditioning Research*. 1998, vol.12(3), pp. 152-156.

16. Raastad T., Bjørø T. Hormonal responses to high-and moderate-intensity strength exercise. *European journal of applied physiology*. 2000, vol.82(1-2), pp. 121-128.
17. Rahimi R. Effect of different rest intervals on the exercise volume completed during squat bouts. *Journal of Sports Science and Medicine*. 2005, vol.4(4), pp. 361-366.
18. Rajabi A., Lotfi N., Abdolmaleki A., Rashid-Amiri S.H. The effects of omega-3 intake on delayed onset muscle soreness in non-athlete men. *Pedagogics, Psychology, Medical-Biological Problems of Physical Training and Sports*. 2013, vol.1, pp. 91-95.
19. Ratamess N. A., Falvo M. J. The effect of rest interval length on metabolic responses to the bench press exercise. *European journal of applied physiology*. 2007, vol.100(1), pp. 1-17.
20. Robinson J. M., Stone M. H. Effects of different weight training exercise/rest intervals on strength, power, and high intensity exercise endurance. *The Journal of Strength & Conditioning Research*. 1995, vol.9(4), pp. 216-221.
21. Snochowski M., Saartok T., Dahlberg E., Eriksson E., Gustafsson J.A. Androgen and glucocorticoid receptors in human skeletal muscle cytosol. *Journal of Steroid Biochemistry*. 1981, vol.14(8), pp. 765-771.
22. Vervoorn C., Quist A. The behaviour of the plasma free testosterone/cortisol ratio during a season of elite rowing training. *International journal of sports medicine*. 1991, vol.12(03), pp. 257-263.
23. Vilmon J.H., Kastil D.L. *Sport Physiology and Physical Activity*. Human kinetics USA. 2008, 200 p.
24. Vitetta L., Anton B. Mind Body Medicine: Stress and Its Impact on Overall Health and Longevity. *Annals of the New York Academy of Sciences*. 2005, vol.1057(1), pp. 492-505.
25. Willardson J.M., Burkett L.N. The effect of rest interval length on bench press performance with heavy vs. light loads. *The Journal of Strength & Conditioning Research*. 2006, vol.20(2), pp. 396-399.

Information about the authors:

Payam Mohamad-panahi: ORCID: 0000-0002-9204-8882; Panahi. payam@gmail.com; Islamic Azad University; Janbazan Sq, 66619-83435, p.o.Box: 161, Ghorveh, Iran

Hadi Rohani: ORCID: 0000-0002-0437-753X; h_Rohani7@yahoo.com; Sport Science Research Institute of Iran; Ladan Blvd, Vakil Abad 54, Mashhad, Iran

Navid Lotfi: ORCID: 0000-0002-1366-5058; navid_lotfi2008@yahoo.com; Islamic Azad University; Janbazan Sq, 66619-83435, p.o.Box: 161, Ghorveh, Iran

Cite this article as: Payam Mohamad-Panahi1, Hadi Rohani2, Navid Lotfi. Hormonal response to different rest intervals during resistance training with light loads. *Pedagogics, psychology, medical-biological problems of physical training and sports*, 2014, vol.2, pp. 67-71. doi:10.6084/m9.figshare.923517

The electronic version of this article is the complete one and can be found online at: <http://www.sportpedagogy.org.ua/html/arhive-e.html>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited (<http://creativecommons.org/licenses/by/3.0/deed.en>).

Received: 28.12.2013
Published: 28.12.2013

Информация об авторах:

Паям Мохамад-панахи: ORCID: 0000-0002-9204-8882; Panahi. payam@gmail.com; Исламский Азад Университет; Джанбазан сквер, 66619-83435, п/я 161, Гхорвех, Иран

Хади Рохани: ORCID: 0000-0002-0437-753X; h_Rohani7@yahoo.com; Научно-исследовательский институт спорта Ирана; Ладан бул, Вакиль Абад 54, Мешхед, Иран

Навид Лотфи: ORCID: 0000-0002-1366-5058; navid_lotfi2008@yahoo.com; Исламский Азад Университет; Джанбазан сквер, 66619-83435, п/я 161, Гхорвех, Иран

Цитируйте эту статью как: Паям Мохамад-Панахи, Хади Рохани, Навид Лотфи. Гормональные реакции на разные интервалы отдыха во время силовых тренировок с небольшими нагрузками // Педагогіка, психологія та медико-біологічні проблеми фізичного виховання і спорту. – 2014. – № 2 – С. 67-71. doi:10.6084/m9.figshare.923517

Электронная версия этой статьи является полной и может быть найдена на сайте: <http://www.sportpedagogy.org.ua/html/arhive.html>

Эта статья Открытого Доступа распространяется под терминами Creative Commons Attribution License, которая разрешает неограниченное использование, распространение и копирование любыми средствами, обеспечивающими должное цитирование этой оригинальной статьи (<http://creativecommons.org/licenses/by/3.0/deed.ru>).

Дата поступления в редакцию: 28.12.2013 г.
Опубликовано: 28.12.2013 г.