

Characteristics of body posture in the sagittal plane and fitness of first-form pupils from rural areas

Żukowska Hanna¹, Szark-Eckardt Mirosława¹, Muszkieta Radosław¹, Iermakova Tetiana²

Kazimierz Wielki University, Bydgoszcz, Poland¹

G.S. Skovoroda Kharkiv National Pedagogical University, Ukraine²

Annotation:

Purpose: to find correlations between characteristics of body posture in the sagittal plane and fitness and endurance of first-form children from rural areas.

Material: an analysis of more than 30 sources of scientific and educational literature. **Results:** the study involved 209 children, including 102 girls and 107 boys. They were children who lived in the country since they were born. To assess particular characteristics of body posture, the children were studied by means of the measuring equipment using the projection Moiré system. Motor skills were estimated using selected EUROFIT physical fitness tests (sitting forward bend, standing broad jump, handgrip, sit-and-reach, bent arm hang and 10 x 5 m shuttle run). The level of physical endurance was evaluated with the Harvard Step Test modified by Montoye.

Conclusions: the conducted research reveals statistically significant correlations between the characteristics of body posture in the sagittal plane and selected EUROFIT physical fitness tests and physical endurance of the children involved in the study.

Keywords:

EUROFIT, correlations, physical endurance, circulatory system, spinal.

Жуковска Анна, Шарк-Ецкардт Мирослава, Мушкета Радослав, Ермакова Татьяна. Характеристики положения тела в сагиттальной плоскости и физическое состояние первоклассников из сельской местности. Цель: определить корреляции между характеристиками осанки в сагиттальной плоскости и физической подготовке и выносливости первоклассников из сельских районов.

Материал: проанализировано более 30 источников научной и учебной литературы. **Результаты:** в исследовании приняли участие 209 детей, из них 102 девочки и 107 мальчиков. Дети, которые жили в деревне с рождения. Для оценки конкретных характеристик осанки, дети были исследованы с помощью измерительного оборудования с использованием системы проекция Muar. Двигательные навыки были оценены с использованием выбранных Eurofit физических тестов (сидя наклон вперед, стоя прыжок в длину, хват рукой, гибкость, челночный бег 10 x 5 м с согнутыми в локтях руками). Уровень физической выносливости оценивали с помощью Гарвард степ тест модифицированного Монтойем.

Выходы: проведенное исследование показывает статистически значимые корреляции между характеристиками осанки в сагиттальной плоскости и выбранным Eurofit физическим тестом испытаний, а также физической выносливости детей, участвующих в исследовании.

Eurofit, корреляции, физическая выносливость, система кровообращения, спинной.

Жуковска Анна, Шарк-Ецкардт Мирослава, Мушкета Радослав, Ермакова Татьяна. Характеристики положения тела в сагиттальной плоскости и физический стан первоклассников из сельской местности. **Мета:** визначити кореляції між характеристиками постави в сагітальній площині і фізичного стану і витривалості першокласників з сільської місцевості.

Матеріал: проаналізовано понад 30 джерел наукової та навчальної літератури. **Результати:** у дослідженні взяли участь 209 дітей, з них 102 дівчинки і 107 хлопчиків. Діти, які жили у селі з народження. Для оцінки конкретних характеристик постави, діти були досліджені за допомогою вимірювального обладнання з використанням системи проекції Muar. Рухові навички були оцінені з використанням вибраних Eurofit фізичних тестів (сидячи нахил вперед, стоячи стрібок у джокину, хват рукою, гнучкість, човниковий біг 10 x 5 м із зігнутими в ліктях руками). Рівень фізичної витривалості оцінювали за допомогою Гарвард степ тест модифікованого Монтойєм.

Висновки: проведене дослідження показує статистично значущі кореляції між характеристиками постави в сагітальній площині і вибраним Eurofit фізичним тестом випробувань, а також фізичної витривалості дітей, що беруть участь в дослідженні.

Eurofit, кореляції, фізична витривалість, система кровообігу, спинний.

Introduction

The current generation of children and teenagers is becoming more robust as Przeweda and Dobosz report [24]. As far as physical development is concerned, an increase in the body height and weight is observed in the young generation. However, as the authors report, the morphological development and its acceleration do not go together with acceleration of physical endurance and fitness. Visible regress can be observed within these indicators [24], which is especially noticeable among rural teenagers and in families of lower social status [17, 24, 25]. Numerous researchers [2, 9, 16, 17, 24, 27, 30] emphasise the influence of environment on the level of biological development and health condition of children and teenagers. As the authors suggest, this decrease in physical endurance and fitness can be also caused by the constant development of mechanization, urbanization and technology. Civilization progress facilitates everyday life but it also reduces people's physical exercise, what leads to a sedentary lifestyle and, in consequence, to different civilization disorders (e.g. obesity).

Human adaptation mechanisms do not keep up with

© Żukowska Hanna, Szark-Eckardt Mirosława, Muszkieta Radosław, Iermakova Tetiana, 2014

doi: 10.6084/m9.figshare.1015583

the dynamic development of environment. In case of teenagers and children, inability to adjust to a big number of stimuli and lack of movement lead to mental disorders (e.g. neuroses) and cause reduced fitness and endurance. This in turn causes decreased immunity which results in pain troubles of the motor system (mainly pain of the cervical and lumbar spine). Disorders of adaptation mechanism in children and teenagers are revealed, among others, by the "epidemic" of faulty postures and lower extremity postural distortions as numerous studies indicate [6, 10, 11, 20, 22, 24, 29]. Therefore, as far as health promotion is concerned a special role should be played by positive health indicators which may comprise the evaluation criteria of the population health condition. These include physical development, physical activity, physical endurance and fitness, among others [7]. The indicators reflect the work of human organs and systems as well as they require constant diagnosing and monitoring. Hence, significant as it seems to be is that particularly children should be subject to constant control towards positive health indicators. Additionally, the body posture of a child expresses its mental and physical state. Anomalies, distortions and faulty postures have an influence on physiological and motor functions of a human, for example, distortions in the thoracic region negatively

affect the lungs and heart work and, in consequence, they lead to circulatory and respiratory impairment. Defects in the region of knees impair a child's motor activity whereas in the region of feet they reduce a child's support and marching activities [23].

Methods

Children studied

The research involved 270 pupils, but finally, there were used the results of 209 children who participated in all research stages and were given their parents' consent. Having divided the children according to their sex the study involved: 102 girls and 107 boys. The children involved in the study lived in the country since they were born (Table 1).

The research was conducted in three rural schools of the Kuyavia-Pomerania Province: Dąbrowa Chełmińska, Nowa Wieś Wielka and Wojnowo. Children from the neighbouring rural areas were driven to schools.

Research Organisation

The rural schools of the Kuyavia-Pomerania Province which were attended by the studied children were chosen using a random search method [14].

The research involved the evaluation of body posture and assessment of motor skills and physical endurance of the children. The evaluation of body posture was conducted at the Body Posture Laboratory of the AWFiS (University of Physical Education and Sport) in Gdansk whereas the study of motor skills of boys and girls was carried out in sports facilities belonging to schools attended by the children involved in the study. During the study of body posture it was essential for a studied person to assume a free, unconstrained standing position with slightly extended legs, straight knee and hip joints, with arms loosely lowered along the trunk and the head in the Frankfurt plane [19, 31]. The evaluation of body posture was conducted in a properly adapted room (lighting, temperature) in the morning.

Body posture evaluation

The study of body posture was conducted at the Body Posture Laboratory of the AWFiS (University of Physical Education and Sport) in Gdansk using the projection

Moiré system. The applied computer system provided three-dimensional coordinates of the studied regions and calculated the parameters of body posture in the sagittal plane at the same time. Having prepared the data, there were obtained 50 parameters of linear and angular posture dimensions. For the purpose of the study, the following data characterizing body posture in the sagittal plane of the studied rural children were applied: angles of depression of the particular spinal sections (degrees): upper thoracic γ , thoracic-lumbar β , lumbar-sacral α and the angle of pelvic anteversion in the sagittal plane (degrees).

Taking the measured angles into consideration, angular dimensions were assessed and small, big and medium angles were determined [18]. Using the Wolański method completed by Zeyland-Malawka [31] the types of posture were determined. Numeric values were also evaluated for: thoracic kyphosis (χ), lordosis (λ), total spinal curvatures (δ) and compensation index (μ).

In order to standardize terminology for the purpose of this scientific paper, the nomenclature of spinal angles presented by Zeyland-Malawka [31] was used. The calculated size of angle α , which means the depression angle of the lumbar-sacrum section according to the Świerc system, determines in this paper the depression angle of the upper thoracic section, angle β – the depression angle of thoracic-lumbar spine remains the same whereas angle γ , which means the depression angle of the upper thoracic section according to the Świerc system, means the depression angle of the lumbar-sacral section in this publication.

Evaluation of fitness and physical endurance

Fitness was examined using selected tests of the EUROFIT physical fitness test and the tests were conducted in accordance with generally binding rules [8].

To evaluate the fitness level the following tests were applied: sitting forward bend, standing broad jump, handgrip (with the stronger hand), sit-and-reach within 30s, bent arm hang and 10 x 5m shuttle run.

In order to estimate the endurance level of the studied boys and girls, the Harvard Step test was applied, which was modified by Montoye and used in the research study

Table 1

Statistical description of the children involved in the study

PLACE OF RESIDENCE	BOYS		GIRLS		TOTAL	
	n	%	n	%	N	%
Dąbrowa Chełmińska	30	46.9	34	53.1	64	30.6
Nowa Wieś Wielka	63	57.8	46	42.2	109	52.1
Wojnowo	14	38.9	22	61.1	36	17.2
TOTAL	107		102		209	

Table 2

Physical fitness criteria developed by Mazur et al. (1975)

Endurance index	Endurance rating
Above 60	excellent
50.1 – 60.0	good
40.1 – 50.0	average
30.1 – 40.0	low average
Below 30	poor

Source: Mazur et al. (1975)

of Polish children by Mazur et al. [15]. This test also reveals the response of cardiac contraction frequency in children during an exercise and after one-minute rest. This test was determined as the Exercise Test in Step-Test. The exercise test used in the step-test was conducted in accordance with recommendations of Mazur et al. [15] for the Harvard Step Test, adapted for children. The exercise took 5 minutes and during this time the studied person stepped up and down at a rate of 30 steps per minute. The pulse was taken by means of a heart rate monitor Polar – S810i one minute after the exercise.

The endurance index was evaluated and for the analysis purpose there were used the endurance criterion (Table 2) proposed by Mazur et al. [15].

Results

Correlations of the characteristics of body posture in the sagittal plane with motor skills in boys and girls

The results achieved by the studied boys in the bent arm hang test reveal a statistically significant correlation at the assumed level $\alpha = 0.05$ with the upper thoracic angle – α (poor correlation $r = -0.19$ and thoracic-lumbar angle – β (moderate correlation $r = -0.3$). These correlations are opposite (when one variable increases – the duration of the hang, the second one decreases – the alpha angle, the beta angle). There is a poor, but statistically insignificant correlation between the time of hang of the boys involved

in the study and the angle of pelvic anteversion ($r = 0.18$) (Table 3).

Table 4 reveals correlations between the analysed angles in the sagittal plane and the sit-and-reach test (Table 4).

When analysing the findings, there can be defined a statistically significant correlation at the assumed level $\alpha = 0.05$ between the sit-and-reach test and the lumbar-sacral angle – γ (poor correlation $r = -0.26$). This correlation is of an opposite character. Additionally, a low, but statistically insignificant correlation was determined between the sit-and-reach test and the upper thoracic angle – α ($r = -0.13$).

As the analysis of the results revealed in table 5 suggests, it appears that the bend forward test correlates only with the thoracic-lumbar angle – β at a low level and turned out to be statistically insignificant ($r = 0.16$).

Between the shuttle run test and spinal angles in the sagittal plane (Table 6) there was noticed one statistically significant correlation, that is with the lumbar-sacral angle γ (poor correlation $r = 0.23$). This correlation is consistent. Apart from that, low correlation can be noticed between the shuttle run test and the upper thoracic angle – α ($r = 0.18$), the thoracic-lumbar β ($r = 0.1$) and the angle of pelvic anteversion ($r = -0.12$). However, these correlations are lower and statistically insignificant.

The standing broad jump test, at a low level,

Table 3

Correlations occurring between the angles characteristic of anterior-posterior pelvic tilts (upper thoracic - α , thoracic-lumbar - β , sacral-lumbar γ) and the angle of pelvic anteversion in the sagittal plane and the bent arm hang test in the studied boys

Bent Arm Hang (s)		BOYS			Angle of (pelvic) anteversion
		α	β	γ	
r		-0.19	-0.3	-0.05	0.18
Correlation power		<i>poor</i>	<i>average</i>	hardly Any	low
t		1.98	3.2	0.47	1.86
Statistical significance		$\alpha = 0.05$	$\alpha = 0.05$	lack	lack

Table 4

Correlations occurring between the angles characteristic of anterior-posterior pelvic tilts (upper thoracic - α , thoracic-lumbar - β , sacral-lumbar γ) and the angle of pelvic anteversion in the sagittal plane and the sit-and-reach test in the boys involved in the study

Sit-and-Reach (number)		BOYS			Angle of (pelvic) anteversion
		α	β	γ	
r		-0.13	0.04	-0.26	-0.06
Correlation power		<i>poor</i>	hardly any	<i>poor</i>	hardly any
t		1.4	0.42	2.74	0.57
Statistical significance		lack	lack	$\alpha = 0.05$	lack

insignificantly correlates with the upper thoracic angle - α ($r = -0.1$) and the lumbar-sacral angle γ ($r = -0.12$) (Table 7).

While analysing the results revealed in table 8 it appears that there are low, but statistically insignificant correlations between the handgrip test and the upper thoracic angle α ($r = 0.12$) and the angle of pelvic anteversion ($r = 0.13$), likewise, within the correlations

between the Step-test and the spinal angles in the sagittal plane there can be noticed a low ($r = -0.18$), statistically insignificant correlation with the upper thoracic angle - α (Table 8, 9).

The results achieved by the girls involved in the study in the bent arm hang test do not significantly correlate at the assumed level $\alpha = 0.05$ with any of the independent variables. Poor, but statistically insignificant correlations

Table 5
Correlations occurring between angles characteristic of anterior-posterior pelvic tilt (upper thoracic - α , thoracic-lumbar - β , sacral-lumbar γ) and the angle of pelvic anteversion in the sagittal plane and the sitting forward bending test in the boys involved in the study

BOYS					
Sitting forward bending (cm)		Angle			Angle of (pelvic) anteversion
		α	β	γ	
r		0.07	0.16	-0.01	0.05
Correlation power		hardly any	poor	hardly any	hardly any
t		0.71	1.67	0.09	0.54
Statistical significance		lack	lack	lack	lack

Table 6
Correlations occurring between the angles characteristic of anterior-posterior pelvic tilts (upper thoracic - α , thoracic-lumbar - β , sacral-lumbar γ) and the angle of pelvic anteversion in the sagittal plane and the shuttle run 10 x 5 test in the boys involved in the study

BOYS					
Shuttle Run 10 x 5 (s)		Angle			Angle of (pelvic) anteversion
		α	β	γ	
r		0.18	0.1	0.23	-0.12
Correlation power		poor	poor	poor	poor
t		1.85	1.01	2.42	1.21
Statistical significance		lack	lack	$\alpha = 0.05$	lack

Table 7
Correlations occurring between the angles characteristic of anterior-posterior pelvic tilts (upper thoracic - α , thoracic-lumbar - β , sacral-lumbar γ) and the angle of pelvic anteversion in the sagittal plane and the standing broad jump test in the boys involved in the study

BOYS					
Standing Broad Jump (cm)		Angle			Angle of (pelvic) anteversion
		α	β	γ	
r		-0.1	0.04	-0.12	-0,001
Correlation power		poor	hardly any	poor	hardly
t		1.02	0.4	1.19	0,02
Statistical significance		lack	lack	lack	lack

were defined between this test and the upper thoracic angle – α ($r = 0.11$), the thoracic-lumbar angle - β ($r = 0.19$) and the lumbar-sacral γ ($r = 0.13$) (Table 10).

When examining the results of the correlations between the sit-and-reach test and the spinal angles in the sagittal plane (Table 11) there can be defined a statistically significant correlation at the assumed level $\alpha = 0.05$ with the following parameters: the thoracic-lumbar angle – β (average correlation $r = 0.32$) and the

angle of pelvic anteversion (poor correlation $r = -0.22$). The correlation between the above mentioned test and the angle β is consistent (when one variable increases and so does the second one) whereas the second correlation is of an opposite character. A poor but statistically insignificant correlation was defined between the sit-and-reach test and the upper thoracic angle ($r = 0.12$).

When conducting the statistical analysis of the bend forward test findings (Table 12) there were noticed

Table 8

Correlations occurring between the angles characteristic of anterior-posterior pelvic tilts (upper thoracic - α , thoracic-lumbar - β , sacral-lumbar γ) and the angle of pelvic anteversion in the sagittal plane and the handgrip test in the boys involved in the study

		BOYS			Angle of (pelvic) anteversion		
		Angle		α	β	γ	
<i>R</i>		0.12	-0.02	-0.03		0.13	
<i>Correlation power</i>		<i>poor</i>	hardly any	hardly any		<i>poor</i>	
<i>T</i>		1.2	0.25	0.31		1.35	
Statistical significance		lack	lack	lack		lack	

Table 9

Correlations occurring between the angles characteristic of anterior-posterior pelvic tilts (upper thoracic - α , thoracic-lumbar - β , sacral-lumbar γ) and the angle of pelvic anteversion in the sagittal plane and the number of points scored in the Step-test in the boys involved in the study

		BOYS			Angle of (pelvic) anteversion		
		Angle		α	β	γ	
<i>r</i>		-0.18	-0.03	0.03		0.001	
<i>Correlation power</i>		<i>poor</i>	hardly any	hardly any		hardly any	
<i>t</i>		1.87	0.34	0.30		0.04	
Statistical significance		lack	lack	lack		lack	

Table 10

Correlations occurring between angles characteristic of anterior-posterior pelvic tilts (upper thoracic - α , thoracic-lumbar - β , sacral-lumbar γ) and the angle of pelvic anteversion in the sagittal plane and the number of points scored in the bent arm hang test in the girls involved in the study

		GIRLS			Angle of (pelvic) anteversion		
		Angle		α	β	γ	
<i>R</i>		0.11	0.19	0.13		-0.19	
<i>Correlation power</i>		<i>poor</i>	poor	poor		hardly any	
<i>T</i>		1.07	1.95	1.35		0.98	
Statistical significance		lack	lack	lack		lack	

Table 11

Correlations occurring between the angles characteristic of anterior-posterior pelvic tilts (upper thoracic - α , thoracic-lumbar - β , sacral-lumbar γ) and the angle of pelvic anteversion in the sagittal plane and the number of points scored in the sit-and-reach test in the girls involved in the study

Sit-and-Reach (number)		Angle			Angle of (pelvic) anteversion
		A	β	γ	
r	0.12	0.32	-0.08	-0.22	
Correlation power	Poor	moderate	hardly any	poor	
t	1.19	3.33	0.77	2.29	
Statistical significance	Lack	$\alpha = 0.05$	lack	lack	$\alpha = 0.05$

Table 12

Correlations occurring between the angles characteristic of anterior-posterior pelvic tilts (upper thoracic - α , thoracic-lumbar - β , sacral-lumbar γ) and the angle of pelvic anteversion in the sagittal plane and the number of points scored in the sitting forward bend test in the girls involved in the study

Sitting forward bend (cm)		Angle			Angle of (pelvic) anteversion
		A	β	γ	
r	0.04	0.1	-0.05	-0.22	
Correlation power	hardly any	poor	hardly any	poor	
t	0.42	1.02	0.54	2.29	
Statistical significance	Lack	lack	lack	lack	$\alpha = 0.05$

statistically significant correlations at the assumed level $\alpha = 0.05$ with the angle of pelvic anteversion (poor correlation $r = -0.22$). The correlation between this test and the angle of pelvic anteversion is opposite. Moreover, there is a poor, statistically insignificant correlation with the thoracic-lumbar angle - β ($r = 0.1$).

The results of the 10 x 5m shuttle run test reveal a statistically significant correlation at the assumed level $\alpha = 0.05$ with the thoracic-lumbar angle - β (poor correlation $r = -0.22$, of an opposite character, i.e. the higher value of the variable the lower result of the shuttle run test). Besides, statistically insignificant correlations at the low level can be observed with the upper thoracic angle α ($r = -0.14$) (Table 13).

Besides, one low and statistically insignificant correlation (at the assumed level $\alpha = 0.05$) was defined between the following tests: the standing broad jump test and the thoracic-lumbar angle - β ($r = 0.22$) (Table 14) as well as the handgrip test and the upper thoracic angle α ($r = 0.21$) (Table 15). The remaining correlations between the above mentioned tests and the step-test and the evaluated angles were at a low level and they were statistically insignificant (Table 14, 15, 16).

When analyzing the correlations between the independent descriptive variable: body posture (correct

and incorrect) and the EUROFIT physical fitness tests, the step-test and the reaction of the circulatory system in the fifth minute of an exercise and after one-minute rest in the studied boys and girls from selected rural schools of the Kuyavia-Pomerania Province, there was defined only one statistically significant correlation at the assumed level $\alpha = 0.05$, with the bent arm hang test ($r = 0.26$) in girls (Table 20). Other correlations both in boys and girls turned out to be statistically insignificant (Table 17, 18, 19, 20). All the correlations in both studied groups had a consistent character whereas better results were achieved in fitness tests by children with the correct body posture.

Discussion

A number of researchers decide to evaluate fitness of different populations and they seek for its correlations with health [1]. Nevertheless, in the available reference books only few papers concern the formation of the fitness level of children with health disorders. Most frequently scholarly publications, when exploring the issue of faulty postures in the context of fitness, refer to children aged 10 years and older [3, 4, 10], but they refer less often to younger children at the school age. These publications concern children from towns and cities [4, 12, 13, 21, 26] more frequently than from rural areas. Since no reference material was found, the

Table 13

Correlations occurring between the angles characteristic of anterior-posterior pelvic tilts (upper thoracic - α , thoracic-lumbar - β , sacral-lumbar γ) and the angle of pelvic anteversion in the sagittal plane and the number of points scored in the 10 x 5 m shuttle run test in the girls involved in the study

Shuttle Run 10 x 5 (s)		Angle			Angle of (pelvic) anteversion
		α	β	γ	
		r	-0.14	-0.22	-0.01
	Correlation power	poor	poor	hardly any	hardly any
	t	1.41	2.20	0.09	0.33
	Statistical significance	lack	$\alpha = 0.05$	lack	lack

Table 14

Correlations occurring between the angles characteristic of anterior-posterior pelvic tilts (upper thoracic - α , thoracic-lumbar - β , sacral-lumbar γ) and the angle of pelvic anteversion in the sagittal plane and the number of points scored in the standing broad jump test in the girls involved in the study

Standing Broad Jump (cm)		Angle			Angle of (pelvic) anteversion
		α	β	γ	
		R	0.16	0.22	-0.01
	Correlation power	poor	poor	hardly any	poor
	T	1.63	2.30	0.06	1.58
	Statistical significance	lack	$\alpha = 0.05$	lack	lack

Table 15

Correlations occurring between the angles characteristic of anterior-posterior pelvic tilts (upper thoracic - α , thoracic-lumbar - β , sacral-lumbar γ) and the angle of pelvic anteversion in the sagittal plane and the number of points scored in the handgrip test in the girls involved in the study

Handgrip Test (KG)		Angle			Angle of (pelvic) anteversion
		α	β	γ	
		R	0.21	0.15	0.09
	Correlation power	poor	poor	hardly any	hardly any
	T	2.15	1.46	0.9	0.4
	Statistical significance	$\alpha = 0.05$	lack	lack	lack

research study developed by authors analysing other age groups were referred to. The data analyses conducted by Chrzanowska and Chojnacki [4], among others, in the research of students, by Prętiewicz-Abacjew et al. [21] in the research of children beginning school education, by Resiak [26] in the research of children at the beginning of primary school education, by Sudera et al. [28] in the research of children in pre-school age, by Cieszkowski et al. [5] in the research of children aged 7-10 years.

The observed persons with incorrect body postures in the research conducted by Chrzanowska and Chojnacki [4] revealed worse endurance and agility, in the research by Prętiewicz-Abacjew et al. [21] and Resiak [26] – flexibility of the spine, trunk muscle strength, explosive strength of lower limbs and endurance, in the research by Cieszkowski et al. [5] – jumping ability, shoulder muscle strength and shoulder girdle strength and endurance, but they revealed better flexibility. The above presented

Table 16

Correlations occurring between the angles characteristic of anterior-posterior pelvic tilts (upper thoracic - α , thoracic-lumbar - β , sacral-lumbar γ) and the angle of pelvic anteversion in the sagittal plane and the number of points scored in the step test in the boys involved in the study

Step-test (points)		Angle			Angle of (pelvic) anteversion
		α	β	γ	
		<i>r</i>	-0.07	0.06	0.09
<i>Correlation power</i>		hardly any	hardly any	hardly any	<i>poor</i>
<i>t</i>		0.68	0.61	0.93	1.1
Statistical significance		lack	lack	lack	lack

Table 17

Correlations occurring between the posture (correct and incorrect) and selected tests of the EUROFIT physical fitness test in the boys involved in the study

Type of posture (correct and incorrect)		BOYS					
		Bent Arm Hang (s)	Sit-and-Reach (number)	Sitting forward bend (cm)	10 x 5 (s) Shuttle Run	Standing Broad Jump	Handgrip Test (KG)
C	0.06	0.11	0.14	0.08	0.13	0.08	
Correlation power	hardly any	poor	poor	hardly any	poor	hardly any	
χ	0.34	1.26	2.13	0.76	1.8	0.69	
Statistical significance	Lack	lack	lack	lack	lack	lack	

Table 18

Correlations occurring between the posture (correct and incorrect) and the number of points scored in the step-test and the reaction of the circulatory system in the fifth minute of an exercise and after one-minute rest in the boys involved in the study

Type of posture (correct and incorrect)		BOYS			
		Step-test (points)	Reaction of the circulatory system		
			in the fifth minute of an exercise	after one-minute rest	
C	0.12		0.14	0.13	
Correlation power	poor		poor	Poor	
χ	1.54		2.21	1.89	
Statistical significance	lack		lack	Lack	

results of other authors, like in own research, reveal a significant influence of body posture on fitness of children and teenagers.

Conclusions

To summarise the correlations between the characteristics of body posture and motor skills and the reaction of the circulatory system in the sagittal plane in the studied children, the following can be stated:

- ✓ As far as boys are concerned, the factors which most frequently reveal a statistically significant correlation with the studied fitness parameters, at the assumed level $\alpha = 0.05m$, include: the

lumbar-sacral angle (the sit-and-reach and 10 x 5 shuttle run tests),

- ✓ Such independent variables as the upper thoracic angle and the thoracic-lumbar angle correlated with the results of only one of the fitness tests (in both cases this was the bent arm hang test),
- ✓ The remaining parameters of body posture did not reveal statistically significant correlations with any of the conducted tests intended to evaluate fitness of the studied boys,
- ✓ As far as girls are concerned, the factors which most frequently correlate with the studied fitness

Table 19

Correlations occurring between the posture (correct and incorrect) and selected tests of the EUROFIT physical fitness test in the boys involved in the study

Type of posture (correct and incorrect)	GIRLS						
		Bent Arm Hang (s)	Sit-and-Reach (number)	Sitting forward bend (cm)	10 x 5 (s) Shuttle Run	Standing Broad Jump	Handgrip Test (KG)
C	0.26	0.23	0.15	0.13	0.06	0.15	
Correlation power	<i>low</i>	low	low	low	hardly any	low	
χ^2	7.08	5.53	3.45	1.66	0.36	2.22	
Statistical significance	$\alpha = 0.05$	lack	lack	lack	Lack	lack	

Table 20

Correlations occurring between the posture (correct and incorrect) and the number of points scored in the step-test and the reaction of the circulatory system in the fifth minute of an exercise and after one-minute rest in the boys involved in the study

Type of posture (correct and incorrect)	GIRLS			
		Step-test (points)	Reaction of the circulatory system	
			in the fifth minute of an exercise	after one-minute rest
C	0.06	0.13	0.15	
Correlation power	hardly any	poor	poor	
χ^2	0.31	1.64	2.48	
Statistical significance	lack	lack	lack	

parameters are: the thoracic-lumbar angle (tests: sit-and-reach, 10 x 5 shuttle run and standing broad jump) and the angle of pelvic anteversion (tests: sit-and-reach and sitting forward bend),

- ✓ The independent variables like the correct posture and the upper thoracic angle correlated with the results of only one of the fitness tests

(the first one with the handgrip test whereas the second one with the step-test),

- ✓ The remaining posture indexes did not reveal any statistically significant correlations with any of the conducted tests intended to evaluate the fitness level of the children involved in the study.

References

1. Bouchard C., Shephard R.J., (1994), Physical activity, fitness, and health: The model and key concepts. [In:] Physical Activity, Fitness Health. (ed.) Bouchard C., Shephard R. J., Stephens T. Champaing, Human Kinetics, p. 77–88.
2. Boziłow W., Krakowiak H., Gworys B., Jeka S., Ćabrić M., Kurlej W., (2003), Analiza zmienności wybranych wskaźników rozwoju somatycznego dzieci i młodzieży wiejskiej województwa kuj.-pom. [W:] Uwarunkowania rozwoju dzieci i młodzieży wiejskiej. (red.) J. Zagórski, M. Skład. Monografie IMW, Lublin. 200 p.
3. Chojnacki K., (1980), Współzależności wybranych wskaźników budowy z postawą ciała i wydolnością fizyczną dzieci ze szkoły sportowej w Ustrzykach Dolnych. [W:] Roczniki Naukowe AWF nr XVII. PWN, Warszawa-Kraków, s. 85– 89.
4. Chrzanowska M., Chojnacki, K., (1976), Z badań nad stanem postawy ciała i sprawnością fizyczną studentów. [W:] Kultura Fizyczna 2, s. 61-64.
5. Cieszkowski S., Lenik J., Lenik P., Szybisty A., (2005), Rozwój motoryczny dzieci z wadami postawy w wieku 7-10 lat. Analiza porównawcza. [W:]. Korektywa i kompensacja zaburzeń w rozwoju fizycznym dzieci i młodzieży. (red.) K. Górnjak. ZWWF, Biala Podlaska, t. II, s. 24-30.
6. Dega W., (1996), Wady postawy. [W:] Ortopedia i rehabilitacja. (red.) W. Dega, A. Senger. PZWL, Warszawa, t. I. 200 p.
7. Drabik J., (2006), Pedagogiczna kontrola pozytywnych mierników zdrowia fizycznego. AWFiS, Gdańsk. 164 p.
8. EUROFIT, (1989), (H. Grabowski, J. Szopa – przekład z języka

References

1. Bouchard C., Shephard R.J. Physical activity, fitness, and health: The model and key concepts. In: Bouchard C., Shephard R. J., Stephens T. ed. *Physical Activity, Fitness Health*. Champaing, Human Kinetics, 1994, p. 77–88.
2. Boziłow W., Krakowiak H., Gworys B., Jeka S., Ćabrić M., Kurlej W. Analysis of the variability of selected indicators of somatic development of children and adolescents rural province Kujawsko-Pomorskie [Analiza zmienności wybranych wskaźników rozwoju somatycznego dzieci i młodzieży wiejskiej województwa Kujawsko-Pomorskiej]. In: J. Zagórski, M. Skład, ed. *Determinants of the development of children and youth from rural areas Kuyavian-Pomeranian Voivodeship* [Uwarunkowania rozwoju dzieci i młodzieży wiejskiej], Lublin, 2003, 200 p.
3. Chojnacki K. Interdependence of selected indicators of the construction of posture and physical performance of children from school sports in the Lower Ustrzyki [Współzależności wybranych wskaźników budowy z postawą ciała i wydolnością fizyczną dzieci ze szkoły sportowej w Ustrzykach Dolnych]. *Scientific yearbook [Roczniki Naukowe]*, 1980, vol. 17, pp. 85– 89.
4. Chrzanowska M., Chojnacki K. Research on the state of the body posture and physical fitness of students [Z badań nad stanem postawy ciała i sprawnością fizyczną studentów]. *Physical Culture [Kultura Fizyczna]*, 1976, vol.2, pp. 61-64.
5. Cieszkowski S., Lenik J., Lenik P., Szybisty A. Motor development of children with posture at the age of 7-10 years. The comparative analysis [Rozwój motoryczny dzieci z wadami postawy w wieku 7-10 lat. Analiza porównawcza]. *Correction and compensation of disturbances in the physical development of children and youth* [Korektywa i kompensacja zaburzeń w rozwoju fizycznym dzieci i

- angielskiego). AWF, Kraków. 204 p.
9. Górniaak K., (2004), Społeczne uwarunkowania skolioz niskostopniowych u dzieci wiejskich (wybrane aspekty). [W:] Uwarunkowania rozwoju dzieci i młodzieży wiejskiej. (red.) J. Zagórski, H. Popławska, M. Skład. Monografie, Instytut Medycyny Wsi, Lublin. s. 394–400.
 10. Górniaak K., Lichota M., (2006), Postawa i budowa ciała dzieci wiejskich ze skoliozą niskostopniową. [W:] Uwarunkowania rozwoju dzieci i młodzieży wiejskie. (red.) J. Saczuk, ZWWF, Biała Podlaska, t. I, s. 274-281.
 11. Kasperekzyk T., (1988) Postawa ciała a wybrane cechy morfologiczne i funkcjonalne u dzieci w wieku 8-15 lat. Wydanie Monograficzne, 32. AWF, Kraków. 240 p.
 12. Lewandowski A., Talar J., (2005), Wady postawy a sprawność fizyczna dzieci w wieku szkolnym na przykładzie badań młodzieży Gimnazjum nr 23 w Bydgoszczy. [W:] Medycyna Sportowa, Vol. 21, nr 2, pp.99-110.
 13. Lichota M., (2003), Pierwszy krytyczny okres posturogenety w badaniach postawy ciała wybranej grupy dzieci z Białej Podlaskiej. (rozprawa doktorska) AWF, Gdańsk. 200 p.
 14. Łobocki M., (2004), Wprowadzenie do metodologii badań pedagogicznych. Wydawnictwo „Impuls”.
 15. Mazur B., Serapata M., Scibich A., Pyda A., (1975) Ocena wydolności fizycznej dzieci szkolnych miasta Katowice na podstawie harwardzkiej próby stopnia. Pediatria Polska nr 7, s.887-892.
 16. Mleczko E., (2006), Rozwój biologiczny dzieci i młodzieży ze środowiska wiejskiego w polskich badaniach auksologicznych. [W:] Uwarunkowania rozwoju dzieci i młodzieży wiejskiej. (red.) J. Saczuk. AWF, Warszawa, ZWWF, Biała Podlaska, t. I, s.39-79.
 17. Napierała M., (2000), Dziecko z regionu kujawsko – pomorskiego: rozwój fizyczny i motoryczny dzieci z klas początkowych. AB, Bydgoszcz. 168 p.
 18. Prętkiewicz-Abacjew E., (1993), Wybrane cechy postawy ciała w płaszczyźnie strzałkowej u chłopców i dziewcząt w okresie dojrzewania płciowego. [W:] Dysfunkcje kręgosłupa – diagnostyka i terapia. (red.) J. Nowotny. AWF, Katowice. 212 p.
 19. Prętkiewicz-Abacjew E., (2002), Kinematyka chodu a postawa ciała dzieci sześciolatnic. AWF, Gdańsk. 240 p.
 20. Prętkiewicz-Abacjew E., (2006), A comparison of body positioning in children with and without genu valgum. *Gait and Posture*. vol. 24, suppl.2, pp.184-186.
 21. Prętkiewicz-Abacjew E., Resiak M., Wróblewska A., (1996), Postawa i sprawność fizyczna dzieci rozpoczynających naukę w szkole. [W:] Nauki o kulturze fizycznej w służbie współczesnego człowieka. (red.) A. Pawłucki. AWF, Gdańsk, s.165-170.
 22. Prętkiewicz-Abacjew E., Zajt-Kwiatkowska J., Rogo B., (2007), Zmiany w postawie ciała chłopców i dziewcząt w latach 1995 – 2004. [W:] Sześciolatne Dziecko Gdańskie w Latach 1995 - 2004. Szkic antropomotoryczny. (red.) J. Drabik, M. Resiak, E., Prętkiewicz-Abacjew. AWFis, Gdańsk, s. 39-50.
 23. Przeweda R., (1981), Rozwój somatyczny i motoryczny. Wydawnictwa Szkolne i Pedagogiczne, Warszawa. 204 p.
 24. Przeweda R., Dobosz J., (2003), Kondycja fizyczna polskiej młodzieży. AWF, Warszawa.
 25. Raczek J., (1995), Hipokinezja i jej skutki jako problem współczesnej cywilizacji. [W:] Nauki o kulturze fizycznej wobec wyzwań współczesnej cywilizacji. Abstrakty na konferencję z okazji 25-lecia AWF w Katowicach. AWF, Katowice. 168 p.
 26. Resiak M., (1996), Wybrane aspekty rozwoju somatycznego, motorycznego i psychicznego dzieci gdańskich u progu szkoły podstawowej. Rozprawa doktorska. AWF, Gdańsk. 200 p.
 27. Siniarska A., Koziol R., Szemik M., (1989) Biological state of rural populations from different geographic environments in Poland. *Studies in Human Ecology*, 8, pp.175-200.
 28. Suder A., Sobiecki J., Kościuk T., Pałosz J., (2002), Sprawność motoryczna a postawa ciała dzieci w wieku przedszkolnym. [W:] Nowiny Lekarskie. Nr 4-5 (71). Poznań, s.230-235.
 29. Śliwa W., Chlewicka E., Chromik K., (2005), Różnice środowiskowe w kształcie krzywizn kręgosłupa chłopców w wieku 5-15 lat. [W:] młodzieży], 2005, vol.3, pp. 24-30.
 6. Dega W., Senger A. *Posture defects* [Wady postawy], Warsaw, PZWL, 1996, T.1, 200 p.
 7. Drabik J. *Pedagogical positive control measures of physical health* [Pedagogiczna kontrola pozytywnych mierników zdrowia fizycznego], Gdańsk, AWFis, 2006, 164 p.
 8. Grabowski H., Szopa J. *EUROFIT*. Krakow, AWF, 1989, 204 p.
 9. Górniaak K. Social determinants of progressive scoliosis in children selected aspects [Społeczne uwarunkowania skolioz niskostopniowych u dzieci wiejskich wybrane aspekty]. In: J. Zagórski, H. Popławska, M. Skład, ed. *Determinants of the development of children and youth from rural areas* [Uwarunkowania rozwoju dzieci i młodzieży wiejskiej]. Institute of Agricultural Medicine, Lublin, 2004, pp. 394–400.
 10. Górniaak K., Lichota M. Posture and body of children with progressive scoliosis [Postawa i budowa ciała dzieci wiejskich ze skoliozą niskostopniową]. In: J. Saczuk, ed. *Determinants of the development of children and youth from rural areas* [Uwarunkowania rozwoju dzieci i młodzieży wiejskiej]. White Podlasie, Faculty of Physical Education, 2006, T.1, pp. 274-281.
 11. Kasperekzyk T. Posture and selected morphological and functional in children aged 8-15 years [Postawa ciała a wybrane cechy morfologiczne i funkcjonalne u dzieci w wieku 8-15 lat], Krakow, Academy of Physical Education, 1988, T.32, 240 p.
 12. Lewandowski A., Talar J. Posture defects and physical fitness of school children as an example to the youth Secondary School No. 23 in Bydgoszcz [Wady postawy a sprawność fizyczna dzieci w wieku szkolnym na przykładzie badań młodzieży Gimnazjum nr 23 w Bydgoszczy], *Sports Medicine* [Medycyna Sportowa], 2005, vol.21(2), pp. 99-110.
 13. Lichota M. *The first critical period posturogenesis research posture selected group of children from the White Podlasie* [Pierwszy krytyczny okres posturogenety w badaniach postawy ciała wybranej grupy dzieci z Białej Podlaskiej]. Dokt. Diss., Gdańsk, 2003, 200 p.
 14. Łobocki M. *Introduction to the methodology of educational research* [Wprowadzenie do metodologii badań pedagogicznych], Impuls Publ., 2004, 160 p.
 15. Mazur B., Serapata M., Scibich A., Pyda A. Evaluation of physical fitness of school children of the city of Katowice on the basis of trial Harvard degree [Ocena wydolności fizycznej dzieci szkolnych miasta Katowic na podstawie harwardzkiej próby stopnia], *Pediatrics Poland* [Pediatria Polska], 1975, vol.7, pp. 887-892.
 16. Mleczko E. Biological development of children and young people from the rural environment in Polish studies auxological [Rozwój biologiczny dzieci i młodzieży ze środowiska wiejskiego w polskich badaniach auksologicznych]. In: J. Saczuk, ed. *Determinants of the development of children and youth from rural areas* [Uwarunkowania rozwoju dzieci i młodzieży wiejskiej]. Academy of Physical Education, Warsaw, Faculty of Physical Education, White Podlasie, 2006, T.1, pp. 39-79.
 17. Napierała M. *Child of the Kujawsko - Pomeranian: physical and motor development of children with classes starting* [Dziecko z regionu kujawsko – pomorskiego: rozwój fizyczny i motoryczny dzieci z klas początkowych], Bydgoszcz, AB, 2000, 168 p.
 18. Prętkiewicz-Abacjew E. Selected characteristics of body posture in the sagittal plane in boys and girls during puberty [Wybrane cechy postawy ciała w płaszczyźnie strzałkowej u chłopców i dziewcząt w okresie dojrzewania płciowego]. In: J. Nowotny, ed. *Dysfunctions of the spine - diagnosis and therapy* [Dysfunkcje kręgosłupa – diagnostyka i terapia], Academy of Physical Education Katowice, 1993, 212 p.
 19. Prętkiewicz-Abacjew E. *Kinematics of gait and posture of children six years* [Kinematyka chodu a postawa ciała dzieci sześciolatnic], AWF, Gdańsk, 2002, 240 p.
 20. Prętkiewicz-Abacjew E. A comparison of body positioning in children with and without genu valgum. *Gait and Posture*. 2006, vol.24(2), pp. 184-186.
 21. Prętkiewicz-Abacjew E., Resiak M., Wróblewska A. Attitude and physical fitness of children at school [Postawa i sprawność fizyczna dzieci rozpoczynających naukę w szkole]. In: A. Pawłucki, ed. *The science about physical culture in the service of modern man* [Nauki o kulturze fizycznej w służbie współczesnego człowieka], AWF, Gdańsk, 1996, pp. 165-170.
 22. Prętkiewicz-Abacjew E., Zajt-Kwiatkowska J., Rogo B. Changes in body posture boys and girls in the years 1995 - 2004 [Zmiany w postawie ciała chłopców i dziewcząt w latach 1995 – 2004]. In: J. Drabik, M. Resiak, E., Prętkiewicz-Abacjew, ed. *A child of six at Gdańsk in Years 1995-2004. Draft antropomotoryczny* [Sześciolatnie Dziecko Gdańskie w Latach 1995 - 2004. Szkic antropomotoryczny],

- Korektywa i kompensacja zaburzeń w rozwoju fizycznym dzieci i młodzieży. (red.) K. Górnjak. AWF, Warszawa, ZWWF, Biala Podlaska, t. I, s.256-262.
30. Wolański N., (2005), Rozwój biologiczny człowieka. Podstawy auksologii, gerontologii i promocji zdrowia. PWN, Warszawa. 216 p.
31. Zeyland-Malawka E., (1997) Badanie postawy ciała. AWF Gdańsk. 280 p.
32. Przewęda R. The development of somatic and motor [Rozwój somatyczny i motoryczny], Publishing school and pedagogical, Warsaw, 1981, 204 p.
33. Przewęda R., Dobosz J. Physical condition of Polish youth [Kondycja fizyczna polskiej młodzieży], Academy of Physical Education, Warsaw, 2003, 216 p.
34. Raczek J. Hypokinesis and its effects as a problem of modern civilization [Hipokinezja i jej skutki jako problem współczesnej cywilizacji]. In: The science of physical culture to the challenges of modern civilization [Nauki o kulturze fizycznej wobec wyzwań współczesnej cywilizacji]. Academy of Physical Education, Katowice, 1995, 168 p.
35. Resiak M. Selected aspects of the development of physical, mental and motor skills of children from Gdańsk on the threshold of primary school [Wybrane aspekty rozwoju somatycznego, motorycznego i psychicznego dzieci gdańskich u progu szkoły podstawowej], Dokt. Diss., AWF, Gdańsk, 1996, 200 p.
36. Siniarska A., Koziol R., Szemik M. Biological state of rural populations from different geographic environments in Poland. *Studies in Human Ecology*, 1989, vol.8, pp. 175-200.
37. Suder A., Sobiecki J., Kościuk T., Pałosz J. Motor efficiency and posture preschool children [Sprawność motoryczna a postawa ciała dzieci w wieku przedszkolnym]. *Medical News [Nowiny Lekarskie]*, 2002, vol.4-5(71), pp. 230-235.
38. Śliwa W., Chlebicka E., Chromik K. Environmental differences in the shape of the spinal curves in boys aged 5-15 years [Różnice środowiskowe w kształcie krzywizn kręgosłupa chłopców w wieku 5-15 lat]. In: K. Górnjak ed. *Correction and compensation of disturbances in the physical development of children and youth* [Korektywa i kompensacja zaburzeń w rozwoju fizycznym dzieci i młodzieży], Academy of Physical Education, Warsaw, Faculty of Physical Education, White Podlasie, 2005, T.1, pp. 256-262.
39. Wolański N. *Biological development of man. Basics auxology, gerontology and health promotion* [Rozwój biologiczny człowieka. Podstawy auksologii, gerontologii i promocji zdrowia], PWN, Warsaw, 2005, 216 p.
40. Zeyland-Malawka E. *Research of body posture* [Badanie postawy ciała], AWF, Gdańsk, 1997, 280 p.

Информация об авторах:

Жуковска Анна: ORCID: 0000-0001-6688-9314; zukowska@ukw.edu.pl; Университет Казимира Великого в Быдгощ; ул. Ходкевича 30, г.Быдгощ 85-064, Польша.;

Шарк-Ецкардт Мирослава: ORCID: 0000-0003-0668-0951; rektor@ukw.edu.pl; Университет Казимира Великого в Быдгощ; ул. Ходкевича 30, г.Быдгощ 85-064, Польша.

Мушкита Радослав Каролиевич: д.п.н., проф.; ORCID: 0000-0001-6057-1583; radek@muszkieta.com; Университет Казимира Великого в Быдгощ; ул. Ходкевича 30, г.Быдгощ 85-064, Польша.

Ермакова Татьяна Сергеевна: ORCID: 0000-0002-3081-0229; tatiana-iermakova@rambler.ru; Харьковский национальный педагогический университет имени Г.С. Сквороды; ул. Артема, 29, г. Харьков, 61002, Украина.

Цитируйте эту статью как: Жуковска Анна, Шарк-Ецкардт Мирослава, Мушкита Радослав, Ермакова Татьяна. Характеристики положения тела в сагиттальной плоскости и физическое состояние первоклассников из сельской местности // Педагогика, психологія та медико-біологічні проблеми фізичного виховання і спорту. – 2014. – № 7. – С. 50-60. doi:10.6084/m9.figshare.1015583

Электронная версия этой статьи является полной и может быть найдена на сайте: <http://www.sportpedagogy.org.ua/html/arhive.html>

Эта статья Открытого Доступа распространяется под терминами Creative Commons Attribution License, которая разрешает неограниченное использование, распространение и копирование любыми средствами, обеспечивающими должное цитирование этой оригинальной статьи (<http://creativecommons.org/licenses/by/3.0/deed.ru>).

Дата поступления в редакцию: 14.03.2014 г.
Опубликовано: 28.03.2014 г.

Information about the authors:

Żukowska Hanna: ORCID: 0000-0001-6688-9314; zukowska@ukw.edu.pl; Kazimierz Wielki University in Bydgoszcz; Chodkiewicza str. 30, 85-064 Bydgoszcz, Poland.

Szark-Eckardt Mirosława: ORCID: 0000-0003-0668-0951; rektor@ukw.edu.pl; Kazimierz Wielki University in Bydgoszcz; Chodkiewicza str. 30, 85-064 Bydgoszcz, Poland.

Muszkieta Radosław: ORCID: 0000-0001-6057-1583; radek@muszkieta.com; Kazimierz Wielki University in Bydgoszcz; Chodkiewicza str. 30, 85-064 Bydgoszcz, Poland.

Iermakova T.S.: ORCID: 0000-0002-3081-0229; tatiana-iermakova@rambler.ru ; G.S. Skovoroda Kharkiv National Pedagogical University ; Artema str., 29, Kharkiv, 61002, Ukraine.

Cite this article as: Żukowska Hanna, Szark-Eckardt Mirosława, Muszkieta Radosław, Iermakova Tetiana. Characteristics of body posture in the sagittal plane and fitness of first-form pupils from rural areas. *Pedagogics, psychology, medical-biological problems of physical training and sports*, 2014, vol.7, pp. 50-60. doi:10.6084/m9.figshare.1015583

The electronic version of this article is the complete one and can be found online at: <http://www.sportpedagogy.org.ua/html/arhive-e.html>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited (<http://creativecommons.org/licenses/by/3.0/deed.en>).

Received: 14.03.2014

Published: 28.03.2014