

The effects of stretching exercise on hemodynamic responses and post-exercise hypotension in normotensive women students

Hamid Mohebbi, Abdulbaset Maroofi, Nazanin Anasri, Aboozar Jorbonian

University of Guilan, Rasht, Iran

Annotation:

Aim: The aim of the present study was to evaluate acute effects of SE on post-exercise hemodynamic responses for 1-h in normotensive sedentary young women. **Methods:** Sixteen women (21.56 ± 1.21 yr; 159.6 ± 0.5 cm; 54.53 ± 6.02 kg) were randomly assigned to SE ($n = 8$) and control (C) groups ($n = 8$). SE group performed 20 stretches for the whole body. Each SE was repeated 2 times. Rest interval between repetitions and movement 10 s were considered. Systolic blood pressure (SBP), diastolic BP (DBP), mean arterial BP (MAP), rate pressure product (RPP), pulse pressure (PP) and heart rate (HR) were measured during 1-h (minutes: 0, 15, 30, 45 and 60) in SE and C groups. **Results:** There were significant decreases ($P < 0.05$) in SBP, DBP, and MAP, after SE from 15 to 30 min of recovery than baseline. RPP also decreased significantly ($P < 0.05$) after SE from 15 to 45 min when compared with baseline. But SE did not cause any significant changes in PP when compared with C group. Hemodynamic responses were not altered during the control trial. **Conclusions:** The results of this study showed that an acute SE lead to PEH. Therefore, SE may be an interesting training strategy to acutely decrease BP in young women.

Хамід Мохеббі, Абдулбасет Мароофі, Наханін Анасрі, Абоозар Джорбоніан. Вплив вправ на розтягування на гемодинамічний статус і профілактику гіпотонії у студентів дівчат, що мають нормальний артеріальний тиск. **Мета:** мета даного дослідження - оцінити вплив вправ на розтягування на профілактику гіпотонії при одноденних заняттях у дівчат, що ведуть малорухливий спосіб життя і що мають нормальний артеріальний тиск. **Матеріал:** 16 жінок (вік - 21.56 ± 1.21 років; зрост - 159.6 ± 0.5 см; вага - 54.53 ± 6.02 кг) було розподілено на експериментальну (ЕГ) і контрольну (КГ) групу ($n = 8$). Експериментальна група мала 2 серії повторень. Експериментальна група виконувала 20 розтягань для всього тіла. Кожен інтервал між повтореннями і вправами складав 10 секунд. Тиск (SBP) систоли, тиск (DBP) діастоли, середній артеріальний тиск BP (MAP), величина нормального тиску (RPP), пульс (PP) і частота сердечних скорочень (HR) були визначені протягом 1 години (хвилини: 0, 15, 30, 45 і 60) в експериментальній і контрольній групі. **Результати:** у експериментальній групі встановлені істотні зменшення ($P < 0.05$) в SBP, DBP і MAP з 15 до 30 хвилин відновлення порівняно з початковими величинами. RPP також значно зменшилася ($P < 0.05$) з 15 до 45 хвилин порівняно з початковими величинами. Але в експериментальній групі не виявлено ніяких істотних змін в РР порівняно з контрольною. Гемодинамічні реакції не були змінені протягом контрольного випробування. **Висновки:** результати цього дослідження показали, що запропоновані заняття для експериментальної групи направлені на профілактику гіпотонії у студентів дівчат, що мають нормальний артеріальний тиск. Тому, програма занять експериментальної групи, можливо, є цікавою тренувальною стратегією для різкого зменшення середнього артеріального тиску у молодих дівчат.

Хамід Мохеббі, Абдулбасет Мароофі, Наханін Анасрі, Абоозар Джорбоніан. Влияние упражнений на растяжение на гемодинамический статус и профилактику гипотонии у студентов девушек, имеющих нормальное артериальное давление. **Цель:** цель настоящего исследования - оценить влияние упражнений на растяжение на профилактику гипотонии при одночасовых занятиях у девушек, ведущих малоподвижный образ жизни и имеющих нормальное артериальное давление. **Материал:** 16 женщин (возраст - 21.56 ± 1.21 лет; рост - 159.6 ± 0.5 см; вес - 54.53 ± 6.02 кг) были распределены на экспериментальную (ЭГ) и контрольную (КГ) группу ($n = 8$). Экспериментальная группа имела 2 серии повторений. Экспериментальная группа выполняла 20 растягиваний для всего тела. Каждый интервал между повторениями и упражнениями составлял 10 секунд. Систолическое давление (SBP), диастолическое давление (DBP), среднее артериальное давление BP (MAP), величина нормального давления (RPP), пульс (PP) и частота сердечных сокращений (HR) были определены в течение 1 часа (минуты: 0, 15, 30, 45 и 60) в экспериментальной и контрольной группе. **Результаты:** в экспериментальной группе установлены существенные уменьшения ($P < 0.05$) в SBP, DBP и MAP с 15 до 30 минут восстановления в сравнении с исходными величинами. RPP также значительно уменьшился ($P < 0.05$) с 15 до 45 минут в сравнении с исходными величинами. Но в экспериментальной группе не обнаружено никаких существенных изменений в РР в сравнении с контрольной. Гемодинамические реакции не были изменены в течение контрольного испытания. **Выводы:** результаты этого исследования показали, что предложенные занятия для экспериментальной группы направлены на профилактику гипотонии у студентов девушек, имеющих нормальное артериальное давление. Поэтому, программа занятий экспериментальной группы, возможно, является интересной тренировочной стратегией для резкого уменьшения среднего артериального давления у молодых девушек.

Keywords:

stretching, blood pressure, hypotension, student, girl.

розтягання, тиск крові, гіпотонія, студент, дівчина.

растягивание, давление крови, гипотония, студент, девушка.

Introduction

High blood pressure or hypertension (HT) is a risk factor for developing cardiovascular disease (CVD) and according to the World Health Organization (WHO) reports, each year more than 7 billion people in the world will die due to this disease (Melo et al., 2006). According to the American College of Sports Medicine (ACSM), even a modest 3 mmHg drop in systolic blood pressure (SBP) and diastolic blood pressure (DBP) at rest is associated with reduced cardiac mortality by 5 to 9 percent (Pescatello et al., 2004). Although pharmacological interventions are effective for reducing the blood pressure (BP), but the side effects of antihypertensive drugs have been proposed (Chobanian et al., 2003). Therefore, a lifestyle change for instant exercise is very important for

the management of BP.

It has been shown that active individual have a lower risk of becoming hypertensive than those that are sedentary (Cardoso et al., 2010). Evidence suggests implementing regular exercise leads to decrease BP and the risk of CVD (Corrick et al., 2013). BP may decrease below pre-exercise baseline levels after different muscular work, this response is known as post-exercise hypotension (PEH) (Halliwill et al., 2013). This phenomenon could be clinically significance, because it tend to maintain BP of hypertensive subjects transiently at lower levels during different times of the day, especially when BP is at its highest level (Moraes et al., 2011). PEH has been investigated with various exercise including aerobic exercise (Mohebbi et al., 2010), resistance exercise (Rezk et al., 2006), water exercise (Terblanche et al., 2012), and concurrent exercise (Keese et al., 2012). The mechanisms

that cause PEH are complex and multifactorial. It may be mediated by decrease in sympathetic nervous system, cardiac output, and vascular resistance or changes in the release of vasodilator substances (MacDonald., 2002). The decrease in total peripheral resistance induced by exercise usually results from increased bioavailability of nitric oxide (NO, a potent vasodilator) produced by endothelial cells in response to autonomic, hemodynamic and humoral stimuli (Santana et al., 2013).

Special populations such as the elderly and obese may have physical or musculoskeletal limitations and some individual due to inaccessibility gymnasium and high costs, may not have the ability to participation in conventional exercise modalities. Stretching exercise (SE) is an important component of fitness training (Farinatti et al., 2011) and it used in exercise programs for muscle injury prevention (Cipriani et al., 2012). Mechanical stress induced by SE can affect hemodynamic responses (Farinatti et al., 2011). Although it is clear that SE enhance flexibility (Cipriani et al., 2012), but its effect on hemodynamic responses and PEH is unclear. Recently, vong et al. observed that eight weeks of SE resulted to decrease in BP in obese postmenopausal women (vong et al., 2013). A previous study has reported that stretch stimulus to rat's pulmonary vascular endothelial cells leads to an increase release of NO (Kuebler et al., 2003). Hotta et al. also demonstrated that acute SE improves peripheral circulated by production of NO (Hotta et al., 2013). Given the role of SE in the production of NO (Hotta et al., 2013), this model of physical activity may lead to PEH. It has been shown that stretched muscle fibers activate mechanoreceptors, which elicit cardiovascular regulation through parasympathetic inhibition and sympathetic activation (Drew et al., 2008). The fall in heart rate (HR) immediately post-exercise is a sign of parasympathetic nervous activity (Cole et al., 1999). Because of the increased vagal activity is associated with a reduced risk of mortality (Cole et al., 1999), we assessed changes in HR during post-exercise recovery. Rate Pressure Product (RPP, index of myocardial oxygen consumption) or Robinson index is the product of HR and SBP (Nagpal et al., 2007). Decrease in RPP post-exercise, indicating improvement in cardiac function (Piotrowicz et al., 2009). Mean arterial pressure (MAP) is a product of cardiac output and systematic vascular resistance (MacDonald., 2002). Given that PEH may be result from reductions in cardiac output, total peripheral resistance, or both (Santana et al., 2013). Therefore, the calculation of this parameter can help us when discussing the results. Pulse pressure (PP) defined as the difference between SBP and DBP, is an indicator of arterial compliance and used in assessing the risk of heart disease (Franklin et al., 1999). Among the indicators of cardiovascular risk assessment, most emphasis is on the PP (Franklin et al., 1999). Thus, evaluate the changes PP post-exercise may be useful. Therefore, because information on the effect of SE on the PEH seems to be lacking, the purpose of these study was to examine the effect of SE on PEH and post-exercise hemodynamic responses in normotensive

sedentary young women.

Material and Methods

Subjects

Sixteen women volunteered for this study. All subjects completed the short version of the International Physical Activity Questionnaire (IPAQ) and were considered sedentary. Subjects were randomly assigned to SE (n = 8) and control groups (C, n = 8). The inclusion criteria were: normotensive, weight normal, no regular exercise training in the previous 12 months. The exclusion criteria were: use of drugs that could affect cardiovascular function, (i.e. beta-blockers and inhibitors of angiotensin-converting enzyme), smoking, have any cardiovascular and osteomyoarticular problems that could affect performance during SE, being in menstrual bleeding cycle, and SBP and DBP greater than 139 and 89 mmHg respectively. Volunteers completed a thorough physical examination, including a medical history, BP assessment, anthropometric, and orthopedic evaluation prior to participation in the experimental sessions. Moreover, all volunteers signed a written consent and were informed about the risks and benefits of the present study which was approved by the ethical Committee of the Department of Sport Sciences, University of Guilan. Demographic characteristics of subjects are shown in Table 1.

Anthropometric, body fat percentage, and hemodynamic parameters

Height was measured using a wall-stadiometer, by precision of 1 mm. weight and body composition was determined by In Body System (0.3, Korea). SBP and DBP using a standard mercury sphygmomanometer (ALPK, Japan) and HR were measured by an electronic HR monitor (Polar PM80, Finland). The measuring procedure was in accordance with the recommendation of the American Heart Association (Fletcher et al., 1995) and was conducted by an experienced researcher. RPP, MAP and PP were estimated by conventional equations.

Acute exercise session

SE group completed familiarization program prior to testing. During the familiarization subjects were advised regarding proper performance of stretches. Experimental session was conducted in a temperature-controlled room ($23 \pm 1^\circ\text{C}$) between 9:00-11:00 am. Subjects refrained for at least 48-h from intake of caffeine or alcoholic beverage and severe activity that demanded high energy before testing. Before the exercise protocol, in order to measure baseline BP and HR, Subjects seated for 10 minutes and then 5 recordings were taken at 2 minute-intervals. The lowest and highest BP recorded values were removed and the remaining three were averaged to determine of baseline BP. At the end of 10 minutes, HR was measured only once. Immediately after the SE, BP and HR were measured at 0, 15, 30, 45, and 60 minutes with subjects seated. Same parameters were measured for C group without performing any type of exercise. Procedure of SE: researcher closely supervised the SE, which were conducted on a floor mat and focused on whole body major muscles: pectorals major and minor, latissimus dorsi, bicep brachial, triceps, deltoid, trapezius,

illioptas, gluteus, quadriceps, hamstring, leg adductors and gastrocnemius. The 20 stretches were performed in the standing, sitting, and lying position. Each SE was repeated 2 times. The stretched muscle was held for 30 s at point of maximal exertion (defined as RPE > 18). Rest interval between repetitions and movement 10 s were considered. Subjects counted from 1 to 30 during the 30 s stretch to avoid holding their breath during the stretching. Total time SE lasted approximately 38 minute.

Statistical Analysis

All data are presented as mean \pm standard deviation (SD). Independent t-tests was used as appropriate and repeated measures analysis of variance was used to determine the degree of difference SBP, DBP, MAP, HR, RPP and PP between baseline values and at 0, 15, 30, 45, and 60 minutes in C and SE groups. Statistical significance was accepted at the $P < 0.05$.

Results

In C group, SBP, DBP, HR, MAP and PP, did not changed significantly. Figure 1 presents mean values of SBP, DBP and MAP during 1-h after the experimental session. The SBP, DBP and MAP was significantly decreased ($P < 0.05$) at time points of 15 to 30 after SE when compared with the baseline values. Table 1 shows the changes in HR, RPP, and PP from baseline values to 1-h after experimental sessions. HR was significantly

increased immediately after SE when compared with baseline value. However, HR at time points of 15 to 30 was significantly reduced. RPP decreased significantly at time points of 15 to 45 when compared with baseline value. But, SE did not any significant changes in PP.

Discussion

The results indicate that SE induced a drop in BP and RPP when compared with the control session. The absence of any decrease in BP during the non-exercise control trial shows that, decreasing the BP levels after SE are due to the exercise effect and not to the normal diurnal BP variations. It was research has demonstrated that acute exercise (aerobic, resistance, water, and concurrent) induce PEH (Mohebbi et al. 2010; Rezk et al. 2006; Terblanche et al. 2012; Keese et al. 2012). But the effects of SE have not been investigated. The novelty of this study was the acute of SE lead to PEH. To the best of knowledge, this is the first report of the effects of stretching exercise on PEH. The mechanisms underlying the PEH are not fully understood. Changes in the neural control of the circulatory system and the release of vasodilator agents are may be related with the PEH (Lockwood et al., 2005). Lansman et al. showed that a single stretch stimulus to vascular endothelial cells instantly enhanced the opening frequency of Ca^{2+} channels in the cell membrane (Lansman et al., 1987). SE

Table 1

Demographic characteristic of subjects

Parameter	Exercise	Control
Age, yr	21.87 \pm 1.25	21.25 \pm 1.64
Height, cm	159.1 \pm 0.06	158.2 \pm 0.05
Weight, kg	55.86 \pm 4.40	53.20 \pm 7.37
BMI, kg/m ²	22.10 \pm 1.94	21.17 \pm 2.06
Fat, %	28.19 \pm 6.31	32.09 \pm 5.84

Values are shown as mean \pm SD or numbers (%). BMI = body mass index

Table 2

HR, RPP and PP variability parameters in the C and SE groups

	Trials	HR	PP	RPP
Exercise	Pre-exercise	80 \pm 5.24	50.75 \pm 4.77	10105 \pm 536.4
	Post-exercise	86 \pm 6.41 * #	49.75 \pm 6.45	10490 \pm 368.6 #
	15 min	77 \pm 5.13 ¥	48.87 \pm 7.30	9222.5 \pm 484.2 *¥#
	30 min	75 \pm 5.13 ¥	50.25 \pm 4.86	9017.5 \pm 811.7 *¥#
	45 min	73.5 \pm 5.21 ¥#	49.12 \pm 5.44	8831 \pm 590.1 *¥#
	60 min	76.5 \pm 5.42 ¥€	51.12 \pm 4.32	9528 \pm 729.5 €
Control	Pre-exercise	84 \pm 8	47.25 \pm 8.41	10024 \pm 1027.5
	Post-exercise	84.2 \pm 7.9	47.23 \pm 8.44	10490 \pm 368.6
	15 min	83 \pm 5.13	47.25 \pm 7.62	9222 \pm 484.2
	30 min	82 \pm 6.05	46.25 \pm 9.76	9017 \pm 811.6
	45 min	82.5 \pm 6.02	46 \pm 8.34	9802 \pm 736.3
	60 min	81.5 \pm 6.02	46 \pm 8.41	9528 \pm 729.5

Values are shown as mean \pm SD or numbers (%). HR = heart rate, PP = pulse pressure, RPP = rate pressure product

* Significant difference from pre-exercise ($P < 0.05$)

€ Significant difference from 45 min ($P < 0.05$)

¥ Significant difference from post-exercise ($P < 0.05$)

Significant difference between the two group ($P < 0.05$)

could increase Ca^{2+} inflow into vascular endothelial cells by stretch activated Ca^{2+} channels (Singer et al., 1982). This increase leads to augmented calcium-dependent NO production from vascular endothelium (Singer et al., 1982). In the present study, we have observed PEH in SBP, DBP and MAP. Therefore, the PEH induced by SE may be due to NO production from vascular endothelial cells. It has been shown that, when muscle

stretching and contraction occur simultaneously (which is property of the static stretching method, due to the muscle spindle reflex), type III fibers and metaboreceptor activated, and may induce parasympathetic inhibition and baroreflex stimulation and contribute to an increase in the hemodynamic responses (Farinatti et al., 2013). It is accepted that during SE there is a parasympathetic withdrawal and sympathetic stimulation, resulting in




Fig. 1. SBP, DBP, MBP changes following exercise
 * Significant difference from pre-exercise (P<0.05)
 \$ Significant difference from 15 min (P<0.05)
 # Significant difference between the two group (P<0.05)

increases of the HR (Goldberger et al., 2006). These effects are reversed during the post-exercise recovery (Goldberger et al., 2006). Our study also revealed that HR was significantly increased immediately after SE and then gradually returns to the baseline value. Although SE did not lead to significantly decrease in HR below baseline level during the recovery period, but RPP for 30 minutes was lower compared with baseline value. Also, the RPP is the product of HR and SBP, any changes of these parameters can affect the RPP value. However, reducing the RPP may be more related with reduction in SBP. The present study does not provide explanations about the mechanism by which SE simultaneously decreases post-exercise BP and RPP in normotensive young women. One possible mechanism underlying the effects of SE on BP and RPP may be related to the decrease in sympathetic nervous activity. Pagani et al. found a significant decrease in low-frequency component SBP (a marker of vascular sympathetic activity) after stretching training (Pagani et al., 1997). Another possible mechanism is an improved endothelial-mediate vasodilation. A recent

study showed that a 15 min SE acutely improved NO-mediated vasodilation (Hotta et al., 2013). Nevertheless, it is possible that SE via releasing of NO and decrease in sympathetic nervous activity contributed to PEH and fall in RPP. The study has limitations that should be considered. The small sample size that was utilized and the voluntary in this study were normotensive women, limiting the extrapolation of these to individual with other characteristics. Moreover, PEH mechanisms were not investigated. Additionally, blood sample measures should be considered for future researches, including the analysis of vasodilator metabolites, particularly NO. The assessment of sympathetic nervous activity following SE.

Conclusions

In normotensive women, SE decreases BP and RPP during the post-exercise period. This decrease persist for 30 min. These findings, if reproducible in hypertensive subjects, may have clinical implications.

Acknowledgments

The authors gratefully acknowledge the all subjects whom cooperated in this investigation.

References

- Cardoso Jr. C. G., Gomides R. S., Queiroz A. C. C., Pinto L. G., Lobo F. D. S., Tinucci T., & Forjaz C. L. D. M. Acute and chronic effects of aerobic and resistance exercise on ambulatory blood pressure. *Clinics*, 2010, vol. 65(3), pp. 317-325.
- Chobanian A. V., Bakris G. L., Black H. R., Cushman W. C., Green L. A., Izzo Jr J. L., & National High Blood Pressure Education Program Coordinating Committee. The seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure: the JNC 7 report. *Jama*, 2003, vol. 289(19), pp. 2560-2571.
- Cipriani D. J., Terry M. E., Haines M. A., Tabibnia A. P., & Lyssanova O. Effect of stretch frequency and sex on the rate of gain and rate of loss in muscle flexibility during a hamstring-stretching program: a randomized single-blind longitudinal study. *The Journal of Strength & Conditioning Research*, 2012, vol. 26(8), pp. 2119-2129.
- Cole C. R., Blackstone E. H., Pashkow F. J., Snader C. E., & Lauer M. S. Heart-rate recovery immediately after exercise as a predictor of mortality. *New England Journal of Medicine*, 1999, vol. 341(18), pp. 1351-1357.
- Corrick K. L., Hunter G. R., Fisher G., & Glasser S. P. Changes in vascular hemodynamics in older women following 16 weeks of combined aerobic and resistance training. *The Journal of Clinical Hypertension*, 2013, vol. 15(4), pp. 241-246.
- Drew R. C., Bell M. P., & White M. J. Modulation of spontaneous baroreflex control of heart rate and indexes of vagal tone by passive calf muscle stretch during graded metaboreflex activation in humans. *Journal of Applied Physiology*, 2008, vol. 104(3), pp. 716-723.
- Farinatti P. T., Soares P. P., Monteiro W. D., Duarte A. F., & Castro L. A. Cardiovascular responses to passive static flexibility exercises are influenced by the stretched muscle mass and the Valsalva maneuver. *Clinics*, 2011, vol. 66(3), pp. 459-464.
- Fletcher G. F., Balady G., Froelicher V. F., Hartley L. H., Haskell W. L., & Pollock M. L. Exercise standards a statement for healthcare professionals from the American Heart Association. *Circulation*, 1995, vol. 91(2), pp. 580-615.
- Franklin S. S., Khan S. A., Wong N. D., Larson M. G., & Levy D. Is pulse pressure useful in predicting risk for coronary heart disease? The Framingham Heart Study. *Circulation*, 1999, vol. 100(4), pp. 354-360.
- Goldberger J. J., Le F. K., Lahiri M., Kannankeril P. J., Ng J., & Kadish A. H. Assessment of parasympathetic reactivation after exercise. *American Journal of Physiology-Heart and Circulatory Physiology*, 2006, vol. 290(6), pp. 2446-2452.
- Halliwill J. R., Buck T. M., Laceywell A. N., & Romero S. A. Postexercise hypotension and sustained postexercise vasodilation: what happens after we exercise?. *Experimental physiology*, 2013, vol. 98(1), pp. 7-18.
- Hotta K., Kamiya K., Shimizu R., Yokoyama M., Nakamura-Ogura M., Tabata M., & Masuda T. Stretching Exercises Enhance Vascular Endothelial Function and Improve Peripheral Circulation in Patients With Acute Myocardial Infarction. *International heart journal*, 2013, vol. 54(2), pp. 59-63.
- Keese F., Farinatti P., Pescatello L., Cunha F. A., & Monteiro W. D. Aerobic exercise intensity influences hypotension following concurrent exercise sessions. *International journal of sports medicine*, 2012, vol. 33(2), pp. 148.
- Kuebler W. M., Uhlig U., Goldmann T., Schael G., Kerem A., Exner K., & Uhlig S. Stretch activates nitric oxide production in pulmonary vascular endothelial cells in situ. *American journal of respiratory and critical care medicine*, 2003, vol. 168(11), pp. 1391-1398.
- Lansman J. B., Hallam T. J., & Rink T. J. single stretch-activated ion channels in vascular endothelial cells as mechanotransducer?. *Nature*, 1987, vol. 325, pp. 811-3.
- Lockwood J. M., Pricher M. P., Wilkins B. W., Holowatz L. A., & Halliwill J. R. Postexercise hypotension is not explained by a prostaglandin-dependent peripheral vasodilation. *Journal of Applied Physiology*, 2005, vol. 98(2), pp. 447-453.
- MacDonald J. R. Potential causes, mechanisms, and implications of post exercise hypotension. *Journal of human hypertension*, 2002, vol. 16(4), pp. 225-236.
- Melo C. M., Alencar Filho A. C., Tinucci T., Mion Jr. D., & Forjaz C. L. Postexercise hypotension induced by low-intensity resistance exercise in hypertensive women receiving captopril. *Blood pressure monitoring*, 2006, vol. 11(4), pp. 183-189.
- Mohebbi H., Rahmani-Nia F., Vatani D. S., & Faraji H. Post-exercise responses in blood pressure, heart rate and rate pressure product in endurance and resistance exercise. *Medicina dello Sport*, 2010, vol. 63(2), pp. 209-219.
- Moraes M. R., Bacurau R. F. P., Simoes H. G., Campbell C. S. G., Pudo M. A., Wasinski F., & Araujo R. C. Effect of 12 weeks of resistance exercise on post-exercise hypotension in stage I hypertensive individuals. *Journal of human hypertension*, 2011, vol. 26(9), pp. 533-539.
- Nagpal S., Walia L., Lata H., Sood N., & Ahuja G. K. Effect of exercise on rate pressure product in premenopausal and postmenopausal women with coronary artery disease. *Indian journal of physiology and pharmacology*, 2007, vol. 51(3), pp. 279.
- Pagani M., Montano N., Porta A., Malliani A., Abboud F. M., Birkett C., & Somers V. K. Relationship between spectral components of cardiovascular variabilities and direct measures of muscle sympathetic nerve activity in humans. *Circulation*, 1997, vol. 95(6), pp. 1441-1448.
- Pescatello L. S., Franklin B. A., Fagard, R., Farquhar W. B., Kelley G. A., & Ray C. A. Exercise and hypertension: American College of Sports

- Medicine Position Stand. *Medicine & Science in Sports & Exercise*, 2004, vol. 36(3), pp. 533-53.
24. Piotrowicz E., Baranowski R., Piotrowska M., Zielinski T., & Piotrowicz R. Variable effects of physical training of heart rate variability, heart rate recovery, and heart rate turbulence in chronic heart failure. *Pacing and Clinical Electrophysiology*, 2009, vol. 32(s1), pp. S113-S115.
25. Rezk C. C., Marrache R. C. B., Tinucci T., Mion Jr D., & Forjaz C. Post-resistance exercise hypotension, hemodynamics, and heart rate variability: influence of exercise intensity. *European journal of applied physiology*, 2006, vol. 98(1), pp. 105-112.
26. Santana H. A., Moreira S. R., Asano R. Y., Sales M. M., Cordova C., Campbell C. S., & Simoes H. G. Exercise intensity modulates nitric oxide and blood pressure responses in hypertensive older women. *Aging clinical and experimental research*, 2013, vol. 25(1), pp. 43-48.
27. Singer H. A., & Peach M. J. Calcium-and endothelial-mediated vascular smooth muscle relaxation in rabbit aorta. *Hypertension*, 1982, vol. 4(3), pp. 19-25.
28. Terblanche E., & Millen A. M. The magnitude and duration of post-exercise hypotension after land and water exercises. *European journal of applied physiology*, 2012, vol.112(12), pp. 4111-4118.
29. Wong A., & Figueroa A. Eight weeks of stretching training reduces aortic wave reflection magnitude and blood pressure in obese postmenopausal women. *Journal of human hypertension*, 2013, pp. 1-5.

Информация об авторах:

Хамид Мохебби: ORCID: 0000-0001-9858-5964; baset_marofy@yahoo.com ; Университет Гилян; п/я 1841, Рашт, Иран.

Абдубасет Мароофи: ORCID: 0000-0003-1524-9530; baset_marofy@yahoo.com ; Университет Гилян; п/я 1841, Рашт, Иран.

Назанин Анасри: ORCID: 0000-0001-5273-5963; baset_marofy@yahoo.com ; Университет Гилян; п/я 1841, Рашт, Иран.

Абоозар Джорбониан: ORCID: 0000-0002-9910-4728; baset_marofy@yahoo.com ; Университет Гилян; п/я 1841, Рашт, Иран

Цитируйте эту статью как: Хамид Мохебби, Абдубасет Мароофи, Назанин Анасри, Абоозар Джорбониан. Влияние упражнений на растяжение на гемодинамический статус и профилактику гипотонии у студентов девушек, имеющих нормальное артериальное давление // Физическое воспитание студентов. – 2014. – № 5 – С. 53-58. doi:10.15561/20755279.2014.0510

Электронная версия этой статьи является полной и может быть найдена на сайте: <http://www.sportpedu.org.ua/html/arhive.html>

Это статья Открытого Доступа распространяется под терминами Creative Commons Attribution License, которая разрешает неограниченное использование, распространение и копирование любыми средствами, обеспечивающими должное цитирование этой оригинальной статьи (<http://creativecommons.org/licenses/by/3.0/deed.ru>).

Дата поступления в редакцию: 25.06.2014 г.
Опубликовано: 30.06.2014 г.

Information about the authors:

Hamid Mohebbi: ORCID: 0000-0001-9858-5964; baset_marofy@yahoo.com ; University of Guilan; P.O. Box 1841, Rasht, Iran.

Abdubaset Maroofi: ORCID: 0000-0003-1524-9530; baset_marofy@yahoo.com ; University of Guilan; P.O. Box 1841, Rasht, Iran.

Nazanin Anasri: ORCID: 0000-0001-5273-5963; baset_marofy@yahoo.com ; University of Guilan; P.O. Box 1841, Rasht, Iran.

Aboozar Jorbonian: ORCID: 0000-0002-9910-4728; baset_marofy@yahoo.com ; University of Guilan; P.O. Box 1841, Rasht, Iran.

Cite this article as: Hamid Mohebbi, Abdubaset Maroofi, Nazanin Anasri, Aboozar Jorbonian. The effects of stretching exercise on hemodynamic responses and post-exercise hypotension in normotensive women students. *Physical education of students*, 2014, vol.5, pp. 53-58. doi:10.15561/20755279.2014.0510

The electronic version of this article is the complete one and can be found online at: <http://www.sportpedu.org.ua/html/arhive-e.html>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited (<http://creativecommons.org/licenses/by/3.0/deed.en>).

Received: 25.06.2014
Published: 30.06.2014