

УДК 519.21

О.Д. Борисенко¹, к.ф.-м.н., доцент
О.В. Борисенко², к.ф.-м.н., доцент

Асимптотична поведінка автономної коливної системи четвертого порядку під дією багатовимірного білого шуму і нецентралованого пауссонівського шуму

¹Київський національний університет імені Тараса Шевченка, 01033, Київ, вул. Володимирська, 64.

²Національний технічний університет України "КПІ", 03056, Київ, просп. Перемоги, 37
e-mail: ¹odb@univ.kiev.ua

У роботі вивчається асимптотична поведінка автономної коливної системи, яка описується диференціальним рівнянням четвертого порядку з малими нелінійними збуреннями типу багатовимірного "білого" та нецентралованого Пауссонівського шумів. Розглядається випадок пари дійсних і пари спряжених чисто уявних коренів характеристичного рівняння.

Ключові слова: асимптотична поведінка, автономна коливна система, стохастичне диференціальне рівняння.

The asymptotic behavior of autonomous oscillating system described by differential equation of fourth order with small non-linear external perturbations of multidimensional "white noise" and non-centered "Poisson noise" types is studied. Every term of external perturbations has own order of small parameter ε . If small parameter is equal to zero, then general solution of obtained non-stochastic fourth order differential equation has an oscillating part. We consider given differential equation with external stochastic perturbations as the system of stochastic differential equations and study the limit behavior of its solution at the time moment t/ε^k , as $\varepsilon \rightarrow 0$. The system of averaging stochastic differential equations is derived and its dependence on the order of small parameter in every term of external perturbations is studied. The case of pair of real and pair of pure imaginary conjugate roots of characteristic equation is considered.

Key Words: asymptotic behavior, autonomous oscillating system, stochastic differential equation.

Communicated by Prof. Kozachenko Yu.V.

1 Introduction

The averaging method proposed by N.M.Krylov, N.N.Bogolyubov and Yu.A.Mytropolskij ([1], [2]) is one of the main tool in studying of the deterministic oscillating systems under the action of small non-linear perturbations. The autonomous and non-autonomous oscillating systems of second order under the action of "white noise" and Poisson type noise perturbations are studied in the papers of O.V.Borysenko ([4]). Particular case of the third order oscillating systems are investigated in articles of O.D.Borysenko, O.V.Borysenko ([5]), O.D.Borysenko, O.V.Borysenko and I.G.Malyshev ([6]). Limit behavior of autonomous and non-autonomous general type third order oscillating

O.D. Borysenko¹, Ph.D., Associate Professor
O.V. Borysenko², Ph.D., Associate Professor
Asymptotic behavior of autonomous oscillating system under the action of multidimensional white noise and non-centered Poisson noise

¹Taras Shevchenko National University of Kyiv, 01033, Kyiv, 64 Volodymyrska st.,

²National Technical University of Ukraine "KPI", 03056, Kyiv, 37 Peremogy prospect
e-mail: ¹odb@univ.kiev.ua

system under the action of external small nonlinear random disturbances such as multidimensional "white noise" and "Poisson noise" was studied in ([7], [8]).

This paper deals with investigation of the behavior, as $\varepsilon \rightarrow 0$, of the fourth order autonomous oscillating system driven by stochastic differential equation

$$\begin{aligned} &x^{IV}(t) + b_1x'''(t) + b_2x''(t) + b_3x'(t) + b_4x(t) = \\ &= \varepsilon^{k_0}f_0(x(t), x'(t), x''(t), x'''(t)) + \\ &+ f_\varepsilon(x(t), x'(t), x''(t), x'''(t)) \end{aligned} \tag{1}$$

with non-random initial conditions $x(0) = x_0^{(1)}$, $x'(0) = x_0^{(2)}$, $x''(0) = x_0^{(3)}$, $x'''(0) = x_0^{(4)}$, where $\varepsilon > 0$ is a small parameter, $f_\varepsilon(x, x', x'', x''')$ is a

random function such that

$$\begin{aligned} \int_0^t f_\varepsilon(x(s), x'(s), x''(s), x'''(s)) ds &= \sum_{i=1}^m \varepsilon^{k_i} \times \\ &\times \int_0^t f_i(x(s), x'(s), x''(s), x'''(s)) dw_i(s) + \varepsilon^{k_{m+1}} \times \\ &\times \int_0^t \int_{\mathbb{R}} f_{m+1}(x(s), x'(s), x''(s), x'''(s), z) \nu(ds, dz), \end{aligned}$$

$k_i > 0, i = \overline{0, m+1}$; f_i are non-random functions $i = \overline{0, m+1}$; $w_i(t), i = \overline{1, m}$ are independent one-dimensional Wiener processes; $\nu(dt, dy)$ is the Poisson measure independent on $w_i(t), i = \overline{1, m}$, $E\nu(dt, dy) = \Pi(dy)dt$, $\tilde{\nu}(dt, dy) = \nu(dt, dy) - \Pi(dy)dt$, $\Pi(A)$ is a finite measure on Borel sets $A \in \mathbb{R}$.

We will consider the equation (1) as the system of stochastic differential equations

$$\begin{aligned} dy_i(t) &= y_{i+1}(t)dt, \quad i = \overline{1, 3} \\ dy_4(t) &= \left[-(b \cdot y(t)) + \varepsilon^{k_0} f_0(y(t)) + \right. \\ &\quad \left. + \varepsilon^{k_{m+1}} \int_{\mathbb{R}} f_{m+1}(y(t), z) \Pi(dz) \right] dt + \\ &\quad + \sum_{i=1}^m \varepsilon^{k_i} f_i(y(t)) dw_i(t) + \\ &\quad + \varepsilon^{k_{m+1}} \int_{\mathbb{R}} f_{m+1}(y(t), z) \tilde{\nu}(dt, dz), \\ y(t) &= \{y_i(t)\}_{i=\overline{1, 4}}, b = (b_4, b_3, b_2, b_1), \\ y_i(0) &= x_0^{(i)}, \quad i = \overline{1, 4} \end{aligned} \tag{2}$$

$(b \cdot y(t))$ – is a scalar product of vectors b and $y(t)$.

In what follows we will use the constant $K > 0$ for the notation of different constants, which are not depend on ε .

2 Main result

We will study the asymptotic behavior of given oscillating system, as $\varepsilon \rightarrow 0$, in the case when there exists stable harmonic oscillations at the system under condition $\varepsilon = 0$. Under this condition corresponding characteristic equation has a form

$$\lambda^4 + b_1 \lambda^3 + b_2 \lambda^2 + b_3 \lambda + b_4 = 0.$$

This paper deals with the following case:

$$b_1 > 0, b_3 > 0, b_1 b_2 > b_3, b_1^2 > 4(b_2 - \frac{b_3}{b_1}),$$

$b_1^2 b_4 = b_3(b_1 b_2 - b_3)$. Characteristic equation has a roots

$$\begin{aligned} \lambda_1 &= -\eta_1, \quad \lambda_2 = -\eta_2, \quad \lambda_{3,4} = \pm i\omega, \text{ where} \\ \eta_{1,2} &= \frac{1}{2} \left(b_1 \pm \sqrt{b_1^2 - 4 \left(b_2 - \frac{b_3}{b_1} \right)} \right), \quad \omega^2 = \frac{b_3}{b_1}. \end{aligned}$$

If $\varepsilon = 0$ then the equation (1) has general solution in the form

$$x(t) = C_1 e^{-\eta_1 t} + C_2 e^{-\eta_2 t} + A_1 \cos \omega t + A_2 \sin \omega t$$

Let us consider the following representation of the process $y(t)$:

$$\begin{aligned} y_1(t) &= N_1(t) + N_2(t) + \\ &\quad + A_1(t) \cos \omega t + A_2(t) \sin \omega t, \\ y_2(t) &= -\eta_1 N_1(t) - \eta_2 N_2(t) - \\ &\quad - A_1(t) \omega \sin \omega t + A_2(t) \omega \cos \omega t, \\ y_3(t) &= \eta_1^2 N_1(t) + \eta_2^2 N_2(t) - \\ &\quad - A_1(t) \omega^2 \cos \omega t - A_2(t) \omega^2 \sin \omega t, \\ y_4(t) &= -\eta_1^3 N_1(t) - \eta_2^3 N_2(t) + \\ &\quad + A_1(t) \omega^3 \sin \omega t - A_2(t) \omega^3 \cos \omega t, \\ N_i(t) &= C_i(t) \exp\{-\eta_i t\}, \quad i = 1, 2. \end{aligned} \tag{3}$$

We can solve the system of linear equations (3) with respect to $(N_1(t), N_2(t), A_1(t), A_2(t))$ and using the Ito formula we derive the system of stochastic differential equations:

$$\begin{aligned} dN_1(t) &= -\eta_1 N_1(t) dt + \frac{1}{(\eta_2 - \eta_1)(\eta_1^2 + \omega^2)} dH(t), \\ dN_2(t) &= -\eta_2 N_2(t) dt - \frac{1}{(\eta_2 - \eta_1)(\eta_2^2 + \omega^2)} dH(t), \\ dA_1(t) &= \frac{-\omega(\eta_1 + \eta_2) \cos \omega t + (\omega^2 - \eta_1 \eta_2) \sin \omega t}{\omega(\eta_1^2 + \omega^2)(\eta_2^2 + \omega^2)} dH(t), \\ dA_2(t) &= \frac{-\omega(\eta_1 + \eta_2) \sin \omega t - (\omega^2 - \eta_1 \eta_2) \cos \omega t}{\omega(\eta_1^2 + \omega^2)(\eta_2^2 + \omega^2)} dH(t), \end{aligned} \tag{4}$$

$$\begin{aligned} dH(t) &= \left[\varepsilon^{k_0} \tilde{f}_0(\omega t, \xi(t)) + \right. \\ &\quad \left. + \varepsilon^{k_{m+1}} \int_{\mathbb{R}} \tilde{f}_{m+1}(\omega t, \xi(t), z) \Pi(dz) \right] dt + \\ &\quad + \sum_{i=1}^m \varepsilon^{k_i} \tilde{f}_i(\omega t, \xi(t)) dw_i(t) + \\ &\quad + \varepsilon^{k_{m+1}} \int_{\mathbb{R}} \tilde{f}_{m+1}(\omega t, \xi(t), z) \tilde{\nu}(dt, dz), \end{aligned}$$

where $\xi(t) = (N_1(t), N_2(t), A_1(t), A_2(t))$, $\tilde{f}_i(\omega t, N_1, N_2, A_1, A_2), i = \overline{0, m}$ are obtained from $f_i(y(t)), i = \overline{0, m}$ and $\tilde{f}_{m+1}(\omega t, N_1, N_2, A_1, A_2, z)$ is obtained from $f_{m+1}(y(t), z)$ using (3).

Theorem 2.1. Let $\Pi(\mathbb{R}) < \infty$, $t \in [0, t_0]$, $k = \min(k_0, 2k_1, \dots, 2k_m, k_{m+1})$. Let us suppose, that functions $f_j, j = \overline{0, m+1}$ bounded and satisfy Lipschitz condition on $y_i, i = \overline{1, 4}$. If given below matrix $\bar{\sigma}^2(A_1, A_2)$ is non-negative definite, then

1. If $k_0 = 2k_i = k_{m+1}$, $i = \overline{1, m}$ then the stochastic process $\xi_\varepsilon(t) = \xi(t/\varepsilon^k)$ weakly converges, as $\varepsilon \rightarrow 0$, to the stochastic process $\bar{\xi}(t) = (0, 0, \bar{A}_1(t), \bar{A}_2(t))$, where $\bar{A}(t) = (\bar{A}_1(t), \bar{A}_2(t))$ is the solution to the system of stochastic differential equations

$$d\bar{A}(t) = \bar{\alpha}(\bar{A}(t))dt + \bar{\sigma}(\bar{A}(t))d\bar{w}(t), \quad (5)$$

$$\bar{A}(0) = (A_1(0), A_2(0)),$$

where

$$\bar{\alpha}(A_1, A_2) = \frac{1}{2\pi} \int_0^{2\pi} \hat{f}_{(1)}(\phi, A_1, A_2) \Psi(\phi) d\phi,$$

$$\begin{aligned} \bar{\sigma}(A_1, A_2) &= \left\{ \bar{B}(A_1, A_2) \right\}^{\frac{1}{2}} = \\ &= \left\{ \frac{1}{2\pi} \int_0^{2\pi} \hat{f}_{(2)}(\phi, A_1, A_2) \Psi(\phi) \Psi^T(\phi) d\phi \right\}^{\frac{1}{2}} \end{aligned}$$

$$\begin{aligned} \Psi(\phi) &= \frac{1}{\omega(\eta_1^2 + \omega^2)(\eta_2^2 + \omega^2)} \times \\ &\times \begin{pmatrix} -\omega(\eta_1 + \eta_2) \cos \phi + (\omega^2 - \eta_1 \eta_2) \sin \phi \\ -\omega(\eta_1 + \eta_2) \sin \phi - (\omega^2 - \eta_1 \eta_2) \cos \phi \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \hat{f}_{(1)}(\phi, A_1, A_2) &= \tilde{f}_0(\phi, 0, 0, A_1, A_2) + \\ &+ \int_{\mathbb{R}} \tilde{f}_{m+1}(\phi, 0, 0, A_1, A_2, z) \Pi(dz), \end{aligned}$$

$$\hat{f}_{(2)}(\phi, A_1, A_2) = \sum_{i=1}^m \tilde{f}_i^2(\phi, 0, 0, A_1, A_2),$$

$\Psi^T(\phi)$ is the vector transpose to vector $\Psi(\phi)$, $\bar{w}(t) = (\bar{w}_i(t), i = 1, 2)$, $\bar{w}_i(t), i = 1, 2$ – independent one-dimensional Wiener processes.

2. If $k < k_0$ then in the averaging equation (5) we must put $\tilde{f}_0 \equiv 0$; if $k < 2k_j$ for some $1 \leq j \leq m$, then in the averaging equation (5) we must put $\tilde{f}_j \equiv 0$ for all such j ; if $k < k_{m+1}$ then in the averaging equation (5) we must put $\tilde{f}_{m+1} \equiv 0$.

Доведення. Let us make a change of variable $t \rightarrow t/\varepsilon^k$ at the system (4) and obtain for the process $\xi_\varepsilon(t) = (N_1^\varepsilon(t), N_2^\varepsilon(t), A_1^\varepsilon(t), A_2^\varepsilon(t)) = (N_1(t/\varepsilon^k), N_2(t/\varepsilon^k), A_1(t/\varepsilon^k), A_2(t/\varepsilon^k))$ the system of stochastic differential equations

$$\begin{aligned} dN_1^\varepsilon(t) &= -\frac{\eta_1}{\varepsilon^k} N_1^\varepsilon(t) dt + \frac{1}{(\eta_2 - \eta_1)(\eta_1^2 + \omega^2)} dH_\varepsilon(t), \\ dN_2^\varepsilon(t) &= -\frac{\eta_2}{\varepsilon^k} N_2^\varepsilon(t) dt - \frac{1}{(\eta_2 - \eta_1)(\eta_2^2 + \omega^2)} dH_\varepsilon(t), \\ dA_1^\varepsilon(t) &= \Psi_1(\omega t/\varepsilon^k) dH_\varepsilon(t), \\ dA_2^\varepsilon(t) &= \Psi_2(\omega t/\varepsilon^k) dH_\varepsilon(t), \end{aligned}$$

$$\begin{aligned} dH_\varepsilon(t) &= \left[\varepsilon^{k_0-k} \tilde{f}_0\left(\frac{\omega t}{\varepsilon^k}, \xi_\varepsilon(t)\right) + \right. \\ &\quad \left. + \varepsilon^{k_{m+1}-k} \int_{\mathbb{R}} \tilde{f}_{m+1}\left(\frac{\omega t}{\varepsilon^k}, \xi_\varepsilon(t), z\right) \Pi(dz) \right] dt + \\ &\quad + \sum_{i=1}^m \varepsilon^{k_i-k/2} \tilde{f}_i\left(\frac{\omega t}{\varepsilon^k}, \xi_\varepsilon(t)\right) dw_i^\varepsilon(t) + \\ &\quad + \varepsilon^{k_{m+1}} \int_{\mathbb{R}} \tilde{f}_{m+1}\left(\frac{\omega t}{\varepsilon^k}, \xi_\varepsilon(t), z\right) \tilde{\nu}_\varepsilon(dt, dz), \end{aligned}$$

where $\Psi_i(\phi), i = 1, 2$ is a components of vector $\Psi(\phi)$, $w_i^\varepsilon(t) = \varepsilon^{k/2} w_i(t/\varepsilon^k)$, $\tilde{\nu}_\varepsilon(t, A) = \nu(t/\varepsilon^k, A) - \Pi(A)/\varepsilon^k$, here A is a Borel set in \mathbb{R} . For each $\varepsilon > 0$ the process $w_i^\varepsilon(t), i = \overline{1, m}$ are independent one-dimensional Wiener processes, and $\tilde{\nu}_\varepsilon(t, A)$ is the centered Poisson measure independent on $w_i^\varepsilon(t), i = \overline{1, m}$.

We have $N_i^\varepsilon(t) = \exp\{-\eta_i t/\varepsilon^k\} C_i(t/\varepsilon^k)$, $i = 1, 2$ and processes $C_i^\varepsilon(t) = C_i(t/\varepsilon^k)$, $i = 1, 2$ satisfy the stochastic differential equations

$$\begin{aligned} dC_1^\varepsilon(t) &= \frac{e^{\eta_1 t/\varepsilon^k}}{(\eta_2 - \eta_1)(\eta_1^2 + \omega^2)} dH_\varepsilon(t), \\ dC_2^\varepsilon(t) &= -\frac{e^{\eta_2 t/\varepsilon^k}}{(\eta_2 - \eta_1)(\eta_2^2 + \omega^2)} dH_\varepsilon(t), \end{aligned}$$

where $|C_i^\varepsilon(0)| \leq K, i = 1, 2$.

From boundedness of functions $f_i, i = \overline{0, m+1}$ and condition $\Pi(\mathbb{R}) < \infty$ we can obtain the estimate

$$\begin{aligned} \mathbb{E}|N_i^\varepsilon(t)|^2 &\leq K [e^{-2\eta_i t/\varepsilon^k} + \varepsilon^k (1 - e^{-2\eta_i t/\varepsilon^k}) \times \\ &\times (t(\varepsilon^{2(k_0-k)} + \varepsilon^{2(k_{m+1}-k)}) + \sum_{i=1}^{m+1} \varepsilon^{2k_i-k})], \\ i &= 1, 2. \end{aligned}$$

Therefore $\lim_{\varepsilon \rightarrow 0} \mathbb{E}|N_i^\varepsilon(t)|^2 = 0, i = 1, 2$ and it is sufficient to study the behavior, as $\varepsilon \rightarrow 0$, of solution to the system of stochastic differential equations

$$dA_i^\varepsilon(t) = \Psi_i(\omega t/\varepsilon^k) d\hat{H}_\varepsilon(t), \quad i = 1, 2 \quad (6)$$

with initial conditions $A_1^\varepsilon(0) = A_1(0)$, $A_2^\varepsilon(0) = A_2(0)$, where

$$\begin{aligned} d\hat{H}_\varepsilon(t) &= \left[\varepsilon^{k_0-k} \tilde{f}_0\left(\frac{\omega t}{\varepsilon^k}, A_1^\varepsilon(t), A_2^\varepsilon(t)\right) + \right. \\ &\quad \left. + \varepsilon^{k_{m+1}-k} \int_{\mathbb{R}} \tilde{f}_{m+1}\left(\frac{\omega t}{\varepsilon^k}, A_1^\varepsilon(t), A_2^\varepsilon(t), z\right) \Pi(dz) \right] dt + \\ &\quad + \sum_{i=1}^m \varepsilon^{k_i-k/2} \tilde{f}_i\left(\frac{\omega t}{\varepsilon^k}, A_1^\varepsilon(t), A_2^\varepsilon(t)\right) dw_i^\varepsilon(t) + \\ &\quad + \varepsilon^{k_{m+1}} \int_{\mathbb{R}} \tilde{f}_{m+1}\left(\frac{\omega t}{\varepsilon^k}, A_1^\varepsilon(t), A_2^\varepsilon(t), z\right) \tilde{\nu}_\varepsilon(dt, dz), \end{aligned}$$

$$\begin{aligned}\hat{f}_j(t, A_1, A_2) &= \tilde{f}_j(t, 0, 0, A_1, A_2), \quad j = \overline{0, m} \\ \hat{f}_{m+1}(t, A_1, A_2, z) &= \tilde{f}_{m+1}(t, 0, 0, A_1, A_2, z).\end{aligned}$$

Let us denote $A_\varepsilon(t) = (A_1^\varepsilon(t), A_2^\varepsilon(t))$. Using conditions on coefficients of equation (6) and properties of stochastic integrals we obtain estimates

$$\begin{aligned}\mathbb{E}\|A_\varepsilon(t)\|^2 &\leq K[1 + t^2(\varepsilon^{2(k_0-k)} + \\ &+ \varepsilon^{2(k_{m+1}-k)}) + t \sum_{i=1}^{m+1} \varepsilon^{2k_i-k}], \\ \mathbb{E}\|A_\varepsilon(t) - A_\varepsilon(s)\|^2 &\leq K[|t-s|^2(\varepsilon^{2(k_0-k)} + \\ &+ \varepsilon^{2(k_{m+1}-k)}) + |t-s| \sum_{i=1}^{m+1} \varepsilon^{2k_i-k}].\end{aligned}$$

Similarly for the process $\zeta_\varepsilon(t) = (\zeta_1^\varepsilon(t), \zeta_2^\varepsilon(t))$, where

$$\zeta_i^\varepsilon(t) = \int_0^t \Psi_i\left(\frac{\omega s}{\varepsilon^k}\right) dM_\varepsilon(s), \quad i = 1, 2,$$

$$\begin{aligned}dM_\varepsilon(t) &= \\ &= \sum_{i=1}^m \varepsilon^{k_i-k/2} \hat{f}_i\left(\frac{\omega t}{\varepsilon^k}, A_1^\varepsilon(t), A_2^\varepsilon(t)\right) dw_i^\varepsilon(t) + \\ &+ \varepsilon^{k_{m+1}} \int_{\mathbb{R}} \hat{f}_{m+1}\left(\frac{\omega t}{\varepsilon^k}, A_1^\varepsilon(t), A_2^\varepsilon(t), z\right) \tilde{\nu}_\varepsilon(dt, dz),\end{aligned}$$

we derive estimates

$$\mathbb{E}\|\zeta_\varepsilon(t)\|^2 \leq Kt \sum_{i=1}^{m+1} \varepsilon^{2k_i-k},$$

$$\mathbb{E}\|\zeta_\varepsilon(t) - \zeta_\varepsilon(s)\|^2 \leq K|t-s| \sum_{i=1}^{m+1} \varepsilon^{2k_i-k}.$$

Therefore for stochastic process $\eta_\varepsilon(t) = (A_\varepsilon(t), \zeta_\varepsilon(t))$ conditions of weak compactness [9] are fulfilled

$$\lim_{h \downarrow 0} \overline{\lim}_{\varepsilon \rightarrow 0} \sup_{|t-s| < h} P\{|\eta_\varepsilon(t) - \eta_\varepsilon(s)| > \delta\} = 0$$

for any $\delta > 0$, $t, s \in [0, T]$,

$$\lim_{N \rightarrow \infty} \overline{\lim}_{\varepsilon \rightarrow 0} \sup_{t \in [0, T]} P\{|\eta_\varepsilon(t)| > N\} = 0,$$

and for any sequence $\varepsilon_n \rightarrow 0$, $n = 1, 2, \dots$ there exists a subsequence $\varepsilon_m = \varepsilon_{n(m)} \rightarrow 0$, $m = 1, 2, \dots$, probability space, stochastic processes $A_{\varepsilon_m}(t) = (\bar{A}_1^{\varepsilon_m}(t), \bar{A}_2^{\varepsilon_m}(t))$, $\zeta_{\varepsilon_m}(t)$, $\bar{A}(t) = (\bar{A}_1(t), \bar{A}_2(t))$, $\bar{\zeta}(t)$ defined on this space, such that $\bar{A}_{\varepsilon_m}(t) \rightarrow \bar{A}(t)$, $\zeta_{\varepsilon_m}(t) \rightarrow \bar{\zeta}(t)$ in probability, as $\varepsilon_m \rightarrow 0$, and finite-dimensional distributions of $\bar{A}_{\varepsilon_m}(t)$, $\bar{\zeta}_{\varepsilon_m}(t)$ are coincide with finite-dimensional

distributions of $A_{\varepsilon_m}(t)$, $\zeta_{\varepsilon_m}(t)$. Since we interesting in limit behaviour of distributions, we can consider processes $A_{\varepsilon_m}(t)$, and $\zeta_{\varepsilon_m}(t)$ instead of $\bar{A}_{\varepsilon_m}(t)$, $\bar{\zeta}_{\varepsilon_m}(t)$. From (6) we obtain equation

$$\begin{aligned}A_{\varepsilon_m}(t) &= A(0) + \int_0^t \alpha_{\varepsilon_m}(s, A_{\varepsilon_m}(s)) ds + \zeta_{\varepsilon_m}(t), \\ A(0) &= (A_1(0), A_2(0)),\end{aligned}\tag{7}$$

where $\alpha_\varepsilon(t, A) = (\alpha_1^\varepsilon(t, A_1, A_2), \alpha_2^\varepsilon(t, A_1, A_2))$,

$$\alpha_i^\varepsilon(t, A_1, A_2) = \Psi_i\left(\frac{\omega t}{\varepsilon^k}\right) \hat{f}_{(1)}^\varepsilon\left(\frac{\omega t}{\varepsilon^k}, A_1, A_2\right),$$

$$\begin{aligned}\hat{f}_{(1)}^\varepsilon(t, A_1, A_2) &= \varepsilon^{k_0-k} \hat{f}_0(t, A_1, A_2) + \\ &+ \varepsilon^{k_{m+1}-k} \int_{\mathbb{R}} \hat{f}_{m+1}(t, A_1, A_2, z) \Pi(dz).\end{aligned}$$

It should be noted that process $\zeta_\varepsilon(t)$ is the vector-valued square integrable martingale with matrix characteristic

$$\begin{aligned}\langle \zeta_\varepsilon^{(l)}, \zeta_\varepsilon^{(n)} \rangle(t) &= \sum_{j=1}^m \int_0^t \sigma_\varepsilon^{(l,j)}(s, A_1^\varepsilon(s), A_2^\varepsilon(s)) \times \\ &\times \sigma_\varepsilon^{(n,j)}(s, A_1^\varepsilon(s), A_2^\varepsilon(s)) ds + \\ &+ \frac{1}{\varepsilon^k} \int_0^t \int_{\mathbb{R}} \gamma_\varepsilon^{(l)}(s, A_1^\varepsilon(s), A_2^\varepsilon(s), z) \times \\ &\times \gamma_\varepsilon^{(n)}(s, A_1^\varepsilon(s), A_2^\varepsilon(s), z) \Pi(dz) ds, \quad l, n = 1, 2,\end{aligned}$$

where

$$\begin{aligned}\sigma_\varepsilon^{(1,j)}(s, A_1, A_2) &= \\ &= \varepsilon^{k_j-k/2} \Psi_1\left(\frac{\omega s}{\varepsilon^k}\right) \hat{f}_j\left(\frac{\omega s}{\varepsilon^k}, A_1, A_2\right), \\ \sigma_\varepsilon^{(2,j)}(s, A_1, A_2) &= \\ &= \varepsilon^{k_j-k/2} \Psi_2\left(\frac{\omega s}{\varepsilon^k}\right) \hat{f}_j\left(\frac{\omega s}{\varepsilon^k}, A_1, A_2\right), \\ \gamma_\varepsilon^{(1)}(s, A_1, A_2, z) &= \\ &= \varepsilon^{k_{m+1}} \Psi_1\left(\frac{\omega s}{\varepsilon^k}\right) \hat{f}_{m+1}\left(\frac{\omega s}{\varepsilon^k}, A_1, A_2, z\right), \\ \gamma_\varepsilon^{(2)}(s, A_1, A_2, z) &= \\ &= \varepsilon^{k_{m+1}} \Psi_2\left(\frac{\omega s}{\varepsilon^k}\right) \hat{f}_{m+1}\left(\frac{\omega s}{\varepsilon^k}, A_1, A_2, z\right).\end{aligned}$$

For processes $A_\varepsilon(t)$ and $\zeta_\varepsilon(t)$ following estimates hold

$$\begin{aligned}\mathbb{E}\|A_\varepsilon(t) - A_\varepsilon(s)\|^4 &\leq \\ &\leq K[(\varepsilon^{4(k_0-k)} + \varepsilon^{4(k_{m+1}-k)})|t-s|^4 + \\ &+ \mathbb{E}\|\zeta_\varepsilon(t) - \zeta_\varepsilon(s)\|^4],\end{aligned}\tag{8}$$

$$\begin{aligned}\mathbb{E}\|\zeta_\varepsilon(t) - \zeta_\varepsilon(s)\|^4 &\leq \\ &\leq K \left[\sum_{j=1}^{m+1} \varepsilon^{4k_j-2k} |t-s|^2 + \right. \\ &\left. + \varepsilon^{4k_{m+1}-3k/2} |t-s|^{3/2} + \varepsilon^{4k_{m+1}-k} |t-s| \right],\end{aligned}\tag{9}$$

$$\begin{aligned} \mathbb{E}\|A_\varepsilon(t) - A_\varepsilon(s)\|^8 &\leq K, \\ \mathbb{E}\|\zeta_\varepsilon(t) - \zeta_\varepsilon(s)\|^8 &\leq K. \end{aligned} \quad (10)$$

Since $A_{\varepsilon_m}(t) \rightarrow \bar{A}(t)$, $\zeta_{\varepsilon_m}(t) \rightarrow \bar{\zeta}(t)$ in probability, as $\varepsilon_m \rightarrow 0$, then, using (10), from (8) and (9) we obtain estimates

$$\begin{aligned} \mathbb{E}\|\bar{A}(t) - \bar{A}(s)\|^4 &\leq K(|t-s|^4 + |t-s|^2), \\ \mathbb{E}\|\bar{\zeta}(t) - \bar{\zeta}(s)\|^4 &\leq C|t-s|^2. \end{aligned}$$

Therefore processes $\bar{A}(t)$ and $\bar{\zeta}(t)$ satisfy the Kolmogorov's continuity condition [10].

Let us consider the case $k_0 = 2k_j = k_{m+1}$, $j = \overline{1, m}$. Under these conditions we have for $l, n = 1, 2$

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \frac{1}{t} \int_0^t \alpha_\varepsilon^{(l)}(s, A_1, A_2) ds &= \bar{\alpha}^{(l)}(A_1, A_2), \\ \lim_{\varepsilon \rightarrow 0} \frac{1}{t} \int_0^t [\sum_{j=1}^m \sigma_\varepsilon^{(l,j)}(s, A_1, A_2) \sigma_\varepsilon^{(n,j)}(s, A_1, A_2) + \\ &+ \frac{1}{\varepsilon^k} \int_R \gamma_\varepsilon^{(l)}(s, A_1, A_2, z) \times \\ &\times \gamma_\varepsilon^{(n)}(s, A_1, A_2, z) \Pi(dz)] ds &= \bar{B}_{ln}(A_1, A_2), \end{aligned} \quad (11)$$

where functions $\bar{\alpha}^{(i)}(A_1, A_2)$ and $\bar{B}(A_1, A_2) = \{\bar{B}_{ij}(A_1, A_2), i, j = 1, 2\}$ are defined in the condition of theorem. Since processes $\bar{A}(t), \bar{\zeta}(t)$ are continuous, then from (Lemma 1 [7]) and relationships (7), (11) it follows

$$\begin{aligned} \bar{A}(t) &= A(0) + \int_0^t \bar{\alpha}(\bar{A}_1(s), \bar{A}_2(s)) ds + \bar{\zeta}(t), \\ A(0) &= (A_1(0), A_2(0)), \end{aligned} \quad (12)$$

where $\bar{\zeta}(t)$ is continuous vector-valued martingale

Список використаних джерел

1. Крълов Н.М. Введение в нелинейную механику/ Н.М. Крълов, Н.Н. Боголюбов. – Киев: Изд-во АН УССР – 1937. – 363 с.
2. Боголюбов Н.Н. Асимптотические методы в теории нелинейных колебаний/ Н.Н. Боголюбов, Ю.А. Митропольский. – М.: Наука. – 1974. – 501 с.
3. Борисенко О.В. Малые случайные возмущения в колебательных системах второго порядка/ О.В. Борисенко // Український математичний журнал. – 1992. – №1. – С. 11 – 17.

with matrix characteristic

$$\langle \bar{\zeta}^{(i)}, \bar{\zeta}^{(j)} \rangle(t) = \int_0^t \bar{B}_{ij}(\bar{A}_1(s), \bar{A}_2(s)) ds, i, j = 1, 2.$$

Hence [11] there exists Wiener process $\bar{w}(t) = (\bar{w}_i(t), i = 1, 2)$, such that

$$\begin{aligned} \bar{\zeta}(t) &= \int_0^t \bar{\sigma}(\bar{A}_1(s), \bar{A}_2(s)) d\bar{w}(s), \\ \bar{\sigma}(A_1, A_2) &= \{\bar{B}(A_1, A_2)\}^{1/2}. \end{aligned} \quad (13)$$

Relationships (12), (13) mean that process $\bar{A}(t)$ satisfies equation (4). Under conditions of theorem the equation (4) has unique solution. Therefore process $\bar{A}(t)$ does not depend on choosing of sub-sequence $\varepsilon_m \rightarrow 0$, and finite-dimensional distributions of process $A_{\varepsilon_m}(t)$ converge to finite-dimensional distributions of process $\bar{A}(t)$. Since processes $A_{\varepsilon_m}(t)$ and $\bar{A}(t)$ are Markov processes then using the conditions for weak convergence of Markov processes we finish the proof of statement 1 of theorem.

Let us consider the case $k < k_0$ or $k < k_{m+1}$. Then the corresponding terms in the coefficients $\alpha_\varepsilon^{(i)}(t, A_1, A_2)$, $i = 1, 2$ of equation (7) tend to zero, as $\varepsilon \rightarrow 0$.

In the case $k < 2k_j$, $j = \overline{1, m}$ in (11) we have $\sigma_\varepsilon^{(l,j)}(t, A_1, A_2) \sigma_\varepsilon^{(n,j)}(t, A_1, A_2) = O(\varepsilon^{2k_j - k})$, $l, n = 1, 2$. Repeating with obvious modifications the proof of statement 1) of theorem we obtain proof of the statement 2).

□

References

1. KRYLOV, N.M. and BOGOLYUBOV, N.N. (1937), *Vvedenie v nelinejnuyu mehaniku*, Kiev: Izd-vo AN USSR, 363 p.
2. BOGOLYUBOV, N.N. and MITROPOLSKIY, Yu.A.(1974), *Asymptoticheskie metody v teorii nelinejnyh kolebanij*, M.: Nauka, 501 p.
3. BORYSENKO, O.V. (1992), "Malye sluchajnye vozmushcheniya v kolebatelnyh sistemah vtorogo poryadka", *Ukrainskii Matematicheskij Zhurnal*, v.44, №1, pp.11-17.

4. Борисенко О.В. Поведение неавтономной колебательной системы под воздействием малых случайных возмущений в резонансном случае / О.В. Борисенко // Украинский математический журнал. – 1992. – **44**. – №12. – С. 1645 – 1651.
5. Борисенко О.Д. Асимптотическое поведение стохастической динамической системы / О.Д. Борисенко, О.В. Борисенко // Теория оптимальных решений. – Киев: Ин-т кибернетики НАН Украины. – 1999. – С. 38 – 44.
6. Borysenko O.D. Asymptotic behavior of third order random oscillation system in the resonance case / O.D. Borysenko, O.V. Borysenko, I.G. Malyshev // Theory of Stoch. Proc. – 2003. – **9(25)**. – №3-4. – P. 1 – 11.
7. Borysenko O.D. Averaging method for autonomous third order random oscillating system / O.D. Borysenko, O.V. Borysenko, D.O. Borysenko // Bulletin of Taras Shevchenko National University of Kyiv. Series: Physics & Mathematics – 2013. – №4. – P. 7 – 11.
8. Borysenko O.D. Averaging method for non-autonomous third order random oscillating system in non-resonance case / O.D. Borysenko, O.V. Borysenko // Bulletin of Taras Shevchenko National University of Kyiv. Series: Physics & Mathematics – 2014. – №1. – P. 7 – 11
9. Скороход А.В. Исследования по теории случайных процессов / А.В. Скороход // К.: Изд. Киевского университета. – 1961.
10. Гихман И.И. Теория случайных процессов, т.1 /И.И. Гихман, А.В. Скороход // М.: Наука. – 1971. – 664 с.
11. N.Ikeda Stochastic Differential Equations and Diffusion Processes /N.Ikeda, S.Watanabe // North Holland, Amsterdam and Kadansha, Tokyo. – 1981. – 464 p.
4. BORYSENKO, O.V. (1992), "Povedenie neavtonomnoi kolebatelnoj sistemy pod vozdejstviem malyh sluchajnyh vozmushchenij v rezonansnom sluchae", *Ukrainskii Matematicheskij Zhurnal*, **v.44**, №12, pp. 1645 – 1651.
5. BORYSENKO, O.D. and BORYSENKO, O.V. (1999), "Asymptoticheskoe povedenie stohasticheskoy dinamicheskoy sistemy", *Teoriya optimalnyh reshenij*, Kiev, Inst.Kibernetiki NANU, pp. 38 – 44.
6. BORYSENKO, O.D., BORYSENKO, O.V. and MALYSHEV, I.G. (2003), "Asymptotic behavior of third order random oscillation system in the resonance case", *Theory of Stoch. Proc.*, **v.9(25)**, №3-4, pp. 1 – 11.
7. BORYSENKO, O.D., BORYSENKO, O.V. and BORYSENKO, D.O. (2013), "Averaging method for autonomous third order random oscillating system", *Bulletin of Taras Shevchenko National University of Kyiv. Series: Physics & Mathematics*, №4, pp. 7 – 11.
8. BORYSENKO, O.D. and BORYSENKO, O.V. (2014), "Averaging method for non-autonomous third order random oscillating system in non-resonance case", *Bulletin of Taras Shevchenko National University of Kyiv. Series: Physics & Mathematics*, №1, pp. 7 – 11.
9. SKOROKHOD, A.V. (1965) *Studies in the Theory of Random Processes*, Addison-Wesley, 199 p.
10. GIKHMAN, I.I. and SKOROKHOD, A.V. (1971), *Teoriya sluchainyh processov*, v.1, M.: Nauka, 664 p.
11. IKEDA, N. and WATANABE, S. (1981), *Stochastic Differential Equations and Diffusion Processes*, North Holland, Amsterdam and Kadansha, Tokyo, 464 p.

Received: 30.09.2014