

## ABSTRACT AND REFERENCES

## MATHEMATICS AND CYBERNETICS – APPLIED ASPECTS

**DOI: 10.15587/1729-4061.2017.108285**  
**DEVELOPMENT OF THE DESCRIPTIVE**  
**BINARY MODEL AND ITS APPLICATION FOR**  
**IDENTIFICATION OF CLUMPS OF TOXIC**  
**CYANOBACTERIA (p. 4-11)**

**Konstantin Nosov**

V. N. Karazin Kharkiv National University,  
 Kharkiv, Ukraine

**ORCID:** <http://orcid.org/0000-0003-4374-7502>

**Grygoriy Zholtkevych**

V. N. Karazin Kharkiv National University,  
 Kharkiv, Ukraine

**ORCID:** <http://orcid.org/0000-0002-7515-2143>

**Marine Georgiyants**

Kharkiv Medical Academy of Postgraduate Education,  
 Kharkiv, Ukraine

**ORCID:** <http://orcid.org/0000-0002-1373-7840>

**Olena Vysotska**

Kharkiv National University of Radio Electronics,  
 Kharkiv, Ukraine

**ORCID:** <http://orcid.org/0000-0003-3723-9771>

**Yurii Balym**

Kharkiv State Zooveterinary Academy, Malaya Danylivka,  
 Dergachi district, Kharkiv region, Ukraine

**ORCID:** <http://orcid.org/0000-0002-2494-1329>

In the paper, a descriptive model of system dynamics for binary data is presented. Binary or dichotomous data are widely spread across various fields of research – in decision making and data mining, marketing, solving of many natural, social and technical problems. The initial data for building the model is a set of states of an autonomous dynamical system with components taking binary values. At the same time, the time order of the states is permissible. The following objectives were stated: to identify the relationships between the components of the system defining its dynamics; on the basis of the identified dynamics, to recover the true order of the system states; to apply the developed model to the problem of visual identification of cyanobacteria in water areas using digital photography.

To solve the problem, we used a mathematical model that enables to describe the relationships between components and transitions between the system states at a simple-for-understanding level. The principle of parsimony underlies the model. According to this principle, the most appropriate model is described by the simplest relations in the sense defined in the work.

As the case study, the problem of recognizing clumps of cyanobacteria from digital satellite imagery was considered. This is a complex, practically important problem that does not have a satisfactory experimental and theoretical solution at the moment. Applying system approaches to the measured colorimetric parameters of digital photography, we developed the index for identification of such clumps. This index uses

the parameters of the digital RGB model of (various parts of) an image and allows us to reveal clumps of cyanobacteria on digital images obtained by aerospace methods. Additionally, digital photography can be performed in the conditions of insufficient visibility (due to precipitation, fog, and other factors), for imitation of which in the case study the original image was distorted by the digital noise.

The studied model can find useful applications in the areas requiring binary dynamical data insights.

**Keywords:** descriptive models, dynamical systems, binary data, parsimony, data mining, clumps of toxic cyanobacteria.

### References

- Zholtkevych, G. N., Bespalov, G. Y., Nosov, K. V., Abhishek, M. (2013). Discrete Modeling of Dynamics of Zooplankton Community at the Different Stages of an Antropogeneous Eutrophication. *Acta Biotheoretica*, 61 (4), 449–465. doi: 10.1007/s10441-013-9184-6
- Zholtkevych, G., Nosov, K., Bespalov, Yu., Rak, L., Vysotskaya, E., Balkova, Y., Kolomiychenko, V. (2016). Descriptive Models of System Dynamics. Proceedings of the 12th International Conference on ICT in Education, Research and Industrial Applications. Integration, Harmonization and Knowledge Transfer, 1614, 57–72
- Carmichael, W. W. (2001). Health Effects of Toxin-Producing Cyanobacteria: “The CyanoHABs.” *Human and Ecological Risk Assessment: An International Journal*, 7 (5), 1393–1407. doi: 10.1080/20018091095087
- Zohary, T., Robarts, R. D. (1990). Hyperscums and the population dynamics of *Microcystis aeruginosa*. *Journal of Plankton Research*, 12 (2), 423–432. doi: 10.1093/plankt/12.2.423
- Karabashev, G. S., Evdoshenko, M. A. (2015). Spektral'nye priznaki cveteniya cianobakterij v Baltijskom more po dannym skanera MODIS. *Sovremennye problemy distancionnogo zondirovaniya Zemli iz kosmosa*, 12 (3), 158–170.
- Vysotska, O. V., Bespalov, Yu. G., Pecherska, A. I., Parvadov, D. A. (2016). Using of margalef succession model in remote detection technologies for indications of human impact on vegetation cover. *Radioelektronni i komp'uterni sistemi*, 2 (76), 15–19.
- Vysotskaya, E. V., Zholtkevych, G. N., Klochko, T. A., Bespalov, Yu. G., Nosov, K. V. (2016). Unmasking the soil cover's disruption by modeling the dynamics of ground vegetation parameters. *Visnyk Natsionalnoho Tekhnichnoho Universytetu Ukrayiny «KPI»*. Seriya – Radiotekhnika. Radioaparaturbuduvannya, 64, 101–109.
- Shulika, B., Porvan, A., Vysotska, O., Nekos, A., Zhemerov, A. (2017). Control over grape yield in the north-eastern region of Ukraine using mathematical modeling. *Eastern-European Journal of Enterprise Technologies*, 2 (3 (86)), 51–59. doi: 10.15587/1729-4061.2017.97969
- De Menezes, F. S., Liska, G. R., Cirillo, M. A., Vivanco, M. J. F. (2017). Data classification with binary response through the Boosting algorithm and logistic

- regression. *Expert Systems with Applications*, 69, 62–73. doi: 10.1016/j.eswa.2016.08.014
10. Pierola, A., Epifanio, I., Alemany, S. (2016). An ensemble of ordered logistic regression and random forest for child garment size matching. *Computers & Industrial Engineering*, 101, 455–465. doi: 10.1016/j.cie.2016.10.013
  11. Ghattas, B., Michel, P., Boyer, L. (2017). Clustering nominal data using unsupervised binary decision trees: Comparisons with the state of the art methods. *Pattern Recognition*, 67, 177–185. doi: 10.1016/j.patcog.2017.01.031
  12. Yamamoto, M., Hayashi, K. (2015). Clustering of multivariate binary data with dimension reduction via L1-regularized likelihood maximization. *Pattern Recognition*, 48 (12), 3959–3968. doi: 10.1016/j.patcog.2015.05.026
  13. Katahira, K. (2016). How hierarchical models improve point estimates of model parameters at the individual level. *Journal of Mathematical Psychology*, 73, 37–58. doi: 10.1016/j.jmp.2016.03.007
  14. Erkan Ozkaya, H., Dabas, C., Kolev, K., Hult, G. T. M., Dahlquist, S. H., Manjeshwar, S. A. (2013). An assessment of hierarchical linear modeling in international business, management, and marketing. *International Business Review*, 22 (4), 663–677. doi: 10.1016/j.ibusrev.2012.10.002
  15. Van Oirbeek, R., Lesaffre, E. (2012). Assessing the predictive ability of a multilevel binary regression model. *Computational Statistics & Data Analysis*, 56 (6), 1966–1980. doi: 10.1016/j.csda.2011.11.023
  16. Asar, Ö., İlk, Ö. (2014). Flexible multivariate marginal models for analyzing multivariate longitudinal data, with applications in R. *Computer Methods and Programs in Biomedicine*, 115 (3), 135–146. doi: 10.1016/j.cmpb.2014.04.005
  17. Zou, G., Donner, A. (2011). Extension of the modified Poisson regression model to prospective studies with correlated binary data. *Statistical Methods in Medical Research*, 22 (6), 661–670. doi: 10.1177/0962280211427759
  18. Schildcrout, J. S., Heagerty, P. J. (2006). Marginalized Models for Moderate to Long Series of Longitudinal Binary Response Data. *Biometrics*, 63 (2), 322–331. doi: 10.1111/j.1541-0420.2006.00680.x
  19. Kupek, E. (2006). Beyond logistic regression: structural equations modelling for binary variables and its application to investigating unobserved confounders. *BMC Medical Research Methodology*, 6 (1). doi: 10.1186/1471-2288-6-13
  20. Rhemtulla, M., Brosseau-Liard, P. É., Savalei, V. (2012). When can categorical variables be treated as continuous? A comparison of robust continuous and categorical SEM estimation methods under suboptimal conditions. *Psychological Methods*, 17 (3), 354–373. doi: 10.1037/a0029315
  21. Pearl, J. (2009). Causal inference in statistics: An overview. *Statistics Surveys*, 3, 96–146. doi: 10.1214/09-ss057
  22. Bollen, K. A., Pearl, J. (2013). Eight Myths About Causality and Structural Equation Models. *Handbook of Causal Analysis for Social Research*, 301–328. doi: 10.1007/978-94-007-6094-3\_15
  23. LANCE-MODIS Collection 6. Available at: <https://lance3.modaps.eosdis.nasa.gov>

---

**DOI: 10.15587/1729-4061.2017.107536**  
**MODELS AND METHODS OF REGRESSION**  
**ANALYSIS UNDER CONDITIONS OF FUZZY**  
**INITIAL DATA (p. 12-19)**

**Lev Raskin**

National Technical University “Kharkiv Polytechnic Institute”, Kharkiv, Ukraine  
**ORCID:** <http://orcid.org/0000-0002-9015-4016>

**Oksana Sira**

National Technical University “Kharkiv Polytechnic Institute”, Kharkiv, Ukraine  
**ORCID:** <http://orcid.org/0000-0002-4869-2371>

**Yuriy Ivanchykhin**

National Technical University “Kharkiv Polytechnic Institute”, Kharkiv, Ukraine  
**ORCID:** <http://orcid.org/0000-0002-3999-6541>

The paper considers the problem of regression analysis with indeterminate explanatory and explained variables. A quality criterion for estimating the regression coefficients is formulated and justified, taking into account possible significant differences in the accuracy of assigning different variables. The study considers a method of calculating the regression coefficients in accordance with the concept of least squares. The proposed approach provides a reasonable compromise between the conflicting requirements: the maximum compactness of the fuzzy value function of the explained variable and the minimal deviation of the solution from the modal one. The problem is solved by minimizing the complex criterion, the terms of which determine the level of satisfaction of these requirements. An additional advantage of the approach is that the original problem, fuzzy by the nature of the initial data, is reduced to solving two usual problems of mathematical programming. The problem of fuzzy comparator identification is considered when the values of the explained variable are not defined but can be ranked by the descending of any chosen indicator. To solve this problem, the study proposes a method for estimating regression coefficients based on solving a fuzzy system of linear algebraic equations.

**Keywords:** fuzzy regression analysis, fuzzy initial data, fuzzy comparator identification.

### References

1. Linear Regression Analysis with Fuzzy Model (1982). *IEEE Transactions on Systems, Man, and Cybernetics*, 12 (6), 903–907. doi: 10.1109/tsmc.1982.4308925
2. Yang, M.-S., Liu, H.-H. (2003). Fuzzy least-squares algorithms for interactive fuzzy linear regression models. *Fuzzy Sets and Systems*, 135 (2), 305–316. doi: 10.1016/s0165-0114(02)00123-9
3. Chen, F., Chen, Y., Zhou, J., Liu, Y. (2016). Optimizing h value for fuzzy linear regression with asymmetric triangular fuzzy coefficients. *Engineering Applications of Artificial Intelligence*, 47, 16–24. doi: 10.1016/j.engappai.2015.02.011
4. Ishibuchi, H., Nii, M. (2001). Fuzzy regression using asymmetric fuzzy coefficients and fuzzified neural networks. *Fuzzy Sets and Systems*, 119 (2), 273–290. doi: 10.1016/s0165-0114(98)00370-4
5. Chang, Y.-H. O., Ayyub, B. M. (2001). Fuzzy regression methods – a comparative assessment. *Fuzzy Sets*

- and Systems, 119 (2), 187–203. doi: 10.1016/s0165-0114(99)00091-3
6. Yen, K. K., Ghoshray, S., Roig, G. (1999). A linear regression model using triangular fuzzy number coefficients. *Fuzzy Sets and Systems*, 106 (2), 167–177. doi: 10.1016/s0165-0114(97)00269-8
  7. Charfeddine, S., Mora-Camino, F. A. C., De Coligny, M. (2004). Fuzzy linear regression: application to the estimation of air transport demand. *FSSCEF 2004, International Conference on Fuzzy Sets and Soft Computing in Economics and Finance*, 350–359. Available at: <https://hal-enac.archives-ouvertes.fr/hal-01022443>
  8. Zaychenko, Yu. P. (2007). Nechetkiy metod gruppovogo ucheta argumentov pri neopredelennykh iskhodnykh dannykh. *Systemni doslidzhennia ta informatsiini tekhnologii*, 3, 100–112.
  9. Zgurovskiy, M., Zaychenko, Yu. (2013). *Modeli i metody prinyatiya resheniy v nechetkiykh usloviyah*. Kyiv: Naukova dumka, 275.
  10. Celmins, A. (1987). Least squares model fitting to fuzzy vector data. *Fuzzy Sets and Systems*, 22 (3), 245–269. doi: 10.1016/0165-0114(87)90070-4
  11. Muzzioli, S., Ruggieri, A., De Baets, B. (2015). A comparison of fuzzy regression methods for the estimation of the implied volatility smile function. *Fuzzy Sets and Systems*, 266, 131–143. doi: 10.1016/j.fss.2014.11.015
  12. Diamond, P. (1988). Fuzzy least squares. *Information Sciences*, 46 (3), 141–157. doi: 10.1016/0020-0255(88)90047-3
  13. Chang, Y.-H. O. (2001). Hybrid fuzzy least-squares regression analysis and its reliability measures. *Fuzzy Sets and Systems*, 119 (2), 225–246. doi: 10.1016/s0165-0114(99)00092-5
  14. Yang, M.-S., Lin, T.-S. (2002). Fuzzy least-squares linear regression analysis for fuzzy input–output data. *Fuzzy Sets and Systems*, 126 (3), 389–399. doi: 10.1016/s0165-0114(01)00066-5
  15. Kao, C., Chyu, C.-L. (2002). A fuzzy linear regression model with better explanatory power. *Fuzzy Sets and Systems*, 126 (3), 401–409. doi: 10.1016/s0165-0114(01)00069-0
  16. Shtovba, S. D. (2016). Nechetkaya identifikatsiya na osnove regressiynykh modeley parametricheskoy funktsii prinalozhnosti. *Problemy upravleniya i informatiki*, 6, 1–8.
  17. Chachi, J., Taheri, S. M. (2016). Multiple fuzzy regression model for fuzzy input-output data. *Iranian Journal of Fuzzy Systems*, 13 (4), 63–78.
  18. Zak, Yu. (2017). Fuzzy-regressiionnye modeli v usloviyah nalichiya v statisticheskoy vyborke nechislovy informatsii. *System Research & Information Technologies*, 1, 88–96.
  19. Jung, H.-Y., Yoon, J. H., Choi, S. H. (2015). Fuzzy linear regression using rank transform method. *Fuzzy Sets and Systems*, 274, 97–108. doi: 10.1016/j.fss.2014.11.004
  20. Pushpa, B., Vasuki, R. (2013). A Least Absolute Approach to Multiple Fuzzy Regression Using Tw- Norm Based Operations. *International Journal of Fuzzy Logic Systems*, 3 (2), 73–84. doi: 10.5121/ijfls.2013.3206
  21. Namdari, M., Yoon, J. H., Abadi, A., Taheri, S. M., Choi, S. H. (2014). Fuzzy logistic regression with least absolute deviations estimators. *Soft Computing*, 19 (4), 909–917. doi: 10.1007/s00500-014-1418-2
  22. Ubale, A. B., Sananse, S. L. (2015). Fuzzy Regression Model and Its Application: A Review. *International Journal of Innovative Research in Science, Engineering and Technology*, 4 (11), 10853–10860.
  23. Kryuchkovskiy, V. V., Petrov, Eh. G., Sokolova, N. A., Hodakov, V. E. (2013). *Vedenie v normativnuyu teoriyu prinyatiya resheniy*. Herson: Grin' D. S., 284.
  24. Zuhovickiy, S. I., Avdeeva, L. I. (1967). *Lineynoe i vypukloe programmirovaniye*. Moscow: Nauka, 460.
  25. Sira, O. V., Al-Shqeerat, K. H. (2009). A New Approach for Resolving Equations with Fuzzy Parameters. *European Journal of Scientific Research*, 38 (4), 619–625.
  26. Seraya, O. V., Demin, D. A. (2012). Linear Regression Analysis of a Small Sample of Fuzzy Input Data. *Journal of Automation and Information Sciences*, 44 (7), 34–48. doi: 10.1615/jautomatinfscien.v44.i7.40
  27. Raskin, L. G., Seraya, O. V. (2008). *Nechetkaya matematika*. Kharkiv: Parus, 352.
  28. Raskin, L., Sira, O. (2016). Method of solving fuzzy problems of mathematical programming. *Eastern-European Journal of Enterprise Technologies*, 5 (4 (83)), 23–28. doi: 10.15587/1729-4061.2016.81292
  29. Pawlak, Z. (1982). Rough sets. *International Journal of Computer & Information Sciences*, 11 (5), 341–356. doi: 10.1007/bf01001956
  30. Raskin, L., Sira, O. (2016). Fuzzy models of rough mathematics. *Eastern-European Journal of Enterprise Technologies*, 6 (4 (84)), 53–60. doi: 10.15587/1729-4061.2016.86739
  31. Raskin, L. G., Kirichenko, I. O., Seraya, O. V. (2013). *Prikladnoe kontinual'noe lineynoe programmirovaniye*. Kharkiv, 293.

DOI: 10.15587/1729-4061.2017.108327

#### CONSTRUCTION OF INTERPOLATION METHOD FOR NUMERICAL SOLUTION OF THE CAUCHY'S PROBLEM (p. 19-27)

Roman Bihun

Ivan Franko National University of Lviv, Lviv, Ukraine

ORCID: <http://orcid.org/0000-0003-4363-4532>

Gregoriy Tsehelyk

Ivan Franko National University of Lviv, Lviv, Ukraine

ORCID: <http://orcid.org/0000-0002-5826-0628>

An interpolation numerical method is developed in order to solve the Cauchy's problem for ordinary first order differential equations using the apparatus of non-classical minorants and diagrams of Newton's functions, assigned in a tabular form. We have proven computational stability of the method, that is, an error of the initial data is not piled up. It is also shown that the method possesses a second order of accuracy and in the case of a convex function produces more accurate results than the Euler's method. The advantages also include simplicity and visual clarity of the method. Given this, it could gain widespread use in many areas, in particular mathematics, physics and mechanics. We also give an example of solving the Cauchy's problem applying the new method, the Euler's method, and the Runge-Kutta fourth order method, with the results compared. The proposed method does not require solving the systems of linear algebraic equations because we do not employ the Bernstein polynomials, and it is not required to superimpose additional conditions, in contrast to the method that applies the Haar functions.

**Keywords:** Newton's minorant, differential equations, Cauchy's problem, Newton's diagram, convex function.

### References

1. Verzhbitskiy, V. M. (2001). Chislennyye metody (matematicheskiy analiz i obyknovennyye differentsialnyye uravneniya). Moscow: Vyschaya shkola, 382.
2. Süli, E., Mayers, D. (2003). An Introduction to Numerical Analysis. Cambridge University Press, 435.
3. Zadachyn, V. M. (2014). Chyselni metody. Kharkiv: Vyd. KhNEU im. S. Kuznetsya, 180.
4. Abdulkawi, M. (2015). Solution of Cauchy type singular integral equations of the first kind by using differential transform method. Applied Mathematical Modelling, 39 (8), 2107–2118. doi: 10.1016/j.apm.2014.10.003
5. Setia, A. (2014). Numerical solution of various cases of Cauchy type singular integral equation. Applied Mathematics and Computation, 230, 200–207. doi: 10.1016/j.amc.2013.12.114
6. De Bonis, M. C., Laurita, C. (2012). Numerical solution of systems of Cauchy singular integral equations with constant coefficients. Applied Mathematics and Computation, 219 (4), 1391–1410. doi: 10.1016/j.amc.2012.08.022
7. Lukomskii, D. S., Lukomskii, S. F., Terekhin, P. A. (2016). Solution of Cauchy Problem for Equation First Order Via Haar Functions. Izvestiya of Saratov University. New Series. Series: Mathematics. Mechanics. Informatics, 16 (2), 151–159. doi: 10.18500/1816-9791-2016-16-2-151-159
8. Lyashenko, M. Ya., Holovan, M. S. (1996). Chysel'ni metody. Kyiv: "Lybid", 285.
9. Varenych, I. I. (2008). Vyshcha matematyka: matematychnyy analiz, dyferentsialni rivnyannya. Kyiv: Dia-Soft, 267.
10. Bihun, R., Tsehelyk, G. (2017). Construction of a numerical method for finding the zeros of both smooth and nonsmooth functions. Eastern-European Journal of Enterprise Technologies, 2 (4 (86)), 58–64. doi: 10.15587/1729-4061.2017.99273
11. Bihun, R. R., Tsehelyk, G. G. (2017). Formulas of minorant type to approximate computing definite integrals. Scientific light, 1 (7), 76–79.
12. Bihun, R. R., Tsehelyk, G. G. (2015). Numerical Method for Finding All Points of Extremum of Random as Smooth and NonSmooth Functions of One Variable. Global Journal of Science Frontier Research: F Mathematics & Decision Sciences, 15 (2), 87–93.

**DOI: 10.15587/1729-4061.2017.108413**  
**DEVELOPMENT OF THE APPROACH TO**  
**PROVING THE SECURITY OF BLOCK CIPHERS**  
**TO IMPOSSIBLE DIFFERENTIAL ATTACK**  
**(p. 28-33)**

**Victor Ruzhentsev**

Kharkiv National University of Radio Electronics,  
 Kharkiv, Ukraine

**ORCID:** <http://orcid.org/0000-0002-1007-6530>

**Yurii Onishchenko**

Kharkiv National University of Internal Affairs,  
 Kharkiv, Ukraine

**ORCID:** <http://orcid.org/0000-0002-7755-3071>

Trends in the development of symmetric cryptography are constantly associated with the increasing of the sizes of keys and blocks. Block ciphers, which are used today in hashing algorithms, usually have a block size of 512 or 1024 bits. One of the main requirements for symmetric crypto algorithms is to provide resistance to known cryptanalytical attacks. Known methods of security estimation against impossible differential attack have too high complexity for such block sizes.

The proposed approach for proving the absence of impossible differentials is applicable to some types of block ciphers and allows proving theoretically the resistance to impossible differentials attack.

Rijndael-like SPN ciphers and Feistel ciphers are analyzed. For the group of Rijndael-like ciphers, the absence of byte impossible differentials for 4 or more rounds is proved. For the group of Feistel ciphers, the absence of byte impossible differentials for 6 or more rounds is proved. The first statement made it possible to prove the absence of byte impossible differentials for 4 or more rounds of the cipher Kalyna (DSTU 7624: 2014) with all block sizes, for 512-bit block ciphers that are used in the hash functions Whirlpool, Groestl and Kupyna (DSTU 7564: 2014). The second statement was used to prove the absence of byte impossible differentials for 6 or more rounds of Tornado and Labyrinth ciphers with a block size of 128 bits.

Computational experiments on the impossible differentials search for these reduced models confirmed the validity of the obtained theoretical conclusions.

**Keywords:** block cipher, impossible differentials attack, impossible differential, Rijndael-like transformations.

### References

1. Bihun, E., Keller, N. (2000). Cryptanalysis of Reduced Variant of Rijndael. The Third Advanced Encryption Standard Candidate Conference. New York.
2. Cheon, J. H., Kim, M., Kim, K., Jung-Yeun, L., Kang, S. (2002). Improved Impossible Differential Cryptanalysis of Rijndael and Crypton. Lecture Notes in Computer Science, 39–49. doi: 10.1007/3-540-45861-1\_4
3. Lu, J., Dunkelman, O., Keller, N., Kim, J. (2008). New Impossible Differential Attacks on AES. Lecture Notes in Computer Science, 279–293. doi: 10.1007/978-3-540-89754-5\_22
4. Bihun, E., Biryukov, A., Shamir, A. (1998). Cryptanalysis of Skipjack Reduced to 31 Rounds using Impossible Differentials. Technion, CS Dept, Tech Report CS0947.
5. Bihun, E., Biryukov, A., Shamir, A. (1999). Miss in the Middle Attacks on IDEA and Khufu. Lecture Notes in Computer Science, 124–138. doi: 10.1007/3-540-48519-8\_10
6. J Lu, J., Kim, J., Keller, N., Dunkelman, O. (2008). Improving the Efficiency of Impossible Differential Cryptanalysis of Reduced Camellia and MISTY1. Lecture Notes in Computer Science, 370–386. doi: 10.1007/978-3-540-79263-5\_24
7. Wu, W.-L., Zhang, W.-T., Feng, D.-G. (2007). Impossible Differential Cryptanalysis of Reduced-Round ARIA and Camellia. Journal of Computer Science and Technology, 22 (3), 449–456. doi: 10.1007/s11390-007-9056-0
8. Kim, J., Hong, S., Sung, J., Lee, S., Lim, J., Sung, S. (2003). Impossible Differential Cryptanalysis for Block Cipher Structures. Lecture Notes in Computer Science, 82–96. doi: 10.1007/978-3-540-24582-7\_6

9. Luo, Y., Wu, Z., Lai, X., Gong, G. (2009). A Unified Method for Finding Impossible Differentials of Block Cipher Structures. IACR Cryptology ePrint Archive.
10. Li, R., Sun, B., Li, C. (2010). Impossible Differential Cryptanalysis of SPN Ciphers. IACR Cryptology ePrint Archive.
11. Yap, H. (2009). Impossible Differential Characteristics of Extended Feistel Networks with Provable Security against Differential Cryptanalysis. *Advances in Security Technology*, 103–121. doi: 10.1007/978-3-642-10240-0\_9
12. Daemen, J., Rijmen, V. (1998). AES proposal: Rijndael. First Advanced Encryption Standard (AES) Conference. Ventura, CA.
13. Gorbenko, I. D., Dolgov, V. I., Oliynykov, R. V., Ruzhentsev, V. I. et. al. (2007). The perspective block symmetrical cipher “Kalyna” – a main positions and specification. *Applied radioelektroniks*, 6 (2), 195–208.
14. V Ruzhentsev, V. I. (2003). About method of the security estimation to truncated differential attack. *Radioelektroniks and informatics*, 4, 130–133.
15. Gorbenko, I. D., Golovashich, S. A. (2003). The Algorithm of the block symmetrical enciphering “Tornado”. The specification of the transformation. *Radiotechnics*, 134, 60–80.
16. Golovashich, S. A. (2007). Specification of the algorithm of the block symmetrical enciphering “Labyrinth”. *Applied Radioelektroniks*, 6 (2), 230–240.

DOI: 10.15587/1729-4061.2017.108395

**FEATURES OF MODELING FAILURES OF RECOVERABLE COMPLEX TECHNICAL OBJECTS WITH A HIERARCHICAL CONSTRUCTIVE STRUCTURE (p. 34-42)**

**Sergey Lenkov**

Military Institute of Taras Shevchenko National University of Kyiv, Kyiv, Ukraine

ORCID: <http://orcid.org/0000-0001-7689-239X>

**Genadiy Zhyrov**

Military Institute of Taras Shevchenko National University of Kyiv, Kyiv, Ukraine

ORCID: <http://orcid.org/0000-0001-7648-7992>

**Dmytro Zaitsev**

Military Institute of Taras Shevchenko National University of Kyiv, Kyiv, Ukraine

ORCID: <http://orcid.org/0000-0002-3784-5790>

**Igor Tolok**

Military Institute of Taras Shevchenko National University of Kyiv, Kyiv, Ukraine

ORCID: <http://orcid.org/0000-0001-6309-9608>

**Evgen Lenkov**

Military Institute of telecommunications and Informatization, Kyiv, Ukraine

ORCID: <http://orcid.org/0000-0001-5819-2656>

**Tetiana Bondarenko**

Military Institute of telecommunications and Informatization, Kyiv, Ukraine

ORCID: <http://orcid.org/0000-0002-2879-2041>

**Yurii Gunchenko**

Odessa I. I. Mechnikov National University, Odessa, Ukraine

ORCID: <http://orcid.org/0000-0003-4423-8267>

**Viktor Zagrebnyuk**

Odessa National Maritime University, Odessa, Ukraine

ORCID: <http://orcid.org/0000-0002-8630-1976>

**Oleksandr Antonenko**

Odessa I. I. Mechnikov National University, Odessa, Ukraine

ORCID: <http://orcid.org/0000-0001-9680-3446>

We developed a methodology and algorithms of forming the optimal sets of elements, which take part in the failure-recovery modeling process of complex technical objects. The methodology is based on the hierarchical constructive structure of the object, takes into account the redundancy of failing elements, as well as the maintainability of the product elements and their cost, which distinguishes this methodology from the known ones. Structurally, the methodology is implemented as a set of three algorithms. The generated optimal sets of elements are used to calculate the predictive reliability characteristics and operating costs of the object. The constructive structure of the object in the model is represented by a graph (tree). The optimality of the sets is understood in the sense of their correspondence to the object maintainability parameters.

With the improvement of maintainability properties, the forecasted values of the mean time between failures and the recovery time are correspondingly improved. Improvement of the operating cost index is not mandatory; provided different input data, there may not be such an improvement. Each variant of the values of object maintainability parameters is conformed to its optimal sets of failing and recoverable elements, under which adequate predictive estimates of reliability indicators and the object operating cost are provided.

The paper provides some examples of modeling, which demonstrate how the optimal sets of failing and recoverable elements are determined, and how the predictive estimates of reliability and object operating cost depend on the choice of these sets.

**Keywords:** complex technical object, hierarchical constructive structure, mean time between failures, unit operating costs, statistical simulation modeling.

## References

1. Lenkov, S. V., Boriak, K. F., Banzak, G. V., Braun, V. O., Osypa, V. A., Pashkov, S. A., Tsytsarev, V. N., Berezovskaia, Iu. V. (2014). Forecasting to reliability complex object radio-electronic technology and optimization parameter their technical usage with use the simulation statistical models. Odessa: VMV, 252.
2. Shannon, R. E. (1975). *Systems Simulation: The Art and Science*. Prentice Hall, 368.
3. Brown, J., Mol, L. (2017). On the roots of all-terminal reliability polynomials. *Discrete Mathematics*, 340 (6), 1287–1299. doi: 10.1016/j.disc.2017.01.024
4. Liang, X. F., Wang, H. D., Yi, H., Li, D. (2017). Warship reliability evaluation based on dynamic bayesian networks and numerical simulation. *Ocean Engineering*, 136, 129–140. doi: 10.1016/j.oceaneng.2017.03.023
5. Tu, H., Lou, W., Sun, Z., Qian, Y. (2017). Structural reliability simulation for the latching mechanism in

- MEMS-based Safety and Arming device. *Advances in Engineering Software*, 108, 48–56. doi: 10.1016/j.advengsoft.2017.02.008
6. Wu, J., Yan, S., Li, J., Gu, Y. (2016). Mechanism reliability of bistable compliant mechanisms considering degradation and uncertainties: Modeling and evaluation method. *Applied Mathematical Modelling*, 40 (23-24), 10377–10388. doi: 10.1016/j.apm.2016.07.006
  7. Okaro, I. A., Tao, L. (2016). Reliability analysis and optimisation of subsea compression system facing operational covariate stresses. *Reliability Engineering & System Safety*, 156, 159–174. doi: 10.1016/j.res.2016.07.018
  8. Cui, L., Li, Y., Shen, J., Lin, C. (2016). Reliability for discrete state systems with cyclic missions periods. *Applied Mathematical Modelling*, 40 (23-24), 10783–10799. doi: 10.1016/j.apm.2016.08.004
  9. Tien, I., Der Kiureghian, A. (2016). Algorithms for Bayesian network modeling and reliability assessment of infrastructure systems. *Reliability Engineering & System Safety*, 156, 134–147. doi: 10.1016/j.res.2016.07.022
  10. Li, Y. Y., Chen, Y., Yuan, Z. H., Tang, N., Kang, R. (2017). Reliability analysis of multi-state systems subject to failure mechanism dependence based on a combination method. *Reliability Engineering & System Safety*, 166, 109–123. doi: 10.1016/j.res.2016.11.007
  11. Li, M., Hu, Q., Liu, J. (2013). Proportional hazard modeling for hierarchical systems with multi-level information aggregation. *IIE Transactions*, 46 (2), 149–163. doi: 10.1080/0740817x.2013.772692
  12. Li, M., Liu, J., Li, J., Uk Kim, B. (2014). Bayesian modeling of multi-state hierarchical systems with multi-level information aggregation. *Reliability Engineering & System Safety*, 124, 158–164. doi: 10.1016/j.res.2013.12.001
  13. Grishko, A., Yurkov, N., Goryachev, N. (2017). Reliability analysis of complex systems based on the probability dynamics of subsystem failures and deviation of parameters. 2017 14th International Conference The Experience of Designing and Application of CAD Systems in Microelectronics (CADSM), 79–82. doi: 10.1109/cadsm.2017.7916109
  14. For, R., Kofman, A., Deni-Papen, M. (1966). *Sovremennaya matematika*. Moscow: Mir, 272.
  15. Strelnikov, V. P., Feduhin, A. V. (2002). *Otsenka i prognozirovanie nadezhnosti elektronnykh elementov i sistem*. Kyiv: Logos, 486.

**DOI: 10.15587/1729-4061.2017.108387**  
**CRITERION OF THE CONTINUATION OF HARMONIC FUNCTIONS IN THE BALL OF n-DIMENSIONAL SPACE AND REPRESENTATION OF THE GENERALIZED ORDERS OF THE ENTIRE HARMONIC FUNCTIONS IN  $\mathbb{R}^n$  IN TERMS OF APPROXIMATION ERROR (p. 43-49)**

**Olga Veselovska**

Lviv Polytechnic National University, Lviv, Ukraine  
**ORCID:** <http://orcid.org/0000-0003-4052-1081>

**Khrystyna Drohomyretska**

Lviv Polytechnic National University, Lviv, Ukraine  
**ORCID:** <http://orcid.org/0000-0002-4987-7810>

**Lubov Kolyasa**

Lviv Polytechnic National University, Lviv, Ukraine  
**ORCID:** <http://orcid.org/0000-0002-9690-8042>

A growth of harmonic functions in the whole space  $\mathbb{R}^n$  is examined. We found the estimate for a uniform norm of spherical harmonics in terms of the best approximation of harmonic function in the ball by harmonic polynomials. An approximation error of harmonic function in the ball is estimated by the maximum modulus of an entire harmonic function in space, as well as the maximum modulus of an entire harmonic function in space in terms of the maximum modulus of some entire function of one complex variable or the maximal term of its power series. These results allowed us to obtain the necessary and sufficient conditions under which a harmonic function in the ball of an  $n$ -dimensional space,  $n \geq 3$ , can be continued to the entire harmonic one. This result is formulated in terms of the best approximation of the given function by harmonic polynomials. In order to characterize growth of an entire harmonic function, we used the generalized and the lower generalized orders. Formulae for the generalized and the lower generalized orders of an entire harmonic function in space are expressed in terms of the approximation error by harmonic polynomials of the function that continues. We also investigated the growth of functions of slow increase. The obtained results are analogues to classical results, which are known for the entire functions of one complex variable.

The conducted research is important due to the fact that the harmonic functions occupy a special place not only in many mathematical studies, but also when applying mathematical analysis to physics and mechanics, where these functions are often employed to describe various stationary processes.

**Keywords:** spherical harmonics, entire harmonic function, generalized order, lower generalized order.

## References

1. Srivastava, G. S., Kumar, S. (2009). Uniform approximation of entire functions on compact sets and their generalized growth. *New Zealand Journal of Mathematics*, 39, 33–43.
2. Srivastava, G. S., Kumar, S. (2009). Approximation of entire functions of slow growth on compact sets. *Archivum Mathematicum (BRNO)*, 45, 137–146.
3. Harfaoui, M. (2010). Generalized Order and Best Approximation of Entire Function in  $p$ -Norm. *International Journal of Mathematics and Mathematical Sciences*, 2010, 1–15. doi: 10.1155/2010/360180
4. Kumar, D. (2011). Generalized growth and best approximation of entire functions in  $L^p$ -norm in several complex variables. *Annali dell'universita' di ferrara*, 57 (2), 353–372. doi: 10.1007/s11565-011-0130-8
5. Datta, S. K., Biswas, T. (2016). Growth analysis of entire functions of two complex variables. *Sahand Communications in Mathematical Analysis*, 3 (2), 13–24.
6. Srivastava, G. S., Kumar, S. (2011). On approximation and generalized type of analytic functions of several complex variables. *Analysis in Theory and Applications*, 27 (2), 101–108. doi: 10.1007/s10496-011-0101-z
7. Harfaoui, M., Kumar, D. (2014). Best Approximation in  $L^p$ -norm and Generalized  $(\alpha, \beta)$ -growth of Analytic Functions. *Theory and Applications of Mathematics & Computer Science*, 4 (1), 65–80.
8. Temliakov, A. A. (1935). K probleme rosta harmonycheskykh funktsiyi trekhmernoho prostranstva. *Matematicheskii sbornik*, 42 (6), 707–718.
9. Kapoor, G. P., Nautiyal, A. (1983). On the growth of harmonic functions in  $\mathbb{R}^3$ . *Demonstr. math.*, 16 (4), 811–819.

10. Veselovskaia, O. V. (1983). O roste tselykh harmonycheskykh v funktsyi. *Yzv. Vuzov. Matem.*, 10, 13–17.
11. Srivastava, G. S. (2008). Generalized growth of entire harmonic functions. *Fasciculi Mathematici*, 40, 79–89.
12. Fugard, T. B. (1980). Growth of entire harmonic functions in  $\mathbb{R}^n$ ,  $n \geq 2$ . *Journal of Mathematical Analysis and Applications*, 74 (1), 286–291. doi: 10.1016/0022-247x(80)90127-4
13. Srivastava, G. S. (2008). Growth of entire harmonic functions in  $\mathbb{R}^n$ , and generalized orders. *Bulletin of the Greek Mathematical Society*, 55, 49–58.
14. Kumar, D., Gupta, S. K. (2012). Growth of universal entire harmonic functions. *TJMM*, 3 (2), 111–116.
15. Kumar, D., Arora, K. N. (2010). Growth and approximation properties of generalized axisymmetric potentials. *Demonstratio Mathematica*, 43 (1), 107–116. doi: 10.1515/dema-2013-0215
16. Kumar, D. (2010) Growth and Chebyshev Approximation of Entire Function Solutions of Helmholtz Equation in  $\mathbb{R}^2$ . *European Journal of Pure and Applied Mathematics*, 3 (6), 1062–1069.
17. Kumar, D., Basu, A. (2015). Growth and Approximation of Generalized Bi-Axially Symmetric Potentials. *Journal of Mathematical Research with Applications*, 35 (6), 613–624.
18. Khan, H. H., Ali, R. (2012). Slow Growth and Approximation of Entire Solution of Generalized Axially Symmetric Helmholtz Equation. *Asian Journal of Mathematics & Statistics*, 5 (4), 104–120. doi: 10.3923/ajms.2012.104.120
19. Kumar, D. (2011) On the  $(p, q)$ -growth of entire function solution of Helmholtz equation. *J. Nonlinear Sci. Appl.*, 4 (2), 92–101.
20. Kumar, D. (2014). Growth and approximation of solutions to a class of certain linear partial differential equations in  $\mathbb{R}^N$ . *Mathematica Slovaca*, 64 (1), 139–154. doi: 10.2478/s12175-013-0192-4
21. Kumar, D. (2013) Slow Growth and Optimal Approximation of Pseudoanalytic Functions on the Disk. *International Journal of Analysis and Applications*, 2 (1), 26–37.
22. Kapoor, G. P., Nautiyal, A. (1982). Approximation of entire harmonic functions in  $\mathbb{R}^3$ . *Indian Journal of Pure and Applied Mathematics*, 13 (9), 1024–1030.
23. Shaker Abdu-Hussein, M., Srivastava, G. S. (2001). On the generalized type and approximation of entire harmonic functions in  $\mathbb{R}^n$  having index pair  $(p, q)$ . *Istanbul Univ. Fem. Fak. Mat. Der.*, 60, 1–17.
24. Šeremeta, M. N. (1970). On the connection between the growth of the maximum modulus of an entire function and the moduli of the coefficients of its power series expansion. *American Mathematical Society Translations: Series 2*, 291–301. doi: 10.1090/trans2/088/11
25. Stein, Y., Veis, H. (1974). *Vvedeniye v harmonycheskyi analiz na evklydovykh prostranstvakh*. Moscow: Myr, 336.
26. Tyman, A. F., Trofymov, V. N. (1968). *Vvedeniye v teoriyu harmonycheskykh funktsii*. Moscow: Nauka, 208.
27. Beitmen, H., Erdeiy, A. (1974). *Vysshye transtsendentnye funktsyy*. 2nd edition. Moscow: Nauka, 296.
28. Sheremeta, M. N. (1968). O svyazy mezhdu rostom tselykh ili analytycheskykh v kruhe funktsyi nulevoho poriadka i koeffytsyentamy ikh stepennykh razlozheniy. *Yzv. vuzov. Matem.*, 6, 115–121.

**DOI: 10.15587/1729-4061.2017.108404**  
**SYNTHESIS OF GENERALIZED NEURAL ELEMENTS BY MEANS OF THE TOLERANCE MATRICES (p. 50-62)**

**Fedir Geche**

Uzhgorod National University, Uzhhorod, Ukraine  
**ORCID:** <http://orcid.org/0000-0002-4757-9828>

**Oksana Mulesa**

Uzhgorod National University, Uzhhorod, Ukraine  
**ORCID:** <http://orcid.org/0000-0002-6117-5846>

**Viktor Buchok**

Uzhgorod National University, Uzhhorod, Ukraine  
**ORCID:** <http://orcid.org/0000-0002-1992-7290>

Application of neuromorphic structures in various spheres of human activity on the basis of generalized neural elements will become possible if effective methods for verifying realizability of the logic algebra functions by one neuron element with a generalized threshold activation function and synthesis of such elements with a large number of entries are developed. A notion of nucleus of Boolean functions in relation to a given system of characters was introduced and algebraic structure of nuclei and reduced nuclei of Boolean functions was investigated. Relation between the nuclei of the logic algebra functions which are realized by one generalized neural element and matrices of tolerance was established. It was shown that the Boolean function is realized by one generalized neuron element if and only if the nucleus of this function admits representation by the matrices of tolerance. If there is no nucleus relative to a specified system of characters for a Boolean function, then such a function is not realized by one generalized neural element in relation to a specified system of characters. On the basis of the properties of the matrices of tolerance, a number of necessary and sufficient conditions for realization of the logic algebra functions by one generalized neural element were obtained. Based on the sufficient conditions, an algorithm for synthesis of integer-valued generalized neural elements with a large number of entries was constructed. In the synthesis of integer-valued generic neural elements for realization of the logic algebra functions, a block representation of the Boolean function nucleus was used and based on the properties of the matrices of tolerance, coordinates of the integer vector of the structure of the generalized neural element were sequentially found.

**Keywords:** matrix of tolerance, nucleus of the Boolean function, group character, spectrum of the Boolean function.

### References

1. Grytskyk, V. V., Tsmots, I. G., Skorokhoda, O. V. (2014). Methods of parallel vertical data processing in neural networks. *Reports of the National Academy of Sciences of Ukraine*, 10, 40–44.
2. Tsmots, I. G., Skorokhoda, O. V., Vavruk, I. Ye. (2012). Principles of construction and methods of VLSI-implementation of real time neural networks. *Scientific Bulletin of Ukrainian National Forestry University*, 6, 292–300.
3. Tkachenko, R., Doroshenko, A. (2009). Neiropodibni struktury mashyny heometrychnykh peretvoren u zavdannakh intelektualnoho analizu danykh. *Kompiuterni nauky ta informatsiini tekhnolohii*, 638, 179–184.

4. Izonin, I. V., Tkachenko, R. O., Peleshko, D. D., Batiyuk, D. A. (2015). Neural network method for change resolution of images. *Information Processing Systems*, 9 (134), 30–34.
5. Marin, D., Aquino, A., Gegundez-Arias, M. E., Bravo, J. M. (2011). A New Supervised Method for Blood Vessel Segmentation in Retinal Images by Using Gray-Level and Moment Invariants-Based Features. *IEEE Transactions on Medical Imaging*, 30 (1), 146–158. doi: 10.1109/tmi.2010.2064333
6. Azarbad, M., Hakimi, S., Ebrahimzadeh, A. (2012). Automatic recognition of digital communication signal. *International journal of energy, information and communications*, 3 (4), 21–33.
7. Zaychenko, Yu. P., Dyakonova, S. V. (2011). Application of fuzzy classifier NEFCLASS to the problem of recognition of buildings in satellite images of ultrahigh resolution. *Computer science, upravlinnya that obchislyvalna tehnika*, 54, 31–35.
8. Amato, F., González-Hernández, J. L., Havel, J. (2012). Artificial neural networks combined with experimental design: A “soft” approach for chemical kinetics. *Talanta*, 93, 72–78. doi: 10.1016/j.talanta.2012.01.044
9. Brougham, D. F., Ivanova, G., Gottschalk, M., Collins, D. M., Eustace, A. J., O’Connor, R., Havel, J. (2011). Artificial Neural Networks for Classification in Metabolic Studies of Whole Cells Using <sup>1</sup>H Nuclear Magnetic Resonance. *Journal of Biomedicine and Biotechnology*, 2011, 1–8. doi: 10.1155/2011/158094
10. Barwad, A., Dey, P., Susheilia, S. (2011). Artificial neural network in diagnosis of metastatic carcinoma in effusion cytology. *Cytometry Part B: Clinical Cytometry*, 82B (2), 107–111. doi: 10.1002/cyto.b.20632
11. Geche, F., Mulesa, O., Geche, S., Vashkeba, M. (2015). Development of synthesis method of predictive schemes based on basic predictive models. *Technology Audit and Production Reserves*, 3 (2 (23)), 36–41. doi: 10.15587/2312-8372.2015.44932
12. Dey, P., Lamba, A., Kumari, S., Marwaha, N. (2011). Application of an artificial neural network in the prognosis of chronic myeloid leukemia. *Analytical and quantitative cytology and histology/the International Academy of Cytology and American Society of Cytology*, 33 (6), 335–339.
13. Geche, F., Batiyuk, A., Mulesa, O., Vashkeba, M. (2015). Development of effective time series forecasting model. *International Journal of Advanced Research in Computer Engineering & Technology*, 4 (12), 4377–4386.
14. Liu, A., Zhu, Q. (2011). Automatic modulation classification based on the combination of clustering and neural network. *The Journal of China Universities of Posts and Telecommunications*, 18 (4), 13–38. doi: 10.1016/s1005-8885(10)60077-5
15. Pathok, A., Wadhvani, A. K. (2012). Data Compression of ECG Signals Using Error Back Propagation (EBP) Algorithm. *International Journal of Engineering and Advance Technology (IJEAT)*, 1 (4), 2249–8958.
16. Bodyanskiy, Y., Grimm, P., Mashtalir, S., Vinarski, V. (2010). Fast Training of Neural Networks for Image Compression. *Lecture Notes in Computer Science*, 165–173. doi: 10.1007/978-3-642-14400-4\_13
17. Shovgun, N. V. (2013). Analiz effektivnosti nechetkih neyronnyh setey v zadache otsenki kreditnogo riska. In: *formation technologies & knowledge. ITHEA IBS ISC*, 7, 286–293.
18. Kertis, Ch., Rayner, I. (1969). *Teoriya predstavleniy konechnykh grupp i assotsiativnykh algebr*. Moscow: Nauka, 667.
19. Golubov, B. I., Efimov, A. V., Skvortsov, V. A. (1987). *Ryady i preobrazovaniya Uolsha. Teoriya i primeneniya*. Moscow: Nauka, 343.
20. Dertouzos, M. (1967). *Porogovaya logika*. Moscow: Mir, 342.
21. Ayzenberg, N. N., Bovdi, A. A., Gergo, E. Y., Geche, F. E. (1980). *Nekotorye algebraicheskie aspekty porogovoy logiki*. *Kibernetika*, 2, 26–30.
22. Yadzhima, S., Ibaraki, T. (1969). *Nizhnnyaya otsenka chisla porogovykh funktsiy*. Vol. 6. *Kiberneticheskiy sbornik: novaya seriya*. Moscow: Mir, 72–81.
23. Geche, F., Mulesa, O., Buchok, V. (2017). Verification of realizability of boolean functions by a neural element with a threshold activation function. *Eastern-European Journal of Enterprise Technologies*, 1 (4 (85)), 30–40. doi: 10.15587/1729-4061.2017.90917

---

**DOI: 10.15587/1729-4061.2017.108976**  
**DESIGN OF THE UNIVERSAL CLASSIFIER**  
**AS A RBF NETWORK BASED ON THE CART**  
**SOLUTION TREE (p. 63-71)**

**Lyudmila Dobrovskaya**

National Technical University of Ukraine «Igor Sikorsky Kyiv Polytechnic Institute», Kyiv, Ukraine  
**ORCID:** <http://orcid.org/0000-0002-4055-6834>

**Iryna Dobrovskaya**

UkrSibbank, Kyiv, Ukraine  
**ORCID:** <http://orcid.org/0000-0003-4537-2011>

The aim of the paper was to develop a universal classifier in the form of a radial basis function network (RBF network) based on the Gaussian function and the CART Solution Tree. The examples of diseases diagnostics classifier were considered. It is noted that during the classifier development, it is necessary to determine the number of RBF neurons and the values of parameters of these neurons (centre, dispersion). For this purpose, a method that allows splitting the space of features into relatively homogeneous domains in the form of hyperparallelepipeds, each of which is associated with one of the RBF neurons, is proposed. The number of RBF neurons and parameters of these neurons are determined automatically directly based on the CART Solution Tree.

As a result of the research, it was found that the proposed classifiers show the highest efficiency on the learning set with a minimal Solution Tree reduction (accuracy from 80 % to 95 %). It was shown that for two and more classes the accuracy of these classifiers on the test set makes 79 % and more, however, provided that the appropriate data sample for the learning set is selected. The possibility of using the RBF network based on the Gaussian function and the CART Solution Tree in the healthcare system for the diseases diagnostics and medical systems (or devices) assistance during decision-making support was proved.

The obtained results could be further applied to improve the universal classifier development method based on the RBF network.



**Keywords:** universal classifier, neural network, RBF network, CART Solution Tree, decision-making support.

### References

1. Dobrovska, L. M., Dobrovska, I. A. (2015). Teoriya ta praktyka neironnykh merezh. Kyiv: NTUU «KPI» Vydvo «Politekhnik», 396.
2. Osovskiy, S. (2004). Neyronnye seti dlya obrabotki informaciy. Moscow: Finansy i statistika, 343.
3. Haykin, S. (2006). Neyronnye seti. Moscow: Izdatel'skiy dom «Vil'yams», 1104.
4. Hagan, M., Demuth, H., Beale, M. (2010). Neural Network Design. USA: Colorado University Bookstore.
5. Kitayama, Y., Kitayama, S., Yamazaki, K. (2008). Pattern Classification by RBF Network. The Proceedings of Design & Systems Conference, 2008.18, 330–332. doi: 10.1299/jsmedsd.2008.18.330
6. Kubat, M. (1998). Decision trees can initialize radial-basis function networks. IEEE Transactions on Neural Networks, 9 (5), 813–821. doi: 10.1109/72.712154
7. Safish, M., Joseph, R. (2012). Data classification with neural classifier using radial basis function with data reduction using hierarchical clustering. ICTACT Journal on Soft Computing, 2 (3), 348–352. doi: 10.21917/ijsc.2012.0054
8. Jia, W., Zhao, D., Ding, L. (2016). An optimized RBF neural network algorithm based on partial least squares and genetic algorithm for classification of small sample. Applied Soft Computing, 48, 373–384. doi: 10.1016/j.asoc.2016.07.037
9. Jang, J.-S., Sun, C.-T., Mizutani, E. (1997). Neuro-fuzzy and soft computing: a computation approach to learning and machine intelligence. USA, 614.
10. Blake, C. L., Merz, C. J. (1998). UCI Repository of machine learning databases. University of California Irvine, Department of Information and Computer Science, Available at <http://archive.ics.uci.edu/ml/datasets.html>
11. Faynzil'berg, L. (2009). Garantirovannaya ocenka ehffektivnosti diagnosticheskikh testov na osnove usilennogo ROC-analiza. Upravlyayushchie sistemy i mashiny, 5, 3–13.
12. Flach, P., Wu, S. (2003). Repairing concavities in ROC curves. Proc. UK Workshop on Comp. Intel, 38–44.