

Mathematical Subject Classification: 34C29, 34A38
UDC 517.911.5

O. Osadcha, N. Skripnik
Odesa I. I. Mechnikov National University

PARTIAL AVERAGING OF IMPULSIVE HYBRID SYSTEMS

Осадча О., Скрипник Н. Схема часткового усереднення для одного класу імпульсних змішаних систем. У статті розглядається обґрунтування схеми часткового усереднення для одного класу імпульсних змішаних систем, коли одне з рівнянь є імпульсним диференціальним рівнянням з похідною Хукухари, а друге – звичайним імпульсним диференціальним рівнянням.

Ключові слова: метод усереднення, змішана система, похідна Хукухари.

Осадчая О., Скрипник Н. Схема частичного усреднения для одного класса импульсных смешанных систем. В статье рассматривается обоснование схемы частичного усреднения для одного класса импульсных смешанных систем, когда одно из уравнений является импульсным дифференциальным уравнением с производной Хукухари, а второе – обыкновенным импульсным дифференциальным уравнением.

Ключевые слова: метод усреднения, смешанная система, производная Хукухари.

Osadcha O., Skripnik N. Partial averaging of impulsive hybrid systems. This paper contains the substantiation of the scheme of partial averaging for one class of impulsive hybrid systems where one equation is an impulsive differential equation with Hukuhara derivative and the second one is an impulsive ordinary differential equation.

Key words: averaging method, hybrid system, Hukuhara derivative.

INTRODUCTION.

In practice, there are often considered the so-called hybrid systems — systems that contain equations of different nature: for example, one of the equations is an equation in partial derivatives and the other one is an ordinary differential equation, or one of the equations is discrete and the other one is differential, etc. In this paper we consider the case of a hybrid system, where one of the equations is a differential equation with Hukuhara derivative and the second one is an ordinary differential equation. The interest in such systems follows from the fact, that some parameters of the model may be accurate, while the rest may contain the noise, errors and inaccuracies.

MAIN DEFINITIONS.

Development of the theory of multivalued mappings led to the question what should be understood as a derivative of a multivalued mapping. The main cause of difficulties for the inducting of such definition was the nonlinearity of the space $comp(R^n)$, which led to the absence of the concept of difference. There are several approaches to define the difference of two sets, one of them is the Hukuhara difference.

Let $conv(R^n)$ be the family of all nonempty compact convex subsets of R^n with the Hausdorff metric

$$h(A, B) = \max\{\max_{a \in A} \min_{b \in B} \|a - b\|, \max_{b \in B} \min_{a \in A} \|a - b\|\},$$

where $\|\cdot\|$ denotes the Euclidean norm in R^n .

Definition 1. [5] Let $X, Y \in \text{conv}(R^n)$. The set $Z \in \text{conv}(R^n)$, where $X = Y + Z$, is called the Hukuhara difference of sets X and Y and is denoted by $X \overset{h}{-} Y$.

Along with the inducted difference there appeared the concept of derivative.

Definition 2. [5] A multivalued mapping $X : I \rightarrow \text{conv}(R^n), I \subset R$, is called differentiable in the sense of Hukuhara at point $t \in I$ if there exists such $D_H X(t) \in \text{conv}(R^n)$ that the limits

$$\lim_{\Delta t \downarrow 0} \frac{1}{\Delta t} \left(X(t + \Delta t) \overset{h}{-} X(t) \right), \quad \lim_{\Delta t \downarrow 0} \frac{1}{\Delta t} \left(X(t) \overset{h}{-} X(t - \Delta t) \right)$$

exist and are equal to $D_H X(t)$. The set $D_H X(t)$ is called the Hukuhara derivative of the multivalued mapping $X : I \rightarrow \text{conv}(R^n)$ at point t .

In M. Hukuhara papers [5] along with the concept of derivative the concept of the integral of a multivalued mapping was inducted and the connection between those two concepts was found. In 1969 F. S. de Blasi and F. Iervolino were first to consider differential equation with the Hukuhara derivative [1, 2, 3, 4]. Its solution is a multivalued mapping. After that various existence and uniqueness theorems were proved, the stability of solutions of this type of equations was considered, integro-differential equations, impulsive differential equations, differential equations with fractional derivatives, control differential equations with Hukuhara derivative were considered, the possibility of applying some averaging schemes for such type of equations was investigated [9, 7, 12, 13, 8, 10, 11, 6].

Consider the hybrid system

$$\begin{cases} D_H X = F(t, X, y), X(t_0) = X_0, \\ \dot{y} = g(t, X, y), y(t_0) = y_0, \end{cases} \quad (1)$$

where $I = [t_0, T] \subset R$; $X : I \rightarrow \text{conv}(R^n)$ is a multivalued mapping; $y : I \rightarrow R^m$ is a vector function; $F : I \times \text{conv}(R^n) \times R^m \rightarrow \text{conv}(R^n)$ is a multivalued mapping; $g : I \times \text{conv}(R^n) \times R^m \rightarrow R^m$ is a vector function; $X_0 \in \text{conv}(R^n)$, $y_0 \in R^m$.

Consider a class S of pairs $(X(\cdot), y(\cdot))$, where $X(\cdot)$ is a continuously differentiable on I in the sense of Hukuhara multivalued mapping, $y(\cdot)$ is a continuously differentiable on I vector-function.

Definition 3. A pair $(X(\cdot), y(\cdot)) \in S$ is called a solution of system (1), if for all $t \in I$ the following equalities fulfill $D_H X(t) = F(t, X(t), y(t))$, $\dot{y}(t) = g(t, X(t), y(t))$ and $X(t_0) = X_0$, $y(t_0) = y_0$.

Theorem 1. Let in the domain

$$Q = \{(t, X, y) : t_0 \leq t \leq t_0 + a, h(X, X_0) \leq b, \|y - y_0\| \leq c\}$$

the multivalued mapping $F(t, X, y)$ and the vector function $g(t, X, y)$ are continuous and satisfy the Lipschitz condition in variables X and y , i. e. there exists such constant $\lambda > 0$ that

$$h(F(t, X_1, y_1), F(t, X_2, y_2)) \leq \lambda [h(X_1, X_2) + \|y_1 - y_2\|],$$

$$\|g(t, X_1, y_1) - g(t, X_2, y_2)\| \leq \lambda [h(X_1, X_2) + \|y_1 - y_2\|].$$

Then system (1) has the unique solution defined on the interval $[t_0, t_0 + d]$ where $d = \min\left(a, \frac{b}{M}, \frac{c}{M}\right)$, constant M satisfies the inequalities $|F(t, X, y)| \leq M$ and $\|g(t, X, y)\| \leq M$ in the domain Q .

MAIN RESULTS.

Consider impulsive hybrid system with a small parameter

$$\begin{aligned} D_H X &= \varepsilon F(t, X, y), \quad t \neq \tau_i, \quad X(0) = X_0, \\ \dot{y} &= \varepsilon g(t, X, y), \quad t \neq \tau_i, \quad y(0) = y_0, \end{aligned} \tag{2}$$

$$\Delta X|_{t=\tau_i} = \varepsilon I_i(X, y), \quad \Delta y|_{t=\tau_i} = \varepsilon J_i(X, y), \tag{3}$$

where $t \in R_+$; the moments of impulses $\tau_{i+1} > \tau_i$; $X : R_+ \rightarrow \text{conv}(R^n)$ is a multivalued mapping; $y : R_+ \rightarrow R^m$ is a vector function; $F : R_+ \times \text{conv}(R^n) \times R^m \rightarrow \text{conv}(R^n)$, $I_i : \text{conv}(R^n) \times R^m \rightarrow \text{conv}(R^n)$ are multivalued mappings; $g : R_+ \times \text{conv}(R^n) \times R^m \rightarrow R^m$, $J_i : \text{conv}(R^n) \times R^m \rightarrow R^m$ are vector functions; $X_0 \in \text{conv}(R^n)$, $y_0 \in R^m$.

Definition 4. A pair $(X(\cdot), y(\cdot))$ is called a solution of system (2), (3) if it is a solution of system (2) on intervals between moments of impulses and satisfies impulse condition (3) in the points of impulses τ_i .

It's easy to notice, that the existence and the uniqueness of a solution of system (2), (3) holds if the right sides of equations (2) satisfy Theorem 1 on intervals between moments of impulses.

Consider the following partially averaged system:

$$\begin{aligned} D_H \bar{X} &= \varepsilon \bar{F}(t, \bar{X}, \bar{y}), \quad \bar{X}(0) = X_0, \\ \dot{\bar{y}} &= \varepsilon \bar{g}(t, \bar{X}, \bar{y}), \quad \bar{y}(0) = y_0, \end{aligned} \tag{4}$$

$$\Delta X|_{t=\sigma_j} = \varepsilon \bar{I}_j(X, y), \quad \Delta y|_{t=\sigma_j} = \varepsilon \bar{J}_j(X, y), \tag{5}$$

where

$$\begin{aligned} \lim_{T \rightarrow \infty} \frac{1}{T} h \left(\int_0^T F(t, X, y) dt + \sum_{0 \leq \tau_i < T} I_i(X, y), \int_0^T \bar{F}(t, X, y) dt + \sum_{0 \leq \sigma_j < T} \bar{I}_j(X, y) \right) &= 0, \\ \lim_{T \rightarrow \infty} \frac{1}{T} \left\| \int_0^T [g(t, X, y) - \bar{g}(t, X, y)] dt + \sum_{0 \leq \tau_i < T} J_i(X, y) - \sum_{0 \leq \sigma_j < T} \bar{J}_j(X, y) \right\| &= 0. \end{aligned} \tag{6}$$

The following theorem that proves the closeness of solutions of systems (2), (3) and (4), (5) holds.

Theorem 2. Let in the domain $Q = \{(t, X, y) : t \geq 0, X \in D_1, y \in D_2\}$ the following conditions hold:

1) the multivalued mappings $F(t, X, y), \bar{F}(t, X, y)$ and vector functions $g(t, X, y), \bar{g}(t, X, y)$ are continuous in t , uniformly bounded with constant M and satisfy the Lipschitz condition in X and y with constant λ ;

2) the multivalued mappings $I_i(X, y), \bar{I}_j(X, y)$ and vector functions $J_i(X, y), \bar{J}_j(X, y)$ are uniformly bounded and satisfy the Lipschitz condition in X and y with constant λ ;

3) limits (6) exist uniformly with respect to $X \in D_1$ and $y \in D_2$;

4) there exists such constant $0 < d \leq \infty$ that

$$\frac{1}{T}i(t, t+T) \leq d, \quad \frac{1}{T}j(t, t+T) \leq d,$$

where $i(t, t+T)[j(t, t+T)]$ are the numbers of points of a sequence $\{\tau_i\}[\{\sigma_j\}]$ on the interval $[t, t+T)$.

5) the solution $(\bar{X}(t), \bar{y}(t))$ of system (4), (5) with the initial condition $\bar{X}(0) = X_0 \in D'_1 \subset D_1, \bar{y}(0) = y_0 \in D'_2 \subset D_2$ is defined for all $t \geq 0, \varepsilon \in (0, \bar{\varepsilon}]$ and $\bar{X}(t)$ belongs with some ρ -neighborhood to the domain $D_1, \bar{y}(t)$ belongs with some ξ -neighborhood to the domain D_2 .

Then for any $\eta > 0$ and $L > 0$ there exists such $\varepsilon_0(\eta, L) \in (0, \bar{\varepsilon}]$, that for $0 < \varepsilon \leq \varepsilon_0$ and $0 \leq t \leq L\varepsilon^{-1}$ the following inequalities fulfill:

$$h(X(t), \bar{X}(t)) \leq \eta, \|y(t) - \bar{y}(t)\| \leq \eta,$$

where $(X(\cdot), y(\cdot))$ and $(\bar{X}(\cdot), \bar{y}(\cdot))$ are the solutions of systems (2), (3) and (4), (5) with the initial conditions $X(0) = \bar{X}(0) \in D'_1, y(0) = \bar{y}(0) \in D'_2$.

Proof. From conditions 1)–3) of the theorem it follows that systems (2), (3) and (4), (5) have unique solutions that are defined for $t \geq 0$ if $X(t)$ and $y(t)$ (accordingly $\bar{X}(t)$ and $\bar{y}(t)$) belong to the domain $D_1 \times D_2$.

Replace systems (2), (3) and (4), (5) with the equivalent system of integral equations:

$$\begin{cases} X(t) = X_0 + \varepsilon \int_0^t F(s, X(s), y(s)) ds + \varepsilon \sum_{0 \leq \tau_i < t} I_i(X(\tau_i), y(\tau_i)), \\ y(t) = y_0 + \varepsilon \int_0^t g(s, X(s), y(s)) ds + \varepsilon \sum_{0 \leq \tau_i < t} J_i(X(\tau_i), y(\tau_i)), \end{cases} \quad (7)$$

$$\begin{cases} \bar{X}(t) = X_0 + \varepsilon \int_0^t \bar{F}(s, \bar{X}(s), \bar{y}(s)) ds + \varepsilon \sum_{0 \leq \sigma_j < t} \bar{I}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j)), \\ \bar{y}(t) = y_0 + \varepsilon \int_0^t \bar{g}(s, \bar{X}(s), \bar{y}(s)) ds + \varepsilon \sum_{0 \leq \sigma_j < t} \bar{J}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j)). \end{cases} \quad (8)$$

Then

$$\begin{aligned}
 & h(X(t), \bar{X}(t)) = \\
 = & h\left(X_0 + \varepsilon \int_0^t F(s, X(s), y(s))ds + \varepsilon \sum_{0 \leq \tau_i < t} I_i(X(\tau_i), y(\tau_i)), \right. \\
 & \left. X_0 + \varepsilon \int_0^t \bar{F}(s, \bar{X}(s), \bar{y}(s))ds + \varepsilon \sum_{0 \leq \sigma_j < t} \bar{I}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j))\right) = \\
 = & \varepsilon h\left(\int_0^t F(s, X(s), y(s))ds + \sum_{0 \leq \tau_i < t} I_i(X(\tau_i), y(\tau_i)), \right. \\
 & \left. \int_0^t \bar{F}(s, \bar{X}(s), \bar{y}(s))ds + \sum_{0 \leq \sigma_j < t} \bar{I}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j))\right) \leq \\
 \leq & \varepsilon h\left(\int_0^t F(s, X(s), y(s))ds, \int_0^t F(s, \bar{X}(s), \bar{y}(s))ds\right) + \\
 + & \varepsilon h\left(\sum_{0 \leq \tau_i < t} I_i(X(\tau_i), y(\tau_i)), \sum_{0 \leq \tau_i < t} I_i(\bar{X}(\tau_i), \bar{y}(\tau_i))\right) + \\
 + & \varepsilon h\left(\int_0^t F(s, \bar{X}(s), \bar{y}(s))ds + \sum_{0 \leq \tau_i < t} I_i(\bar{X}(\tau_i), \bar{y}(\tau_i)), \right. \\
 & \left. \int_0^t \bar{F}(s, \bar{X}(s), \bar{y}(s))ds + \varepsilon \sum_{0 \leq \sigma_j < t} \bar{I}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j))\right) \leq \\
 \leq & \varepsilon \int_0^t h(F(s, X(s), y(s)), F(s, \bar{X}(s), \bar{y}(s))) ds + \\
 + & \varepsilon \sum_{0 \leq \tau_i < t} h(I_i(X(\tau_i), y(\tau_i)), I_i(\bar{X}(\tau_i), \bar{y}(\tau_i))) + \\
 + & \varepsilon h\left(\int_0^t F(s, \bar{X}(s), \bar{y}(s))ds + \sum_{0 \leq \tau_i < t} I_i(\bar{X}(\tau_i), \bar{y}(\tau_i)), \right. \\
 & \left. \int_0^t \bar{F}(s, \bar{X}(s), \bar{y}(s))ds + \varepsilon \sum_{0 \leq \sigma_j < t} \bar{I}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j))\right) \leq \\
 \leq & \varepsilon \lambda \int_0^t [h(X(s), \bar{X}(s)) + |y(s) - \bar{y}(s)|] ds + \\
 + & \varepsilon \lambda \sum_{0 \leq \tau_i < t} [h(X(\tau_i), \bar{X}(\tau_i)) + |y(\tau_i) - \bar{y}(\tau_i)|] + \\
 + & \varepsilon h\left(\int_0^t F(s, \bar{X}(s), \bar{y}(s))ds + \sum_{0 \leq \tau_i < t} I_i(\bar{X}(\tau_i), \bar{y}(\tau_i)), \right. \\
 & \left. \int_0^t \bar{F}(s, \bar{X}(s), \bar{y}(s))ds + \varepsilon \sum_{0 \leq \sigma_j < t} \bar{I}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j))\right). \tag{9}
 \end{aligned}$$

Similarly,

$$\begin{aligned}
 \|y(t) - \bar{y}(t)\| & \leq \varepsilon \lambda \int_0^t [h(X(s), \bar{X}(s)) + \|y(s) - \bar{y}(s)\|] ds + \\
 + & \varepsilon \lambda \sum_{0 \leq \tau_i < t} [h(X(\tau_i), \bar{X}(\tau_i)) + \|y(\tau_i) - \bar{y}(\tau_i)\|] + \\
 + & \varepsilon \left\| \int_0^t g(s, \bar{X}(s), \bar{y}(s))ds + \sum_{0 \leq \tau_i < t} J_i(\bar{X}(\tau_i), \bar{y}(\tau_i)) - \right.
 \end{aligned}$$

$$\left\| - \int_0^t \bar{g}(s, \bar{X}(s), \bar{y}(s)) ds - \varepsilon \sum_{0 \leq \sigma_j < t} \bar{J}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j)) \right\|. \quad (10)$$

Divide the interval $[0, L\varepsilon^{-1}]$ in m equal intervals by the points $t_p = \frac{pL}{\varepsilon m}$. Define by $(\bar{X}_p, \bar{y}_p) = (\bar{X}(t_p), \bar{y}(t_p))$ the solution of system (4), (5) in division points.

Let us estimate in the interval $[t_k, t_{k+1}]$, where $0 \leq k \leq m - 1$ the expression

$$\begin{aligned} & \varepsilon h \left(\int_0^t F(s, \bar{X}(s), \bar{y}(s)) ds + \sum_{0 \leq \tau_i < t} I_i(\bar{X}(\tau_i), \bar{y}(\tau_i)), \right. \\ & \left. \int_0^t \bar{F}(s, \bar{X}(s), \bar{y}(s)) ds + \varepsilon \sum_{0 \leq \sigma_j < t} \bar{I}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j)) \right) = \\ & = \varepsilon h \left(\sum_{p=0}^{k-1} \left[\int_{t_p}^{t_{p+1}} F(s, \bar{X}(s), \bar{y}(s)) ds + \sum_{t_p \leq \tau_i < t_{p+1}} I_i(\bar{X}(\tau_i), \bar{y}(\tau_i)) \right] + \right. \\ & \quad \left. + \int_{t_k}^t F(s, \bar{X}(s), \bar{y}(s)) ds + \sum_{t_k \leq \tau_i < t} I_i(\bar{X}(\tau_i), \bar{y}(\tau_i)), \right. \\ & \quad \left. \sum_{p=0}^{k-1} \left[\int_{t_p}^{t_{p+1}} \bar{F}(s, \bar{X}(s), \bar{y}(s)) ds + \sum_{t_p \leq \sigma_j < t_{p+1}} \bar{I}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j)) \right] + \right. \\ & \quad \left. + \int_{t_k}^t \bar{F}(s, \bar{X}(s), \bar{y}(s)) ds + \sum_{t_k \leq \sigma_j < t} \bar{I}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j)) \right) \leq \\ & \leq \varepsilon h \left(\sum_{p=0}^{k-1} \left[\int_{t_p}^{t_{p+1}} F(s, \bar{X}(s), \bar{y}(s)) ds + \sum_{t_p \leq \tau_i < t_{p+1}} I_i(\bar{X}(\tau_i), \bar{y}(\tau_i)) \right] + \right. \\ & \quad \left. \sum_{p=0}^{k-1} \left[\int_{t_p}^{t_{p+1}} \bar{F}(s, \bar{X}(s), \bar{y}(s)) ds + \sum_{t_p \leq \sigma_j < t_{p+1}} \bar{I}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j)) \right] \right) + \\ & \quad + \varepsilon h \left(\int_{t_k}^t F(s, \bar{X}(s), \bar{y}(s)) ds + \sum_{t_k \leq \tau_i < t} I_i(\bar{X}(\tau_i), \bar{y}(\tau_i)), \right. \\ & \quad \left. \int_{t_k}^t \bar{F}(s, \bar{X}(s), \bar{y}(s)) ds + \sum_{t_k \leq \sigma_j < t} \bar{I}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j)) \right) \leq \\ & \leq \varepsilon \sum_{p=0}^{k-1} \left[h \left(\int_{t_p}^{t_{p+1}} F(s, \bar{X}(s), \bar{y}(s)) ds + \sum_{t_p \leq \tau_i < t_{p+1}} I_i(\bar{X}(\tau_i), \bar{y}(\tau_i)), \int_{t_p}^{t_{p+1}} F(s, \bar{X}_p, \bar{y}_p) ds + \sum_{t_p \leq \tau_i < t_{p+1}} I_i(\bar{X}_p, \bar{y}_p) \right) + \right. \\ & \quad \left. + h \left(\int_{t_p}^{t_{p+1}} F(s, \bar{X}_p, \bar{y}_p) ds + \sum_{t_p \leq \tau_i < t_{p+1}} I_i(\bar{X}_p, \bar{y}_p), \int_{t_p}^{t_{p+1}} \bar{F}(s, \bar{X}_p, \bar{y}_p) ds + \sum_{t_p \leq \sigma_j < t_{p+1}} \bar{I}_j(\bar{X}_p, \bar{y}_p) \right) + \right. \\ & \quad \left. + h \left(\int_{t_p}^{t_{p+1}} \bar{F}(s, \bar{X}_p, \bar{y}_p) ds + \sum_{t_p \leq \sigma_j < t_{p+1}} \bar{I}_j(\bar{X}_p, \bar{y}_p), \int_{t_p}^{t_{p+1}} F(s, \bar{X}(s), \bar{y}(s)) ds + \sum_{t_p \leq \sigma_j < t_{p+1}} \bar{I}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j)) \right) \right] + \\ & \quad + \varepsilon \left[h \left(\int_{t_k}^t F(s, \bar{X}(s), \bar{y}(s)) ds + \sum_{t_k \leq \tau_i < t} I_i(\bar{X}(\tau_i), \bar{y}(\tau_i)), \int_{t_k}^t F(s, \bar{X}_k, \bar{y}_k) ds + \sum_{t_k \leq \tau_i < t} I_i(\bar{X}_k, \bar{y}_k) \right) + \right. \\ & \quad \left. + h \left(\int_{t_k}^t F(s, \bar{X}_k, \bar{y}_k) ds + \sum_{t_k \leq \tau_i < t} I_i(\bar{X}_k, \bar{y}_k), \int_{t_k}^t \bar{F}(s, \bar{X}_k, \bar{y}_k) ds + \sum_{t_k \leq \sigma_j < t} \bar{I}_j(\bar{X}_k, \bar{y}_k) \right) + \right. \end{aligned}$$

$$\begin{aligned}
 & +h \left(\int_{t_k}^t \bar{F}(s, \bar{X}_k, \bar{y}_k) ds + \sum_{t_k \leq \sigma_j < t} \bar{I}_j(\bar{X}_k, \bar{y}_k), \int_{t_k}^t \bar{F}(s, \bar{X}(s), \bar{y}(s)) ds + \sum_{t_k \leq \sigma_j < t} \bar{I}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j)) \right) \Big] \leq \\
 & \leq \varepsilon \sum_{p=0}^{k-1} \left[\int_{t_p}^{t_{p+1}} h(F(s, \bar{X}(s), \bar{y}(s)), F(s, \bar{X}_p, \bar{y}_p)) ds + \sum_{t_p \leq \tau_i < t_{p+1}} h(I_i(\bar{X}(\tau_i), \bar{y}(\tau_i)), I_i(\bar{X}_p, \bar{y}_p)) \right) + \\
 & +h \left(\int_{t_p}^{t_{p+1}} F(s, \bar{X}_p, \bar{y}_p) ds + \sum_{t_p \leq \tau_i < t_{p+1}} I_i(\bar{X}_p, \bar{y}_p), \int_{t_p}^{t_{p+1}} \bar{F}(s, \bar{X}_p, \bar{y}_p) ds + \sum_{t_p \leq \sigma_j < t_{p+1}} \bar{I}_j(\bar{X}_p, \bar{y}_p) \right) + \\
 & + \int_{t_p}^{t_{p+1}} h(\bar{F}(s, \bar{X}_p, \bar{y}_p), \bar{F}(s, \bar{X}(s), \bar{y}(s))) ds + \sum_{t_p \leq \sigma_j < t_{p+1}} h(\bar{I}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j)), \bar{I}_j(\bar{X}_p, \bar{y}_p)) \Big] + \\
 & + \varepsilon \left[\int_{t_k}^t h(F(s, \bar{X}(s), \bar{y}(s)), F(s, \bar{X}_k, \bar{y}_k)) ds + \sum_{t_k \leq \tau_i < t} h(I_i(\bar{X}(\tau_i), \bar{y}(\tau_i)), I_i(\bar{X}_k, \bar{y}_k)) \right) + \\
 & +h \left(\int_{t_k}^t F(s, \bar{X}_k, \bar{y}_k) ds + \sum_{t_k \leq \tau_i < t} I_i(\bar{X}_k, \bar{y}_k), \int_{t_k}^t \bar{F}(s, \bar{X}_k, \bar{y}_k) ds + \sum_{t_k \leq \sigma_j < t} \bar{I}_j(\bar{X}_k, \bar{y}_k) \right) + \\
 & + \int_{t_k}^t h(\bar{F}(s, \bar{X}_k, \bar{y}_k), \bar{F}(\bar{X}(s), \bar{y}(s))) ds + \sum_{t_k \leq \sigma_j < t} h(\bar{I}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j)), \bar{I}_j(\bar{X}_k, \bar{y}_k)) \Big] \leq \\
 & \leq \varepsilon \sum_{p=0}^k \left[\int_{t_p}^{t_{p+1}} h(F(s, \bar{X}(s), \bar{y}(s)), F(s, \bar{X}_p, \bar{y}_p)) ds + \sum_{t_p \leq \tau_i < t_{p+1}} h(I_i(\bar{X}(\tau_i), \bar{y}(\tau_i)), I_i(\bar{X}_p, \bar{y}_p)) \right) + \\
 & + \int_{t_p}^{t_{p+1}} h(\bar{F}(s, \bar{X}_p, \bar{y}_p), \bar{F}(s, \bar{X}(s), \bar{y}(s))) ds + \sum_{t_p \leq \sigma_j < t_{p+1}} h(\bar{I}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j)), \bar{I}_j(\bar{X}_p, \bar{y}_p)) \Big] + \\
 & + \varepsilon \sum_{p=0}^{k-1} h \left(\int_{t_p}^{t_{p+1}} F(s, \bar{X}_p, \bar{y}_p) ds + \sum_{t_p \leq \tau_i < t_{p+1}} I_i(\bar{X}_p, \bar{y}_p), \int_{t_p}^{t_{p+1}} \bar{F}(s, \bar{X}_p, \bar{y}_p) ds + \sum_{t_p \leq \sigma_j < t_{p+1}} \bar{I}_j(\bar{X}_p, \bar{y}_p) \right) + \\
 & + \varepsilon h \left(\int_{t_k}^t F(s, \bar{X}_k, \bar{y}_k) ds + \sum_{t_k \leq \tau_i < t} I_i(\bar{X}_k, \bar{y}_k), \int_{t_k}^t \bar{F}(s, \bar{X}_k, \bar{y}_k) ds + \sum_{t_k \leq \sigma_j < t} \bar{I}_j(\bar{X}_k, \bar{y}_k) \right).
 \end{aligned}$$

Similarly

$$\begin{aligned}
 & \varepsilon \left\| \int_0^t g(s, \bar{X}(s), \bar{y}(s)) ds + \sum_{0 \leq \tau_i < t} J_i(\bar{X}(\tau_i), \bar{y}(\tau_i)) - \int_0^t \bar{g}(s, \bar{X}(s), \bar{y}(s)) ds - \right. \\
 & \left. - \sum_{0 \leq \sigma_j < t} \bar{J}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j)) \right\| \leq \varepsilon \sum_{p=0}^k \left[\int_{t_p}^{t_{p+1}} \|g(s, \bar{X}(s), \bar{y}(s)) - g(s, \bar{X}_p, \bar{y}_p)\| ds + \right. \\
 & + \sum_{t_p \leq \tau_i < t_{p+1}} \|J_i(\bar{X}(\tau_i), \bar{y}(\tau_i)) - J_i(\bar{X}_p, \bar{y}_p)\| + \int_{t_p}^{t_{p+1}} \|\bar{g}(s, \bar{X}_p, \bar{y}_p) - \bar{g}(s, \bar{X}(s), \bar{y}(s))\| ds + \\
 & \left. + \sum_{t_p \leq \sigma_j < t_{p+1}} \|\bar{J}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j)) - \bar{J}_j(\bar{X}_p, \bar{y}_p)\| \right] + \\
 & + \varepsilon \sum_{p=0}^{k-1} \left\| \int_{t_p}^{t_{p+1}} [g(s, \bar{X}_p, \bar{y}_p) - \bar{g}(s, \bar{X}_p, \bar{y}_p)] ds + \sum_{t_p \leq \tau_i < t_{p+1}} J_i(\bar{X}_p, \bar{y}_p) - \sum_{t_p \leq \sigma_j < t_{p+1}} \bar{J}_j(\bar{X}_p, \bar{y}_p) \right\| + \\
 & + \varepsilon \left\| \int_{t_k}^t [g(s, \bar{X}_k, \bar{y}_k) - \bar{g}(s, \bar{X}_k, \bar{y}_k)] ds + \sum_{t_k \leq \tau_i < t} J_i(\bar{X}_k, \bar{y}_k) - \sum_{t_k \leq \sigma_j < t} \bar{J}_j(\bar{X}_k, \bar{y}_k) \right\|.
 \end{aligned}$$

Notice that

$$\begin{aligned}
h(\bar{X}(s), \bar{X}_p) &= h(\bar{X}(s), \bar{X}(t_p)) \leq \varepsilon \int_s^s h(\bar{F}(\bar{X}(v), \bar{y}(v)), \{0\}) dv + \\
&+ \sum_{t_p \leq \sigma_j < s} h(\bar{I}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j)), \{0\}) \leq \varepsilon M(d+1)(s - t_p), \\
\|\bar{y}(s) - \bar{y}_p\| &= \|\bar{y}(s) - \bar{y}(t_p)\| \leq \\
\leq \varepsilon \int_{t_p}^s \|\bar{g}(\bar{X}(v), \bar{y}(v))\| dv + \sum_{t_p \leq \sigma_j < s} \|\bar{J}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j))\| &\leq \varepsilon M(d+1)(s - t_p).
\end{aligned}$$

Then

$$\begin{aligned}
&\varepsilon \sum_{p=0}^k \int_{t_p}^{t_{p+1}} h(F(s, \bar{X}(s), \bar{y}(s)), F(s, \bar{X}_p, \bar{y}_p)) ds \leq \\
&\leq \varepsilon \sum_{p=0}^k \int_{t_p}^{t_{p+1}} \lambda [h(\bar{X}(s), \bar{X}_p) + \|\bar{y}(s) - \bar{y}_p\|] ds \leq \\
&\leq \varepsilon \lambda \cdot 2\varepsilon M(d+1) \sum_{p=0}^k \int_{t_p}^{t_{p+1}} (s - t_p) ds = 2\varepsilon^2 \lambda M(d+1) \sum_{p=0}^k \frac{(t_{p+1} - t_p)^2}{2} = \\
&= \varepsilon^2 \lambda M(d+1) \cdot (k+1) \cdot \left(\frac{L}{\varepsilon m}\right)^2 \leq \frac{\lambda M L^2 (d+1)}{m}, \\
&\varepsilon \sum_{p=0}^k \sum_{t_p \leq \tau_i < t_{p+1}} h(I_i(\bar{X}(\tau_i), \bar{y}(\tau_i)), I_i(\bar{X}_p, \bar{y}_p)) \leq \\
&\leq \varepsilon \sum_{p=0}^k \sum_{t_p \leq \tau_i < t_{p+1}} \lambda [h(\bar{X}(\tau_i), \bar{X}_p) + \|\bar{y}(\tau_i) - \bar{y}_p\|] \leq \\
&\leq \varepsilon \sum_{p=0}^k \sum_{t_p \leq \tau_i < t_{p+1}} 2\lambda \varepsilon M(d+1)(\tau_i - t_p) = 2\varepsilon^2 \lambda M(d+1) \sum_{p=0}^k \sum_{t_p \leq \tau_i < t_{p+1}} (\tau_i - t_p) \leq \\
&\leq 2\varepsilon^2 \lambda M(d+1) \cdot \frac{L}{\varepsilon m} \sum_{p=0}^k \sum_{t_p \leq \tau_i < t_{p+1}} 1 \leq \frac{2\varepsilon \lambda M L (d+1)}{m} \cdot d \cdot \frac{L}{\varepsilon m} \cdot (k+1) \leq \frac{2\lambda M L^2 d(d+1)}{m}, \\
&\varepsilon \sum_{p=0}^k \int_{t_p}^{t_{p+1}} h(\bar{F}(\bar{X}_p, \bar{y}_p), \bar{F}(\bar{X}(s), \bar{y}(s))) ds \leq \\
&\leq \varepsilon \sum_{p=0}^k \int_{t_p}^{t_{p+1}} \lambda [h(\bar{X}_p, \bar{X}(s)) + \|\bar{y}_p - \bar{y}(s)\|] ds \leq \frac{\lambda M L^2 (d+1)}{m}, \\
&\varepsilon \sum_{p=0}^k \sum_{t_p \leq \sigma_j < t_{p+1}} h(\bar{I}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j)), \bar{I}_j(\bar{X}_p, \bar{y}_p)) \leq \frac{2\lambda M L^2 d(d+1)}{m}.
\end{aligned}$$

Similarly

$$\begin{aligned}
&\varepsilon \sum_{p=0}^k \int_{t_p}^{t_{p+1}} \|g(s, \bar{X}(s), \bar{y}(s)) - g(s, \bar{X}_p, \bar{y}_p)\| ds \leq \frac{\lambda M L^2 (d+1)}{m}, \\
&\varepsilon \sum_{p=0}^k \sum_{t_p \leq \tau_i < t_{p+1}} \|J_i(\bar{X}(\tau_i), \bar{y}(\tau_i)) - J_i(\bar{X}_p, \bar{y}_p)\| \leq \frac{2\lambda M L^2 d(d+1)}{m}, \\
&\varepsilon \sum_{p=0}^k \int_{t_p}^{t_{p+1}} \|\bar{g}(\bar{X}_p, \bar{y}_p) - \bar{g}(\bar{X}(s), \bar{y}(s))\| ds \leq \frac{\lambda M L^2 (d+1)}{m}, \\
&\varepsilon \sum_{p=0}^k \sum_{t_p \leq \sigma_j < t_{p+1}} \|\bar{J}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j)) - \bar{J}_j(\bar{X}_p, \bar{y}_p)\| \leq \frac{2\lambda M L^2 d(d+1)}{m}.
\end{aligned}$$

Using condition 3) of the theorem there exist such monotone decreasing functions $f_1(t)$ and $f_2(t)$ that tend to zero as $t \rightarrow \infty$, that for all $(X, y) \in D_1 \times D_2$ we have:

$$h \left(\int_0^t F(s, \bar{X}, \bar{y}) ds + \sum_{0 \leq \tau_i < t} I_i(\bar{X}, \bar{y}), \int_0^t \bar{F}(s, \bar{X}, \bar{y}) ds + \sum_{0 \leq \sigma_j < t} \bar{I}_j(\bar{X}, \bar{y}) \right) \leq t \cdot f_1(t),$$

$$\left\| \int_0^t (g(s, \bar{X}, \bar{y}) - \bar{g}(s, \bar{X}, \bar{y})) ds + \sum_{0 \leq \tau_i < t} J_i(\bar{X}, \bar{y}) - \sum_{0 \leq \sigma_j < t} \bar{J}_j(\bar{X}, \bar{y}) \right\| \leq t \cdot f_2(t).$$

Then

$$\begin{aligned} & \varepsilon h \left(\int_{t_p}^{t_{p+1}} F(s, \bar{X}_p, \bar{y}_p) ds + \sum_{t_p \leq \tau_i < t_{p+1}} I_i(\bar{X}_p, \bar{y}_p), \int_{t_p}^{t_{p+1}} \bar{F}(s, \bar{X}_p, \bar{y}_p) ds + \sum_{t_p \leq \sigma_j < t_{p+1}} \bar{I}_j(\bar{X}_p, \bar{y}_p) \right) = \\ & = \varepsilon h \left(\int_0^{t_{p+1}} F(s, \bar{X}_p, \bar{y}_p) ds - \int_0^{t_p} F(s, \bar{X}_p, \bar{y}_p) ds + \sum_{0 \leq \tau_i < t_{p+1}} I_i(\bar{X}_p, \bar{y}_p) - \sum_{0 \leq \tau_i < t_p} I_i(\bar{X}_p, \bar{y}_p), \right. \\ & \quad \left. \int_0^{t_{p+1}} \bar{F}(s, \bar{X}_p, \bar{y}_p) ds - \int_0^{t_p} \bar{F}(s, \bar{X}_p, \bar{y}_p) ds + \sum_{0 \leq \sigma_j < t_{p+1}} \bar{I}_j(\bar{X}_p, \bar{y}_p) - \sum_{0 \leq \sigma_j < t_p} \bar{I}_j(\bar{X}_p, \bar{y}_p) \right) \leq \\ & \leq \varepsilon \left[h \left(\int_0^{t_{p+1}} F(s, \bar{X}_p, \bar{y}_p) ds + \sum_{0 \leq \tau_i < t_{p+1}} I_i(\bar{X}_p, \bar{y}_p), \int_0^{t_{p+1}} \bar{F}(s, \bar{X}_p, \bar{y}_p) ds + \sum_{0 \leq \sigma_j < t_{p+1}} \bar{I}_j(\bar{X}_p, \bar{y}_p) \right) \right. \\ & \quad \left. + h \left(\int_0^{t_p} F(s, \bar{X}_p, \bar{y}_p) ds + \sum_{0 \leq \tau_i < t_p} I_i(\bar{X}_p, \bar{y}_p), \int_0^{t_p} \bar{F}(s, \bar{X}_p, \bar{y}_p) ds + \sum_{0 \leq \sigma_j < t_p} \bar{I}_j(\bar{X}_p, \bar{y}_p) \right) \right] \leq \\ & \leq \varepsilon [t_{p+1} \cdot f_1(t_{p+1}) + t_p \cdot f_1(t_p)] \leq 2 \sup_{\tau \in [0, L]} \tau f_1\left(\frac{\tau}{\varepsilon}\right) = \gamma_1(\varepsilon), \end{aligned}$$

where $\tau = \varepsilon t$, $\lim_{\varepsilon \rightarrow 0} \gamma_1(\varepsilon) = 0$. Similarly

$$\varepsilon \left\| \int_{t_p}^{t_{p+1}} [g(s, \bar{X}_p, \bar{y}_p) - \bar{g}(s, \bar{X}_p, \bar{y}_p)] ds + \sum_{t_p \leq \tau_i < t_{p+1}} J_i(\bar{X}_p, \bar{y}_p) - \sum_{t_p \leq \sigma_j < t_{p+1}} \bar{J}_j(\bar{X}_p, \bar{y}_p) \right\| \leq \gamma_2(\varepsilon),$$

where $\lim_{\varepsilon \rightarrow 0} \gamma_2(\varepsilon) = 0$;

$$\begin{aligned} & h \left(\int_{t_k}^t F(s, \bar{X}_k, \bar{y}_k) ds + \sum_{t_k \leq \tau_i < t} I_i(\bar{X}_k, \bar{y}_k), \int_{t_k}^t \bar{F}(s, \bar{X}_k, \bar{y}_k) ds + \sum_{t_k \leq \sigma_j < t} \bar{I}_j(\bar{X}_k, \bar{y}_k) \right) \leq \gamma_1(\varepsilon), \\ & \varepsilon \left\| \int_{t_k}^t (g(s, \bar{X}_k, \bar{y}_k) - \bar{g}(s, \bar{X}_k, \bar{y}_k)) ds + \sum_{t_k \leq \tau_i < t} J_i(\bar{X}_k, \bar{y}_k) - \sum_{t_k \leq \sigma_j < t} \bar{J}_j(\bar{X}_k, \bar{y}_k) \right\| \leq \gamma_2(\varepsilon). \end{aligned}$$

So

$$\begin{aligned} & \varepsilon h \left(\int_0^t F(s, \bar{X}(s), \bar{y}(s)) ds + \sum_{0 \leq \tau_i < t} I_i(\bar{X}(\tau_i), \bar{y}(\tau_i)), \right. \\ & \quad \left. \int_0^t \bar{F}(s, \bar{X}(s), \bar{y}(s)) ds + \sum_{0 \leq \sigma_j < t} \bar{I}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j)) \right) \leq \end{aligned}$$

$$\begin{aligned} &\leq \frac{\lambda ML^2(d+1)}{m} + \frac{4\lambda ML^2}{m}d(d+1) + \frac{\lambda ML^2(d+1)}{m} + (k+1)\gamma_1(\varepsilon) \leq \\ &\leq \frac{2\lambda ML^2}{m}(2d+1)(d+1) + m\gamma_1(\varepsilon) \equiv \varphi_1(\varepsilon, m), \end{aligned} \quad (11)$$

$$\begin{aligned} &\varepsilon \left\| \int_0^t [g(s, \bar{X}(s), \bar{y}(s)) - \bar{g}(s, \bar{X}(s), \bar{y}(s))] ds + \right. \\ &\quad \left. + \sum_{0 \leq \tau_i < t} J_i(\bar{X}(\tau_i), \bar{y}(\tau_i)) - \sum_{0 \leq \sigma_j < t} \bar{J}_j(\bar{X}(\sigma_j), \bar{y}(\sigma_j)) \right\| \leq \\ &\leq \frac{\lambda ML^2(d+1)}{m} + \frac{4\lambda ML^2}{m}d(d+1) + \frac{\lambda ML^2(d+1)}{m} + (k+1)\gamma_2(\varepsilon) \leq \\ &\leq \frac{2\lambda ML^2}{m}(2d+1)(d+1) + m\gamma_2(\varepsilon) \equiv \varphi_2(\varepsilon, m). \end{aligned} \quad (12)$$

If we substitute (11) in (9) and (12) in (10), we will get

$$\begin{aligned} h(X(t), \bar{X}(t)) &\leq \varepsilon \lambda \int_0^t [h(X(s), \bar{X}(s)) + \|y(s) - \bar{y}(s)\|] ds + \\ &+ \varepsilon \lambda \sum_{0 \leq \tau_i < t} [h(X(\tau_i), \bar{X}(\tau_i)) + \|y(\tau_i) - \bar{y}(\tau_i)\|] + \varphi_1(\varepsilon, m), \\ \|y(t) - \bar{y}(t)\| &\leq \varepsilon \lambda \int_0^t [h(X(s), \bar{X}(s)) + \|y(s) - \bar{y}(s)\|] ds + \\ &+ \varepsilon \lambda \sum_{0 \leq \tau_i < t} [h(X(\tau_i), \bar{X}(\tau_i)) + \|y(\tau_i) - \bar{y}(\tau_i)\|] + \varphi_2(\varepsilon, m). \end{aligned}$$

Adding these two inequalities and applying the analogue of Gronwall—Bellmann lemma [14] we get

$$\begin{aligned} h(X(t), \bar{X}(t)) + \|y(t) - \bar{y}(t)\| &\leq (\varphi_1(\varepsilon, m) + \varphi_2(\varepsilon, m)) (1 + \varepsilon \lambda)^{i(0,t)} e^{\varepsilon \lambda t} \leq \\ &\leq (\varphi_1(\varepsilon, m) + \varphi_2(\varepsilon, m)) (1 + \varepsilon \lambda)^{d \cdot t} e^{\varepsilon \lambda t} \leq \\ &\leq (\varphi_1(\varepsilon, m) + \varphi_2(\varepsilon, m)) (1 + \varepsilon \lambda)^{d \cdot \frac{t}{\varepsilon}} e^{\lambda L} \leq \\ &\leq (\varphi_1(\varepsilon, m) + \varphi_2(\varepsilon, m)) e^{(d+1)\lambda L} = \\ &= \left(\frac{4\lambda ML^2}{m}(2d+1)(d+1) + m\gamma_1(\varepsilon) + m\gamma_2(\varepsilon) \right) e^{(d+1)\lambda L}. \end{aligned}$$

Then for every summand the inequality holds:

$$\begin{aligned} h(X(t), \bar{X}(t)) &\leq \left(\frac{4\lambda ML^2}{m}(2d+1)(d+1) + m\gamma_1(\varepsilon) + m\gamma_2(\varepsilon) \right) e^{(d+1)\lambda L}, \\ \|y(t) - \bar{y}(t)\| &\leq \left(\frac{4\lambda ML^2}{m}(2d+1)(d+1) + m\gamma_1(\varepsilon) + m\gamma_2(\varepsilon) \right) e^{(d+1)\lambda L}. \end{aligned}$$

Let $\eta_1 = \min\{\rho, \eta, \xi\}$. Choose m to satisfy the inequality

$$e^{(d+1)\lambda L} \frac{\lambda ML^2}{m} (2d+1)(d+1) < \frac{\eta_1}{12}.$$

Then fix m and choose $\varepsilon_0 \in (0, \bar{\varepsilon}]$ such that for $\varepsilon \in (0, \varepsilon_0]$ the inequalities hold

$$e^{(d+1)\lambda L} m\gamma_1(\varepsilon) \leq \frac{\eta_1}{3}, e^{(d+1)\lambda L} m\gamma_2(\varepsilon) \leq \frac{\eta_1}{3}.$$

Then $h(X(t), \bar{X}(t)) \leq \eta_1$ and $\|y(t) - \bar{y}(t)\| \leq \eta_1$ if the solution $(X(t), y(t))$ belongs to the domain $D_1 \times D_2$. And it follows from condition 3) of the theorem as $\eta_1 = \min\{\eta, \rho, \xi\}$.

So, we get that for any $\eta > 0$ and $L > 0$ there exists such $\varepsilon_0 \in (0, \bar{\varepsilon}]$ that for $\varepsilon \in (0, \varepsilon_0]$ and $t \in [0, L\varepsilon^{-1}]$ the following inequalities fulfill

$$h(X(t), \bar{X}(t)) \leq \eta, \|y(t) - \bar{y}(t)\| \leq \eta,$$

where $(X(t), y(t))$ and $(\bar{X}(t), \bar{y}(t))$ are the solutions of systems (2), (3) and (4), (5) with the initial conditions $X(0) = \bar{X}(0) \in D'_1$, $y(0) = \bar{y}(0) \in D'_2$.

The theorem is proved.

Example 1. Consider the impulsive hybrid system

$$X \in \text{conv}(R^2), y \in R.$$

$$\begin{cases} D_H X = \varepsilon \left[\begin{pmatrix} 1 & \sin t \\ \cos 2t & 2y^2 \end{pmatrix} X + \Pi_{2+e^{-t}+\sin(\ln(1+t)),1} \begin{pmatrix} \sin t \\ \cos t \end{pmatrix} \right], \\ X_0 = S_{0.5} \begin{pmatrix} -1 \\ 0 \end{pmatrix}, \\ \dot{y} = \varepsilon \frac{y}{|X|} \sin^2 t, y_0 = 0.1. \end{cases}$$

$$\Delta X|_{t=\tau_i} = \varepsilon X, \Delta y|_{t=\tau_i} = -\varepsilon y, \tau_i = 2\pi i, i = \overline{1, \infty}.$$

The averaged system is:

$$\begin{cases} D_H \bar{X} = \varepsilon \left[\begin{pmatrix} 1 & 0 \\ 0 & 2\bar{y}^2 \end{pmatrix} \bar{X} + \Pi_{2+\sin(\ln(1+t)),1} \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right], \\ \bar{X}_0 = S_{0.5} \begin{pmatrix} -1 \\ 0 \end{pmatrix}, \\ \dot{\bar{y}} = \varepsilon \frac{\bar{y}}{2|\bar{X}|}, \bar{y}_0 = 0.1. \end{cases}$$

$$\Delta \bar{X}|_{t=\tau_j} = \varepsilon \bar{X}, \Delta \bar{y}|_{t=\tau_j} = -\varepsilon \bar{y}, \tau_j = 2\pi j, j = \overline{1, \infty}.$$

The graphs of the solutions of initial system and averaged system see on next page.

CONCLUSION. This paper contains the substantiation of the scheme of partial averaging for one class of impulsive hybrid systems where one equation is a differential equation with Hukuhara derivative and the other one is an ordinary differential equation. In case when the right-hand sides are periodic in time one can obtain a better estimate. Namely one can show that for any $L > 0$ there exist $C(L) > 0$ and $\varepsilon_0(L) > 0$ such that the conclusion of the theorem holds with $\eta = C\varepsilon$.

a) When $\varepsilon = 0.1$:




Image 1. (a) The graph of the solutions of initial system $X(t)$ (black) and averaged system $\bar{X}(t)$ (gray). (b) The graph of the solutions of initial system $y(t)$ (black) and averaged system $\bar{y}(t)$ (gray).

b) When $\varepsilon = 0.05$:




Image 2. (a) The graph of the solutions of initial system $X(t)$ (black) and averaged system $\bar{X}(t)$ (gray). (b) The graph of the solutions of initial system $y(t)$ (black) and averaged system $\bar{y}(t)$ (gray).

1. **de Blasi F. S.** On the differentiability of multifunctions / F. S. de Blasi // Pacific J. Math. – 1976. – Vol. 66, Is. 1. – P. 67 – 81.
2. **de Blasi F. S.** Equazioni differenziali con soluzioni a valore compatto convesso / F. S. de Blasi, F. Iervolino // Boll. Unione Mat. Ital. – 1969. – Vol. 2, Is. 4 – 5. – P. 491 – 501.

3. **de Blasi F. S.** Euler method for differential equations with set-valued solutions / F. S. de Blasi, F. Iervolino // *Boll. Unione Mat. Ital.* – 1971. – Vol. 4, Is. 4. – P. 941 – 949.
4. **Brandao Lopes Pinto A. J.** Uniqueness and existence theorems for differential equations with compact convex valued solutions / A. J. Brandao Lopes Pinto, F. S. de Blasi, F. Iervolino // *Boll. Unione Mat. Ital.* – 1970. – Is. 4. – P. 534 – 538.
5. **Hukuhara M.** Integration des applications mesurables dont la valeur est un compact convexe / M. Hukuhara // *Func. Ekvacioj.* – 1967. – Is. 10. – P. 205 – 223.
6. **Skripnik N. V.** Averaging of impulsive differential equations with Hukuhara derivative. / N. V. Skripnik // *Visn. Yuriy Fedkovich Chernivtsy National University.* – 2008. – Vol. 374. – P. 109 – 115. (in Ukrainian)
7. **Differential** equations with impulse effects: multivalued right-hand sides with discontinuities (De Gruyter Studies in Mathematics: 40) / [Perestyuk N. A., Plotnikov V. A., Samoilenko A. M., Skripnik N. V.]. – Berlin/Boston: Walter De Gruyter GmbH Co., 2011. – 307 p.
8. **Plotnikov V. A.** Differential equations with a multivalued right-hand side. Asymptotic methods / V. A. Plotnikov, A. V. Plotnikov, A. N. Vityuk. – Odessa: Astroprint, 1999. – 354 p. (in Russian)
9. **Kisielewicz M.** Method of Averaging for Differential Equations with Compact Convex Valued Solutions / M. Kisielewicz // *Rend. Math.* – 1976. – Vol. 9, Is. 3. – P. 397–408.
10. **Plotnikov V. A.** Existence, continuous dependence and averaging in differential equations with Hukuhara derivative and delay / V. A. Plotnikov, P. I. Rashkov // "Mathematics and education in mathematics": Proceedings of Twenty Sixth Spring Conference of the Union of Bulgarian Mathematicians (April 22–25, 1997). – Plovdiv, Bulgaria, 1997. – P. 179–184
11. **Plotnikov V. A.** Averaging in differential equations with Hukuhara derivative and delay / V. A. Plotnikov, P. I. Rashkov // *Funct. Differ. Equ.* – 2001. – Vol. 8. – P. 371 – 381.
12. **Plotnikov V. A.** Averaging of controlled equations with the Hukuhara derivative / V. A. Plotnikov, O. D. Kichmarenko // *Nonlinear Oscil. (N. Y.).* – 2006. – Vol. 9: 3. – P. 365 – 374.
13. **Plotnikov V. A.** Averaging of equations with Hukuhara derivative, multivalued control and delay / V. A. Plotnikov, O. D. Kichmarenko // *Bulletin of the Odessa National University.* – 2007. – Vol. 12: 7. – P. 130 – 139. (in Russian)
14. **Samoilenko A. M.** Impulsive Differential Equations / A. M. Samoilenko, N. A. Perestyuk. – Singapore: World Scientific, 1995. – 462 p.