

tions are in accordance with the species biological requirements. And on the contrary, incompatibility between the natural conditions and the requirements of living organisms leads to low yield capacity crop production.

The aim of the article is to study the impact of climatic factors in Lubny district of Poltava region on winter wheat yield capacity as winter wheat is the main grain crop of the area.

In the research there was performed the analysis of long-term (1996-2015) data concerning winter wheat yield rates and the meteorological parameters, which characterize the temperature and moisture regimes in Lubny district of Poltava region. The study showed the interrelation between climate conditions and winter wheat productivity. Using the data collected by Lubny meteorological station we calculated the classical index of heat supply in the region during the period of 1996-2015. In the course of the research the analytical methods, long-term statistical analysis, methods of comparison and logical generalization were applied.

The analysis shows that the unevenness of the dynamic changes in winter wheat yield capacity is determined by the interrelation of the climatic territorial conditions, which considerably influence the performance of all the components in agroecosystems. Taking into account the significant impact of the agro-climatic factors on the crop plantings, plant heat supply index, moisture regime, crop damage with diseases, as well as weed contamination index have been analyzed over a 20-year period and showed their influence on winter wheat productivity have been shown.

Analyzing the dynamics of winter wheat capacity depending on climatic changes within a 20-year period, it has been found that it ranged from 0.8 t/ha to 4.9 t/ha. Whereas in 1997, 2000, 2003 the yield capacity dropped to 1.8, 1.0 and 0.8 t/ha, respectively, and that was presumably caused by the adverse weather behaviour during the above-mentioned years. In particular, the winter season in these years was characterized by the unstable changes in certain climatic parameters. As a result, there was observed the formation of thick ice crust. The reduction of the crop plantings comprised 25-32% in the early spring of 2000, which led to the increase in weed contamination. At the same time, heavy rains in June with strong winds and hail caused crop lodging.

Taking into consideration the low winter wheat yields in 1997, 2000, and 2003 it was necessary to analyze the disease infestation and weed contamination indexes. The statistical data showed that weed contamination of winter wheat comprised 2-3 points out of 3. Moreover, the unfavourable weather conditions of the summer season in 1997 caused a fungal disease (*Septoria-anodorum*).

The analysis of data which embraced a 20-year period (1996-2015) showed a tendency to the increase in classical index of heat supply and the reduction of annual precipitation compared to the long-term norms, which influenced the winter wheat productivity.

Further changes in climate parameters may cause the changes in the weather conditions in the cultivation area and lead to considerable reduction in the crop production. Therefore, to obtain permanent high winter wheat yields in the forest-steppe zone it is necessary to take into account the climatic resources of the region.

Key words: winter wheat, crop productivity, heat supply index, moisture content index, weed contamination, agro-climatic resources.

Надійшла 08.04.2016 р.

UDK 633.111"324":575.1/2:631.527.5

LOZINSKYI M., candidate of agricultural science
Bila Tserkva National Agrarian University

INHERITANCE AND GRAIN WEIGHT TRANSGRESSIVE VARIABILITY PER PLANT IN HYBRID WINTER WHEAT (*T. AESTIVUM L.*), OBTAINED FROM THE HYBRIDIZATION OF VARIOUS ECOTYPES

Висвітлено особливості успадкування маси зерна з рослини у гібридів F_1 пшениці м'якої озимої. Встановлено, що успадкування маси зерна з рослини гібридами першого покоління в переважній більшості комбінацій проходило за типом позитивного наддомінування. Ступінь фенотипового домінування (h_p) становив 1,4-64,0. Істинний гетерозис за масою зерна з рослини спостерігався у дев'яти з десяти гібридів F_1 з показником 41,1-68,9 %. Частота позитивних трансгресивних рекомбінантів за масою зерна з рослини у гібридів F_2 , отриманих від схрещування степового еко типу з лісостеповим становила 36,0-80,2 %. У гібридних популяцій, отриманих від схрещування віддалених еколого-географічних форм, найбільша кількість позитивних трансгресій спостерігалась в популяціях Гайтун / Олеся і Гайтун / Білоцерківська напівкарликова – 68,8 і 57,1 % відповідно.

Ключові слова: пшениця м'яка озима, маса зерна з рослини, еко типи, комбінації схрещування, гібриди, успадкування, гетерозис, ступінь домінування, ступінь і частота трансгресій.

Introduction. One of the priorities of agriculture of Ukraine is a significant increase and stabilization of grain production. Soft wheat winter is the basic and the most important food crop in the world. It is grown in most countries.

The main objective in wheat breeding is to create a soft winter varieties with high productivity. In recent years, due to global climate change, much attention is paid to breeding varieties with enhanced adaptive capacity [1, 2, 3].

Analysis of domestic and international breeding results reveals that broad scientific and reasonable use in breeding programs of different source material is crucial in breeding new modern varieties. Therefore, the study of winter wheat sort samples collection for economically valuable traits provides establishing their breeding value creation for further selecting the varieties with high productivity, grain quality and adaptability to specific soil and climatic cultivation conditions.

Of particular urgency is the experimental verification of the properties of new winter wheat sort samples of various ecological-geographical and genetic origin. Their involvement in hybridization contributes to identifying their breeding value under certain soil and climatic conditions for breeding varieties with agronomic traits complex.

Analysis of recent research and publications. The weight of grain per plant is the main feature of the structure of individual grain productivity of wheat and, according to most scholars, it is one of the most effective means to improve the crop performance [4, 5, 6]. The level of the manifestation of the symptoms depends on many elements, each of them has its own inheritance character and variation range. Number of plants per area unit and their performance determine the level of productivity.

The most effective method of winter wheat selection is intraspecific genetic recombination characteristics with applying different types of crossbreeding with the following single and multiple selection [7]. Gene D, which provided hexaploid wheat tetraploid two-genome transition from level to three-genome in combination with the first two genomes Au and B resulted in large intraspecific polymorphism and diversity of *T. aestivum* L ecotypes, which makes it possible for its improvement [4, 8]. As a result of crossing varieties that differ in morphological, biological and physiological characteristics, with different levels of genetic potential productivity and resistance to biotic and abiotic adverse environmental factors, large quantities of genetically modified form of several parents united in a single genotype. This diversity of recombinant is a starting material for the subsequent creation of a new genotype closely related to environmental conditions [9]. Successful selection of the crossbreed pairs requires determining the direction of selection and studying the agri-environmental conditions for the planned cultivation areas, taking into account factors that may limit the potential yield of the variety.

Varieties of different ecotypes are adapted to the conditions of ecological and climatic zones for which they are derived. They differ in terms of duration of the growing season, plant height, morphological features, endurance to adverse environmental conditions. Ye.M. Sins'ka launched the doctrine of the development of the ecological system of selection [10]. The current ecological and geographical differentiation of plant genetic resources provides limitless opportunities of their use in producing varieties with desirable biological properties.

The probability of positive breeding in respect of transgressions increases in crossing the varieties which belong to different ecological types of development. The essence of this phenomenon is not geographical distance and differences in genetic hybridization components which are caused by the results of selection in different natural and historical conditions. The presence of genetic differences in the components of hybridization is the main condition for the recombinant alleles system, which provides the best possible expression of quantitative traits of wheat productivity [4].

Knowledge on the genetic nature of quantitative traits determining plants productivity are of great importance for breeding work [11, 12]. M.I. Vavilov explained insufficient study of quantitative traits by their complexity, the presence of transitional forms, genetic determination of traits ignorance [13].

The aim of research was to establish the nature of inheritance of grain weight per plant by F1 hybrids and the extent and frequency of positive transgressions in F2 hybrid populations of soft winter wheat, obtained by crossing parent forms belonging to different ecological groups.

Research material and methods. The study was conducted in Bila Tserkva experimental breeding station (BTEBS) of the Institute of crops bioenergy and sugar beet in 2011-2013.

Parental forms were the varieties of breeding establishments located in different ecological and geographical areas, namely: Missia Odes'ka (Miss. Od.) (Plant Breeding and Genetics Institute), Vidrada, Lybid', Olesya, Rostavytsya, Bilotserkivska semi-dwarf (BTEBS), Dryada 1 (ETE "Dryada a"), Polis'ka 90 (Institute of agriculture), NAZ (Kazakhstan), Haytun and Pekin (China), belonging to various environmental groups. We have studied 10 hybrid combinations: Miss. Od. / Vidrada, Miss. Od. / Lybid, Dryada a 1 / Olesya, Dryada 1 / Rostavytsya, NAZ / Olesya, NAZ / Polis'ka 90 Haytun / Olesya, Haytun / B.TS. s / d, Pekin / Olesya, Pekin / B.TS. s / d. Seeds F1-2 were sown with breeding drill SSKF-7M on the scheme: parent form, hybrid, parent form. Hybrid generation was worked by pedihri method. During

the growing season we conducted phenological observations and structural analysis of sheaves was done after the complete ripeness [14-15].

The degree of phenotypic dominance (h_p) of grain weight per plant in the hybrids, was determined according to the formula by G.M. Bailey and R.I. Atkins [16], the degree and frequency of positive transgressions – according to the formulas suggested by H.S. Voskresenskaya, V.I. Shpot [17], A.P. Orlyuk and V.V. Bazaliy [18], true heterosis according to the formula proposed by Kh. Daskalev [19].

Biometric analyzes was carried out on an average sample of 25 plants in triple repetition. The results of the experimental data were treated statistically by the "Statistica" program, version 5.0.

Results and discussion. It has been found out that the inheritance of grain mass per plant hybrids of the first generation of winter wheat soft in most combinations held by the type of positive overdominance. In crossing the steppe and forest-steppe ecotypes the degree of dominance was within 1.4-64.0. In five of the six hybrids obtained by hybridization of remote ecological and geographical forms the rate of phenotypic forms of domination was within 5.9-64.0. Only in the combination of crossing NAZ / Polis'ka 90 the determination of the trait "grain weight per plant" developed on the negative dominance type (Fig. 1).




Fig. 1. The degree of phenotypic domination of grain weight in soft winter wheat F1 hybrids (2012).

Crossing combinations: 1 – Miss. Od. / Vidrada; 2 – Miss. Od. / Lybid'; 3 – Dryada 1 / Olesya; 4 – Dryada 1 / Rostavytsya; 5 – NAZ / Olesya; 6 – NAZ / Polis'ka 90; 7 – Haytun / Olesya; 8 – Haytun / B.TS. s/d; 9 – Pekin / Olesya; 10 – Pekin / B.TS. s/d.

The index of phenotypic dominance does not give grounds to judge the value of heterosis effect – it only determines the nature of the manifestation of the trait studied: its values are essential only within 1.1 – (1.1). A more objective assessment of trait inheritance characteristics can be obtained by calculating the degree of heterosis [20].

The degree of heterosis (H%) was determined by comparing the rate of grain weight in F1 hybrid plants with this indicator in a better parent form.

True heterosis for grain weight per plant was marked in nine of the ten first-generation hybrids. In hybrids Dryada 1 / Rostavytsya (crossing steppe and forest steppe ecotypes) and the NAZ / Olesya (crossing remote eco-geographical forms) we observed the highest heterosis rates of 68.9 and 60.9 %, respectively (Fig. 2).

It has been found out that nine out of the ten first-generation hybrids with a grain weight per plant of 5.85-9.06 g exceeded the rates of the parental forms with a bigger display by 0.46-3.28 g (Table. 1).

Hybrids F2 exceeded the parental forms with bigger trait display by 0.19-2.68 g for grain weight per plant, except for Miss. Od. / Lybid. The maximum weight of grain per plants in the F2 hybrid populations goexceeded the parental forms and reached 4.82-8.30 g. Variation of trait in F1-2 hybrids and parental forms (except for Haytun, 2013) is significant.




Fig. 2. Heterosis for grain weight per plant in the F1 soft winter wheat hybrids (2012).
Crossing combinations: 1 – Miss. Od. / Vidrada; 2 – Miss. Od. / Lybid’; 3 – Dryada I / Olesya;
 4 – Dryada I / Rostavytsya; 5 – NAZ / Olesya; 6 – NAZ / Polis’ka 90; 7 – Haytun / Olesya;
 8 – Haytun / B.TS. s/d; 9 – Pekin / Olesya; 10 – Pekin / B.TS. s/d.

An integral part of studying soft winter wheat breeding is the selection of transgressive recombinants in hybrid populations both by the elements of the crop structure in the complex of traits that make up the adaptive potential of modern varieties.

Using transgressive fission, a breeder can extend the existing limits and intensity of desirable traits display in wheat. Obtaining these forms of winter wheat by certain economically valuable traits and their complex is an urgent and one of the most difficult problems of the crop breeding. By transgressive splitting it is possible to create new features or, more commonly, a new level of intensity of manifestation of existing signs [4].

Table 1 – Display of statistical indicators of weight variation in grain plants F1-2 hybrids and their parental forms

Crossing combinations and parental forms	F ₁ hybrids, 2012				F ₂ hybrids, 2013			
	$(\bar{x} \pm S \bar{x}), r$	Lim (r)		V, %	$(\bar{x} \pm S \bar{x}), r$	Lim (r)		V, %
		min	max			min	max	
Steppe ecotype / Forest-Steppe ecotype								
♀ Miss. Od.	4,52 ± 0,31	2,07	6,29	27,7	3,33 ± 0,23	2,28	4,42	21,7
Miss. Od. / Vidrada	7,04 ± 0,47	4,42	11,03	24,3	4,93 ± 0,37	2,24	8,30	31,4
♂ Vidrada	4,43 ± 0,46	1,84	9,01	36,6	2,70 ± 0,26	1,58	3,85	30,1
Miss. Od. / Lybid’	7,37 ± 0,52	2,59	12,02	23,8	2,96 ± 0,44	1,90	6,80	55,5
♂ Lybid’	4,88 ± 0,45	2,11	7,18	32,9	2,09 ± 0,15	1,37	2,72	22,9
♀ Driada I	3,08 ± 0,34	1,46	5,79	42,7	1,14 ± 0,10	0,62	1,85	35,1
Driada I / Olesya	5,85 ± 0,43	2,97	7,83	26,9	4,44 ± 0,42	2,04	7,40	31,1
♂ Olesya	5,39 ± 0,49	2,31	8,85	32,0	2,95 ± 0,26	1,42	4,15	27,5
Driada I / Rostavytsya	7,38 ± 0,50	3,93	12,22	23,6	5,33 ± 0,43	3,51	7,59	25,4
♂ Rostavytsya	4,37 ± 0,44	2,23	6,73	37,7	2,65 ± 0,36	1,00	4,24	42,9
Remote ecological geographical forms crossing								
♀ NAZ	4,59 ± 0,49	2,08	9,02	37,8	3,75 ± 0,37	2,13	5,97	31,1
NAZ / Olesya	8,67 ± 0,67	4,12	17,09	23,4	3,94 ± 0,44	2,18	6,94	38,7
NAZ / Polis’ka 90	4,60 ± 0,26	2,75	6,32	25,0	4,80 ± 0,45	2,20	7,25	32,1
♂ Polis’ka 90	4,81 ± 0,42	2,19	8,34	35,1	2,65 ± 0,26	1,41	3,58	31,1
♀ Haytun	5,97 ± 0,43	2,91	9,39	28,7	3,01 ± 0,16	2,26	3,82	17,3
Haytun / Olesya	9,06 ± 0,69	4,60	17,42	22,1	4,87 ± 0,28	3,06	6,92	23,2
Haytun / B.Ts. s/d	8,48 ± 0,53	4,64	14,0	20,9	4,85 ± 0,47	2,48	7,17	31,9
♂ B.Ts. s/d	4,95 ± 0,41	2,32	8,56	34,0	2,86 ± 0,28	1,47	4,54	31,3
♀ Pekin	5,48 ± 0,39	2,87	8,78	30,1	3,13 ± 0,26	2,12	4,45	26,2
Pekin / Olesya	8,00 ± 0,49	4,70	13,43	21,8	3,37 ± 0,33	1,52	5,77	37,9
Pekin / B.Ts. s/d	7,73 ± 0,47	4,15	10,95	21,8	3,37 ± 0,30	1,96	4,82	29,4
Podolyanka (St)	5,77 ± 0,52	2,14	10,62	30,6	2,53 ± 0,23	1,63	4,16	28,2

The degree of positive transgressions by grain weight per plant in the studied populations of the second generation hybrid was in the range of 8.3 % (Pekin x B.Ts. s/d) to 87.8 % (Miss. Od. / Vidrada) (Table 2).

Frequency of transgressive recombinants by weight of corn plants in hybrids derived from crossing steppe and forest steppe ecotypes was 36.0-80.2 %. In hybrid populations obtained by crossing remote eco-geographical forms, the highest number of transgressions was obtained in populations Haytun / Olesya and Haytun x B.TS. s / d – 68.8 and 57.1 % respectively.

Table 2 – Degree and frequency of positive transgressions by weight per plant in F2 hybrids (2013)

Crossing combinations	h_p degree in F_1	Transgression degree, %	Transgression frequency, %
Steppe ecotype / Forest-Steppe ecotype			
Miss. Od. / Vidrada	64	87,8	36,0
Miss. Od. / Lybid'	14,8	53,8	30,8
Dryada 1 / Olesya	1,4	78,3	33,3
Dryada 1 / Rostavytsya	5,7	79,0	80,2
Remote ecological geographical forms crossing			
NAZ / Olesya	9,2	16,2	8,3
NAZ / Polis'ka 90	-0,9	21,4	20,0
Haytun / Olesya	11,7	66,7	68,8
Haytun / B.TS. s/d	5,9	57,9	57,1
Pekin / Olesya	64	29,7	20,0
Pekin / B.TS. s/d	9,7	8,3	5,3

Conclusions and recommendations for further research. 1. F1 hybrids inheritance of soft winter wheat grain weight per plant, obtained by crossing parental ecotypes of different forms, developed, in most combinations, according to the type of positive overdominance. The degree of phenotypic dominance (h_p) was 1.4-64.0.

2. The highest heterosis rate for the grain weight per plant was observed in hybrids Dryad 1 / Rostavytsya (steppe and forest-steppe ecotypes crossing) and in NAZ / Olesya (Remote ecological geographical forms crossing) – 68.9 and 60.9 % respectively.

3. Frequency of transgressive recombinants by grain weight per plant in F2 hybrid populations, obtained by steppe and forest-steppe ecotypes crossing was 36.0-80.2 %, received. The highest number of transgressions in remote eco-geographical forms crossing was obtained in populations Haytun / Olesya and Haytun B.Ts. s/d – 68.8 and 57.1 % respectively.

4. Engaging local adapted varieties of other ecotypes in hybridization allows creating significant reserve genotypic variability by grain weight per plant.

The prospect of further research is selection and evaluation of the obtained recombinants by a complex of economically valuable traits aimed to create a new source material for breeding varieties with high productivity and adaptability to adverse environmental conditions.

LIST OF REFERENCES

1. За маркерними ознаками. Оцінювання стійких сортів озимої м'якої пшениці проти шкідників / Трибель С.О., Стругун О.О., Гетьман М.В., Топчій Т.В. // Насінництво. – 2010. – № 10. – С. 4-8.
2. Бурденюк-Тарасевич Л. Пшеница. Глубина генетического потенциала / Л. Бурденюк-Тарасевич // Зерно. – 2010. – № 4 (48). – С. 49-51.
3. Колюча Г.С. Природні і штучні види пшениці та амфідиплоїди як джерело генетичного різноманіття при створенні вихідного матеріалу для селекції озимої пшениці / Г.С. Колюча, В.Т. Колючий // Науково-технічний бюлетень Миронівського інституту пшениці ім. В.М. Ремесла. – Миронівка, 2009. – Вип. 9. – С. 25-32.
4. Орлюк А.П. Генетика пшениці з основами селекції: [Монографія] / А.П. Орлюк. – Херсон: Айлант, 2012. – 436 с.
5. Власенко В.А. Оцінка адаптивності сортів пшениці озимої за врожайністю та висотою рослин / В.А. Власенко, Л.А. Коломієць, Г.С. Басанець // Проблеми підвищення адаптивного потенціалу системи рослинництва у зв'язку зі змінами клімату: Тези доп. міжнар. наук.-практ. конф., 26-28 лютого 2008 року, м. Біла Церква / Білоцерківський ДАУ. – Біла Церква, 2008. – С. 16.
6. Васильківський С.П. Особливості використання хімічного мутагенезу при створенні вихідного матеріалу для селекції пшениці: автореф. дис...д-ра с.-г. наук / С.П. Васильківський. – Одеса, 1999. – 35 с.

7. Бурденюк-Тарасевич Л.А. Принципи підбору пар для гібридизації в селекції озимої пшениці *T. aestivum* L. на адаптивність до умов довкілля / Л.А. Бурденюк-Тарасевич, М.В. Лозінський // Фактори експериментальної еволюції організмів: зб. наук. пр. / Національна академія наук України, АН України, Інститут молекулярної біології і генетики, Укр. т-во генетиків і селекціонерів ім. М.І. Вавилова; редкол.: В.А. Кунах (голов. ред.) [та ін.]. – К.: Укр. т-во генетиків і селекціонерів ім. М.І. Вавилова, 2015. – Т.16. – С. 92-96.
8. Лукьяненко П.П. Гибридизация отдаленных эколого-географических форм озимой пшеницы / П.П. Лукьяненко // С.-х. биология. – М., 1968. – № 1. – С. 3-11.
9. Жученко А.А. Экологическая генетика культурных растений (адаптация, рекомбиногенез, агробиоценоз / А.А. Жученко. – Кишинев: Штиинца, 1980. – 588 с.
10. Синская Е.Н. Экологическая система селекции кормовых культур / Е.Н. Синская. – Л.: ВИР, 1933. – 44 с.
11. Лисничук Г.Н. Характер генотипической корреляции урожая зерна озимой пшеницы с элементами его структуры / Г.Н. Лисничук // Селекция и семеноводство. – М.: Колос, 1985. – № 4. – С. 16-17.
12. Адаптивная селекция. Теория и технология на современном этапе / [П.П. Литун, В.В. Кириченко, В.П. Петренкова, В.П. Коломацкая]. – Харьков, 2007. – 263 с.
13. Вавилов Н.И. Избранные сочинения / Н.И. Вавилов. – М.: Колос, 1966. – 460 с.
14. Методика державного випробування сортів рослин на придатність до поширення в Україні: Заг. част. // Охорона прав на сорти рослин: Офіційний бюл. / Гол. ред. В.В. Волкодав. – К.: Алефа, 2003. – Вип.1, ч. 3. – 106 с.
15. Доспехов Б.А. Методика полевого опыта / Б.А. Доспехов. – М.: Агропромиздат, 1985. – 352 с.
16. Beil C.M. Inheritance of quantitative characters in grain sorghum / C.M. Beil, P.E. Atkins // Jowa J. Sci., 1965. – Vol. 39. – № 3. – P. 345-358.
17. Воскресенская Г.С. Трансгрессия признаков Brassica и методика количественного учета этого явления / Г.С. Воскресенская, В.И. Шпота // Доклады ВАСХНИЛ. – М., 1967. – № 7. – С. 18-20.
18. Орлюк А.П. Принципы трансгрессивной селекции пшеницы / А.П. Орлюк, В.В. Базалий. – Херсон, 1998. – 274 с.
19. Даскалев Хр. Гетерозис при доминанте / Хр. Даскалев, М. Йорданом, А. Огнянова. – София: Българска академия на науките, 1967. – 179 с.
20. Аналіз успадкування деяких кількісних ознак гороху посівного / Мамалига В.С., Кондратенко М.І., Бугайов В.Д., Янчук В.І. // Фактори експериментальної еволюції організмів: Зб. наук. пр. / НАН України, НААН України, НАМН України, Інститут молекулярної біології і генетики НАН України, Укр. т-во генетиків і селекціонерів ім. М.І. Вавилова; редкол.: В.А. Кунах (голов. ред.) [та ін.]. – К.: Логос, 2013. – С. 214-219.

REFERENCES

1. Evaluation of winter wheat varieties resistance against pests by the marker signs / S.O. Triybel', O.O. Struhun, M.V. Get'man, T.V. Topchiy // Seed breeding. – 2010. – № 10. – P. 4-8.
2. Burdeynyuk-Tarasevych L. Wheat. Genetic capacity depth / L. Burdeynyuk-Tarasevych // Cereals. – 2010. – № 4 (48). – P. 49-51.
3. Kolyucha G.S. Natural and artificial types of wheat and amphidybrids as a source of genetic diversity in creating original material for winter wheat breeding / G.S. Kolyucha, V.T. Kolyuchiy // Scientific and Technical Bulletin of Myronivka Institute of wheat named after V.N. Remeslo. – Myronivka, 2009. – Vol. 9. – P. 25-32.
4. Orlyuk A.P. Wheat genetics with breeding fundamentals [Manual] / A.P. Orlyuk. – Kherson: Ailant, 2012. – 436 p.
5. Vlasenko V.A. Evaluation of winter wheat varieties adaptability by the yield and plant height / V.A. Vlasenko, L.A. Kolomiets', G.S. Basanets' // Problems of crops breeding adaptive capacity increasing in terms of climate change: Theses of rep. of intern. science practical. conf., 26-28 February 2008, Bila Tserkva / Bila Tserkva State Agrarian University. – Bila Tserkva, 2008. – P. 16.
6. Vasykivskiy S.P. Some peculiarities of using chemical mutagenesis in creating original material wheat for breeding: Author. dis ... for Dr of agricultural sciences / S.P. Vasykivskiy. – Odessa, 1999. – 35 p.
7. Burdeynyuk-Tarasevych L.A. Principles of pairs selection for hybridization in *T. aestivum* L. winter wheat breeding for the adaptability to environmental conditions / L.A. Burdeynyuk-Tarasevych, M.V. Lozinskyi // Factors of experimental evolution of organisms: Bulletin. / National Academy of Sciences of Ukraine, Academy of Sciences of Ukraine, Institute of Molecular Biology and Genetics, Ukr. society of geneticists and breeders named after M.I Vavilov; Ed.: V.A. Kunaх (ed.-in-chief) et al. – K.: Ukr. society of geneticists and breeders named after M.I Vavilov, 2015. – Vol.16. – P. 92-96.
8. Luk'yanenko P.P. Hybridization of remote eco-geographical forms of winter wheat / P.P. Luk'yanenko // Agr. biology. – М., 1968. – № 1. – P. 3-11.
9. Zhuchenko A.A. Environmental genetics of cultural plants (Adaptation, recombination, agrobiocenosis / A.A. Zhuchenko. – Kishinev: Shtyynsa, 1980. – 588 p.
10. Sinskaya Ye.N. Environmental system of crops breeding selection / Ye.N. Sinskaya. – L.: VIR, 1933. – 44 p.
11. Lysnychuk G.N. The nature of genotype correlation of winter wheat grain yield with the element of its structure / G.N. Lysnychuk // Selection and seeds breeding. – М.: Kolos, 1985. – № 4. – P. 16-17.
12. Adaptive selective breeding. Theory and current technology / [P.P. Litun, V.V. Kirichenko, V.P. Petrenkova, V.P. Kolomatskaya]. – Kharkiv, 2007. – 263 p.
13. Vavilov N.I. Selected papers / N.I. Vavilov. – М.: Kolos, 1966. – 460 p.
14. Methods of state plants varieties for testing suitability for distribution in Ukraine: Gen. part // Protection of rights for crops varieties: Official Bull. / Ed.-in-chief V.V. Volkodav. – K.: Alepha, 2003. – Vol.1, part 3. – 106 p.
15. Dospekhov B.A. Methods of field experience / B.A. Dospekhov. – М.: Агропромиздат, 1985. – 352 p.
16. Beil C.M. Inheritance of quantitative characters in grain sorghum / C.M. Beil, P.E. Atkins // Jowa J. Sci., 1965. – Vol. 39. – № 3. – P. 345-358.

17. Voskresenskaya G.S. Transgression of Brassica traits and accounting methodology of the phenomenon / G.S. Voskresenskaya, V.I. Shpota // VASHNYL thesis. – M., 1967. – № 7. – С. 18-20.

18. Orlyuk A.P. Principles wheat transgressive selection / A.P. Orlyuk, V.V. Bazaliy. – Kherson, 1998. – 274 p.

19. Daskalev Chr. Heterosis at domatyte / Chr. Daskalev, M. Yordanom A. Ohnyanova. – Sofia: Bulgaria academy of science, 1967. – 179 p.

20. The analysis of inheritance of some pea quantitative traits / Mamalyha V.S., Kondratenko M.I., Bugayov V.D., Yanchuk V.I. // Factors of experimental evolution of organisms: Bull. / NAS of Ukraine, Ukrainian Academy of Agricultural Sciences, NAMS of Ukraine, Institute of Molecular Biology and Genetics, NAS of Ukraine, Ukr. society of geneticists and breeders named after M.I. Vavilov; Ed.: V.A. Kunah (ed.-in-chief) et al.. – K.: Logos, 2013. – P. 214-219.

Наследование и трансгрессивная изменчивость массы зерна с растения в гибридов пшеницы озимой (*T. aestivum L.*), полученных от гибридизации разных экотипов

Н.В. Лозинский

Показано особенности наследования массы зерна с растения гибридами F₁ пшеницы мягкой озимой. Степень фенотипического доминирования (h_p) составляла 1,4-64,0. Истинный гетерозис по массе зерна с растения наблюдался в девяти с десяти гибридов F₁ с показателем 41,1-68,9 %. Частота положительных трансгрессивных рекомбинантов по массе зерна с растения у гибридов F₂, полученных от скрещивания степного экотипа с лесостепным составляла 36,0-80,2 %. У гибридных популяций, полученных от скрещивания отдаленных эколого-географических форм, наибольшее количество позитивных трансгрессий наблюдалось в популяциях Гайтун / Олеся и Гайтун / Белоцерковская – полукарликовая 68,8 и 57,1 % соответственно.

Ключевые слова: пшеница мягкая озимая, масса зерна с растения, экотипы, комбинации скрещивания, гибриды, наследование, гетерозис, степень доминирования, степень и частота трансгрессий.

Inheritance and grain weight transgressive variability per plant in hybrid winter wheat (*T. aestivum L.*), obtained from the hybridization of various ecotypes

M. Lozinskyi

The peculiarities of grain weight inheritance per plant in the F₁ soft winter wheat hybrids is highlighted. It has been found out that grain weight inheritance per plant in first generation hybrids in most combinations develops on the type of positive overdominance. The degree of phenotypic dominance (h_p) was 1.4-64.0. True heterosis for grain weight per plant was observed in nine of the ten F₁ hybrids with the index of 41.1-68.9 %. The frequency of positive transgressive recombinants by crop grain weight in F₂ hybrids, obtained by crossing the steppe and forest-steppe ecotypes was 36.0-80.2 %. The highest number of the positive transgressions in hybrid populations obtained with crossing remote eco-geographical forms, was observed in Haytun / Olesya and Haytun / Bilotserkivska, semi-dwarf populations – 68.8 and 57.1 % respectively.

Key words: soft winter wheat, the grain weight per plant, ecotypes, crossbreeding combinations, hybrids, inheritance, heterosis, dominance degree, transgressions degree and frequency.

Надійшла 14.04.2016 р.

УДК 636.085.2.51:632.51:632.93: 631.584.5

ГРАБОВСЬКИЙ М.Б., ГРАБОВСЬКА Т.О., ОБРАЖІЙ С.В., кандидати с.-г. наук
Білоцерківський національний аграрний університет

**ФОРМУВАННЯ ПРОДУКТИВНОСТІ
СУМІСНИХ ПОСІВІВ КУКУРУДЗИ І СОРГО ЦУКРОВОГО
ЗАЛЕЖНО ВІД ЗАХОДІВ ЗАХИСТУ РОСЛИН ВІД БУР'ЯНІВ**

Наведено результати досліджень з вивчення впливу заходів захисту від бур'янів на продуктивність сумісних посівів кукурудзи та сорго цукрового. Механізований догляд за посівами забезпечує знищення 80,1 % бур'янів, технічна ефективність гербіцидів становить – 58,6-69,5 %. Використання післясходового гербіциду Примекстра TZ Голд забезпечує максимальну урожайність зеленої маси – 77,5 т/га і збору сухої речовини – 19,8 т/га. Дослідженнями встановлено, що застосування гербіцидів є оптимальним заходом захисту від бур'янів рослин кукурудзи і сорго цукрового для реалізації їх біологічного потенціалу в сумісних посівах.

Ключові слова: кукурудза, сорго цукрове, сумісні посіви, продуктивність, суха речовина, гербіциди, механізований догляд.

Постановка проблеми. Одним з факторів, які стримують збільшення виробництва кукурудзи і сорго, є бур'яни. Через негативний вплив на рівень вологозабезпеченості посівів, а також надходження поживних речовин вони блокують 35–60 % всіх факторів росту й розвитку цих культур [16].

Бур'яни досить добре пристосувались до умов існування в агроценозах сільськогосподарських культур і відзначаються можливістю розвитку за несприятливих умов. Набагато витриваліші