



UDC 582-152.42 (477.85)

**CHOROLOGICAL, ECOLOGICAL AND COENOTIC CHARACTERISTICS
OF *SYMPHYOTRICHUM CILIATUM* (LINDL.) NESOM (ASTERACEAE)
IN THE BUKOVINIAN CIS-CARPATHIAN**

A. I. Tokaryuk¹, I. I. Chorney¹, V. V. Budzhak¹, V. V. Protopopova², M. V. Shevera^{3,2}

¹ Yuriy Fedkovych Chernivtsi National University, 11, Fedkovych St., Chernivtsi 58022, Ukraine
e-mail: iichorney@ukr.net, vbudzhak@gmail.com

² Ferenc Rakoczi II Transcarpathian Hungarian Institute
6, Koshut Sq., Beregovo 90200, Ukraine
e-mail: protopopova.vira@gmail.com

³ M.G. Kholodny Institute of Botany, NAS of Ukraine
2, Tereschenkivska St., Kyiv 01004, Ukraine
e-mail: shevera.myroslav@gmail.com

The results of study on distribution, ecological and coenotic characteristics of alien species *Symphotrichum ciliatum* (Lindl.) Nesom (Asteraceae) in the Bukovinian Cis-Carpathian (Chernivtsi Region, Ukraine) are presented. Five associations of synanthropic vegetation species were found in the region namely: *Eragrostio-Amarantheum albi* Morariu 1943, class *Digitario sanguinalis-Eragrostietea minoris* Mucina, Lososová et Šilc in Mucina et al. 2016, *Lolio-Plantaginetum majoris* (Linkola 1921) Beger and *Polygonetum arenastri* Gams 1927, class *Polygono-Poetea annuae* Rivas-Mart. 1975, *Odontito-Ambrosietum artemisiifoliae* Mucina 1993 and *Convolvulo-Agropyretum repentis* Felföldy 1943, class *Artemisietea vulgaris* Lohmeyer et al. in Tx. ex von Rochow 1951. These species are noted in the anthropogenic ecotopes and in habitat of Nature 2000. The indices of main edaphic and climatic factors for the species were determined for the first time. According to the main climatic indices, such as thermoregulation, ombrorezhime and criorezhime, *S. ciliatum* is characterized by stenotopic amplitude and belongs to submezoterms, subaridophytes and hemicriophytes, etc. *S. ciliatum* in the region is characterized as potential invasive species.

Keywords: *Symphotrichum ciliatum*, Asteraceae, distribution, ecological and coenotic characteristics, the Bukovinian Cis-Carpathian, Ukraine

INTRODUCTION

Symphotrichum ciliatum (Lindl.) Nesom (*Brachyactis ciliata* (Ledeb.) Ledeb.) is an alien species with the debatable systematic position caused its high variability, vague diagnostic morphological features, disjunctive area etc. Previously florists and taxonomists have considered this species as a member of different genera, e.g. *Aster* L.,

Erigeron L., *Conyza* Less., *Tripolium* Nees. In Europe it was often referred to the genus *Brachyactis* Ledeb. [12, 26, 32] or to the genus *Symphotrichum* Nees [2, 17, 23, 31]. Due to the karyological studies carried out by Nesom [17] it was established that species of genera *Symphotrichum* and *Brachyactis* had the different number of chromosomes, ($2n=14$) and ($2n = 18$) respectively. This and also significant distinction between taxa have been allowed consider the studied species as a member of the section *Conyzopsis* (Torr & A. Gray) G.L. Nesom, subgenus *Symphotrichum*, genus *Symphotrichum* [2, 17, 23, 31].

An interpretation of the origin of this species is also debatable: as *S. ciliatum*, sub nom. *B. ciliate* it maybe Asian [1, 9–10, 12, 24–26, 32] or North American [13]; as *S. ciliatum* it maybe North American, North and Middle Asian [30]. However study conducted by G.L. Nesom [17] showed that European populations as well as North American ones have the number of chromosomes $2n=14$ and therefore it is necessarily to include them in the genus *Symphotrichum*.

The natural area of *S. ciliatum* is North America (Rocky Mountains) where it grows in dry places. Later the species migrated to the prairie of the Great Plains, and also through Alaska and the Bering Strait to the North-Eastern of Asia.

The general area of *S. ciliatum* is Circumpolar, disjunctive. It distributes in North America, West and East Sibirian, Far East and Central Asia [2, 12, 17, 26]. At present its active distribution is observed in the East of the American continent [4] and in Europe (Ukraine, Moldova, Romania, Poland, Slovakia, Russia) [7–10, 18, 24, 31], where *S. ciliatum* manifests a wide ecological amplitude growing on different substrata from dry sands to saline, moist soils, but mostly prefers some anthropogenic habitats.

In Ukraine, *S. ciliatum* (sub nom. *B. ciliate*) was first recorded in Danube Biosphere Reserve [7]. Later it has been found in Steppe, mainly in the anthropogenic and semi-natural (coastal, seaside and steppe) ecotopes [21]. According to *CHER* and *KW* herbarium collections, it is also known from Transcarpathia, Bukovinian Cis-Carpathian, Prut-Dnister interfluve, Zhytomyr Polissia and Roztochchya [13].

In Bukovinian Cis-Carpathian, *S. ciliatum* was first recorded in 2007 [22]. Later some more new localities were found. Very likely, this species penetrated in the The Bukovinian Cis-Carpathian from Romania where it had been determined as potential invasive [24–25]. The status of the species in Poland the same [30].

The aim of the article is to study of current distribution of *S. ciliatum* in the Bukovinian Cis-Carpathian, rate and character of dispersal, to determine ecological and coenotic characteristics and estimate the possible consequences of its naturalization in the region.

MATERIAL AND METHODS

Investigations are based on the original data obtained during fieldwork over the years 2007–2016 and an examination of herbarium collections of the Yuri Fedkovych Chernivtsi National University (*CHER*) and M.G. Kholodny Institute of Botany, NAS of Ukraine (*KW*).

The schematic map of the *S. ciliatum* distribution in the The Bukovinian Cis-Carpathian was prepared by the grid method of plant mapping using MapInfo Professional version 12.5 software packages, according to Atlas Florae European adapted for regional territories [3]. The phytosociological surveys were performed with using ecological and floristic criteria of plant community releves. The phytosociological releves were carried out by the method of transformation of phytocoenotic tables with using the program

package FICEN 2 [28]. The calculation of score synphytoindication indices was performed with the use of JUICE 6.5 program [29].

Ecological analysis was made according to the phytoindicatory scales [6, 20]. Syntaxonomic scheme of plant communities with the participation of *S. ciliatum* was compiled following by Vegetation of Europe [14–16].

RESULTS

The information on the distribution of *S. ciliatum* in the The Bukovinian Cis-Carpathian generalized on based of the herbarium collections *CHER* and *KW* data and fieldwork is given in fig. 1 and list mentioned below.

List of localities of *S. ciliatum* in Chernivtsi Region:


1. Glyboka Distr., Glyboka-Bukovynska station, along railway tracks, 12.10. 2007, I. Chorney, V. Budzhak, A. Tokaryuk (*CHER*); the same, 12.10. 2007, V.V. Protopopova, M.V. Shevera (*KW*).
2. Glyboka Distr., vicinity of Cherepkivtsi vill., Vadul-Seret railway station, along railway tracks, 12.10. 2007, I. Chorney, V. Budzhak, A. Tokaryuk (*CHER*); the same, 12.10. 2007, V.V. Protopopova, M.V. Shevera (*KW*).
3. Storozhynets Distr., Storozhynets, railway station, 20.09. 2009, A. Tokaryuk (*CHER*).
 - Storozhynets Distr., Storozhynets, along railway, 18.10. 2009, A. Tokaryuk (*CHER*).
4. Storozhynets Distr., Storozhynets, along road, 21.11. 2009, A. Tokaryuk (*CHER*);
 - Storozhynets Distr., Storozhynets, along field road, 03.11. 2013, A. Tokaryuk (*CHER*).
5. Storozhynets Distr., Storozhynets city, left bank of Siret River, 30.09. 2012, A. Tokaryuk (*CHER*).

S. ciliatum actively distributes in the The Bukovinian Cis-Carpathian in anthropogenic transformed plant communities. In an early stage of invasion and colonization of ruderalized plots with denuded soil by this species it is characterized by high density and number. In the overgrowing stage of the plots, when the total cover of grass increases and vegetation becomes closer, the mentioned population indices of the species decrease.

The species is an annual/terophyte. Due to some biological characteristics such as the short period of ontogenesis and high reproductively it can extend to rather a long distance from parent individuals mainly by virtue of disturbed plots with a significant proportion of denuded soil. However, a long keep of colonizing territory do not observed especially in case of the process of overgrowing.

The densest and numerous populations of *S. ciliatum* in the The Bukovinian Cis-Carpathian were revealed in Glyboka city in plant community related to artificial rubble plots between railway tracks within the Glyboka-Bukivynska railway station. Floristic composition of the community is rather poor and composed of 19 species. Its total projective cover comes to 30–40 % and projective cover of *S. ciliatum* to 5–7 %. Diagnostic *Eragrostis minor* Host and *Amaranthus albus* L. dominated. Among the invasion species there are *Conyza canadensis* (L.) Cronq.), *Echinochloa crusgalli* (L.) P. Beauv., *Galinsoga parviflora* Cav. and *Phalacrolooma annuum* (L.) Dumort. (Table 1, relevé 1). According to ecological and floristic classification, the plant community belongs to

association *Eragrostio-Amarantheum albi* Morariu 1943, alliance *Eragrostion cilianensis-minoris* Tx. ex Oberd. 1954, order *Eragrostietalia* J. Tx. ex Poli 1966 and class *Digitario sanguinalis-Eragrostietea minoris* Mucina, Lososová et Šilc in Mucina et al. 2016.



Map of localities of *Symphyotrichum ciliatum* (Lindl.) Nesom in the Bukovinian Cis-Carpathian
Карта поширення *Symphyotrichum ciliatum* (Lindl.) Nesom у Буковинському Передкарпатті

Table 1. Phytocoenotic characteristics of plant communities with participation of *Symphyotrichum ciliatum* (Lindl.) Nesom in the Bukovinian Cis-Carpathian

Таблиця 1. Фітоценотична характеристика рослинних угруповань за участю *Symphyotrichum ciliatum* (Lindl.) Nesom у Буковинському Передкарпатті

| Number of relevés | 1 | 2 | 3 | 4 | 5 | 6 |
|--|----------|----------|----------|----------|----------|----------|
| Total projective cover, % | 30–40 | 70 | 50 | 70 | | 100 |
| Total number of species | 19 | 24 | 24 | 20 | 30 | 26 |
| Number of syntaxa | 1 | 2 | | 3 | 4 | 5 |
| <i>Symphyotrichum ciliatum</i> | 2 | 1 | + | + | 2 | + |
| D.s. Ass. <i>Eragrostio-Amarantheum albi</i> | | | | | | |
| <i>Eragrostis minor</i> | 2 | . | . | . | . | . |
| <i>Amaranthus albus</i> | 1 | . | . | . | . | . |
| D.s. Cl. <i>Digitario sanguinalis-Eragrostietea minoris</i> | | | | | | |
| <i>Echinochloa crusgalli</i> | + | . | 1 | 1 | . | . |
| <i>Setaria glauca</i> | . | . | . | 1 | + | . |
| <i>Portulaca oleracea</i> | + | . | . | . | . | . |
| D.s. Ass. <i>Lolio-Plantaginietum majoris</i> | | | | | | |
| <i>Plantago major</i> | . | 2 | 3 | 4 | . | + |
| <i>Lolium perenne</i> | . | 1 | . | . | . | . |
| <i>Lepidotheca suaveolens</i> | . | + | . | . | . | . |

| D.s. Ass. Polygonetum avicularis | | | | | | |
|---|---|---|---|---|---|---|
| <i>Polygonum aviculare</i> | . | 1 | 2 | 2 | . | . |
| <i>Poa annua</i> | . | . | 1 | . | . | . |
| D.s. Cl. Polygono-Poetea annua | | | | | | |
| <i>Taraxacum officinale</i> | + | + | + | + | + | + |
| <i>Medicago lupulina</i> | + | 1 | + | . | + | . |
| <i>Lepidium ruderales</i> | . | + | . | . | . | . |
| <i>Potentilla anserina</i> | . | . | . | + | . | . |
| D.s. Ass. Odontito-Ambrosietum artemisiifoliae | | | | | | |
| <i>Ambrosia artemisiifolia</i> | . | 5 | 2 | 2 | 3 | 1 |
| <i>Odontites vulgaris</i> | . | . | . | . | 1 | . |
| D.s. Ass. Convolvulo-Agropyretum repentis | | | | | | |
| <i>Calamagrostis epigeios</i> | . | . | . | . | . | 1 |
| <i>Elytrigia repens</i> | . | 1 | . | . | 1 | 4 |
| D.s. Cl. Artemisietea vulgaris | | | | | | |
| <i>Phalacrolooma annuum</i> | 1 | + | + | 1 | 1 | 1 |
| <i>Artemisia vulgaris</i> | 1 | + | . | + | . | + |
| <i>Erigeron canadensis</i> | + | . | + | + | + | . |
| <i>Daucus carota</i> | . | 1 | . | 1 | + | 1 |
| <i>Cirsium arvense</i> | + | . | . | + | . | . |
| <i>Arctium lappa</i> | . | . | . | + | . | 1 |
| <i>Cichorium intybus</i> | . | . | . | . | + | + |
| <i>Arctium tomentosum</i> | . | + | . | . | . | . |
| <i>Lactuca serriola</i> | . | . | + | . | . | . |
| <i>Linaria vulgaris</i> | . | . | . | . | + | . |
| <i>Melilotus officinalis</i> | . | . | . | . | . | + |
| D.s. Cl. Sisymbrietea | | | | | | |
| <i>Amaranthus retroflexus</i> | . | . | + | + | + | . |
| <i>Galinsoga urticifolia</i> | . | + | + | . | . | . |
| <i>Matricaria perforata</i> | . | + | + | . | . | . |
| <i>Atriplex patula</i> | + | . | . | . | . | . |
| <i>Chenopodium album</i> | + | . | . | . | . | . |
| <i>Galinsoga parviflora</i> | 1 | . | . | . | . | . |
| <i>Ranunculus sardous</i> | + | . | . | . | . | . |
| <i>Xanthoxalis dillenii</i> | . | . | + | . | . | . |
| D.s. Cl. Epilobietea angustifolii | | | | | | |
| <i>Galega officinalis</i> | . | + | . | . | . | + |
| <i>Glechoma hederacea</i> | . | + | . | . | . | . |
| <i>Urtica dioica</i> | . | + | . | . | . | . |
| <i>Ranunculus repens</i> | . | . | . | . | + | . |
| <i>Rumex confertus</i> | . | . | . | . | . | + |

| D.s. Cl. <i>Molinio-Arrhenatheretea</i> | | | | | | |
|---|---|---|---|---|---|---|
| <i>Plantago lanceolata</i> | . | . | 1 | 1 | 1 | + |
| <i>Trifolium repens</i> | . | 1 | 1 | . | + | . |
| <i>Trifolium pratense</i> | . | . | + | . | + | + |
| <i>Poa pratensis</i> | . | . | + | 1 | . | 1 |
| <i>Agrostis tenuis</i> | . | . | . | . | 1 | 1 |
| <i>Festuca pratensis</i> | . | . | . | . | . | + |
| <i>Phleum pratense</i> | . | . | . | . | . | 1 |
| <i>Dactylis glomerata</i> | . | . | . | . | . | 2 |
| <i>Cerastium holosteoides</i> | + | . | . | . | . | . |
| <i>Trifolium hybridum</i> | . | . | . | + | . | . |
| <i>Mentha arvensis</i> | . | . | . | + | . | . |
| Other species: | | | | | | |
| <i>Achillea submillefolium</i> | + | . | + | . | + | + |
| <i>Lotus ucrainicus</i> | . | . | + | . | + | 1 |
| <i>Polygonum persicaria</i> | . | + | . | . | . | + |
| <i>Bidens tripartita</i> | 1 | . | + | . | . | . |
| <i>Barkhausia setosa</i> | . | . | + | . | + | . |
| <i>Chaenorhinum minus</i> | . | . | + | . | + | . |
| <i>Potentilla reptans</i> | . | . | . | . | + | + |

The species occurred only in one relevé: *Anchusa officinalis* (5 [+]); *Anisantha tectorum* (2 [1]); *Berteroa incana* (5 [+]); *Echium vulgare* (5 [+]); *Eragrostis pilosa* (4 [1]); *Galium verum* (5 [+]); *Geum urbanum* (6 [+]); *Inula britannica* (2 [+]); *Juncus bufonius* (2 [1]); *Juncus tenuis* (6 [+]); *Lepidium densiflorum* (5 [+]); *Oenothera biennis* (5 [+]); *Poa compressa* (3 [+]); *Polygonum convolvulus* (1 [+]); *Potentilla supina* (4 [+]); *Swida sanguinea* (5 [+]); *Xanthium albinum* (5 [1])

Види, які трапляються в одному описі: *Anchusa officinalis* (5 [+]); *Anisantha tectorum* (2 [1]); *Berteroa incana* (5 [+]); *Echium vulgare* (5 [+]); *Eragrostis pilosa* (4 [1]); *Galium verum* (5 [+]); *Geum urbanum* (6 [+]); *Inula britannica* (2 [+]); *Juncus bufonius* (2 [1]); *Juncus tenuis* (6 [+]); *Lepidium densiflorum* (5 [+]); *Oenothera biennis* (5 [+]); *Poa compressa* (3 [+]); *Polygonum convolvulus* (1 [+]); *Potentilla supina* (4 [+]); *Swida sanguinea* (5 [+]); *Xanthium albinum* (5 [1])

Association localities of the relevés:

1. Glyboka, railway station, along railway, 12.10. 2007, A. Tokaryuk;
2. Glyboka, railway station, along railway, 29.07. 2008, A. Tokaryuk;
3. Storozhynets, railway station, 21.09. 2009, A. Tokaryuk;
4. Storozhynets, railway station, 21.09.2 009, A. Tokaryuk;
5. Glybotskyi Distr., Cherepkivtsi vill., Vadul-Siret railway station, 12.10. 2007, A. Tokaryuk;
6. Glyboka, railway station, 12.10. 2007; 29.07. 2008, A. Tokaryuk

Описи виконано:

1. смт Глибока, залізничний вокзал, уздовж залізничної колії, 12.10. 2007, А. Токарюк;
2. смт Глибока, залізничний вокзал, уздовж залізничної колії, 29.07. 2008, А. Токарюк;
3. м. Сторожинець, залізничний вокзал, 21.09. 2009, А. Токарюк;
4. м. Сторожинець, залізничний вокзал, 21.09.2 009, А. Токарюк;
5. Глибоцький р-н, с. Черепківці, залізнична станція Вадул-Сірет, 12.10. 2007, А. Токарюк;
6. смт Глибока, залізничний вокзал, 12.10. 2007, 29.07. 2008, А. Токарюк

Number of syntaxa: 1 – *Eragrostio-Amarantheum albi*; 2 – *Lolio-Plantaginetum majoris*; 3 – *Polygonetum arenastris*; 4 – *Odontito-Ambrosietum artemisiifoliae*; 5 – *Convolvulo-Agropyretum repentis*

Номери синтаксонів: 1 – *Eragrostio-Amarantheum albi*; 2 – *Lolio-Plantaginetum majoris*; 3 – *Polygonetum arenastris*; 4 – *Odontito-Ambrosietum artemisiifoliae*; 5 – *Convolvulo-Agropyretum repentis*

The plant community of association *Lolio-Plantaginetum majoris* (Linkola 1921) Beger with participation of *S. ciliatum* was found along a path at a beet collection in the territory Glyboka-Bukovynska station. Its total projective cover comes to 80 % and projective cover of *S. ciliatum* is 1–2 %. Floristic composition of association composed of 24 species. The diagnostic *Lolium perenne* L., *Plantago major* L. and *Lepidotheca suaveolens* (Pursh) Nuttm dominate. Components of the plant community are invasive *Ambrosia artemisiifolia* L. (50–60 %), *Galinsoga urticifolia* (Kunth) Benth. and *Phalacrolooma annuum* (Table 1, relevé 2).

The association belongs to alliance *Polygono-Coronopodion* Sissingh 1969 order *Polygono arenastri-Poetalia annuae* Tx. in Géhu et al. 1972 corr. Rivas-Mart. et al. 1991, class *Polygono-Poetea annuae* Rivas-Mart. 1975 [26].

Within alliance *Polygono-Coronopodion* *S. ciliatum* have been found in plant community of association *Polygonetum arenastri* Gams 1927 confined to a roadside on the territory of Storozhynets railway station. The number of species in the coenosis varies from 20 to 25. Its total projective cover is 50–70 %. Diagnostic species of association *Plantago major* L. (20–50 %), *Poa annua* L. (2–3 %), and *Polygonum aviculare* L. (5–7 %) dominate. *S. ciliatum* occurs rare. Among invasive species *Ambrosia artemisiifolia* (5–7 %), *Conyza canadensis*, *Echinochloa crusgalli* (2–3 %), *Galinsoga urticifolia*, *Phalacrolooma annuum*, *Setaria glauca* (L.) P. Beauv. (2–3 %), and *Xanthoxalis dillenii* (Jacq.) Holub are observed (Tabl. 1, relevés 3–4).

Plant community of association *Odontito-Ambrosietum artemisiifoliae* Mucina 1993 with participation of *S. ciliatum* was revealed in Vadul-Siret railway station near Cherepkivtsi village of Glyboka District related to open places along path between railway tracks. Total projective cover is 80 %, projective cover of *S. ciliatum* is 1–2 %. Diagnostic species: *Ambrosia artemisiifolia* is dominant and *Odontites vulgaris* Moench is co-dominant in the plant community. The plant community composes of 30 species of vascular plants, including invasive *Cichorium intybus* L., *Conyza canadensis*, *Phalacrolooma annuum*, *Setaria glauca* and *Xanthium albinum* (Widder) H.Scholz (Table 1, relevé 5). It belongs to alliance *Onopordion acanthii* Br.-Bl. 1926, order *Onopordetalia acanthii* Br.-Bl. et Tx. ex Klika et Hadač 1944, class *Artemisietea vulgaris* Lohmeyer et al. ex von Rochow 1951 [11].

The isolated individuals of *S. ciliatum* occurs also in plant community of association *Convolvulo-Agropyretum repentis* Felföldy 1943 distributed along ground roads within the Glyboko-Bukovynska railway station territory. In this plant community (total projective cover 100 %) *Elytrigia repens* dominates with negligible dirt of *Agrostis capillaris* L., *Calamagrostis epigeios* (L.) Roth, *Dactylis glomerata* L., *Festuca pratensis* Huds., *Phleum pratense* L., and *Poa pratensis* L. Components of the association are invasive *Ambrosia artemisiifolia*, *Cichorium intybus*, *Phalacrolooma annuum* and *Juncus tenuis* Willd. (Table 1, relevé 6). Plant community of association belongs to alliance *Convolvulo arvensis-Agropyrion repentis* Görs 1967, order *Agropyretalia intermedio-repentis* T. Müller et Görs 1969, class *Artemisietea vulgaris* Lohmeyer et al. in Tx. ex von Rochow 1951.

Thus, on the basis of phytosociological relevé it was established that *S. ciliatum* in the Bukovinian Cis-Carpathian grows in five association, four order and three classes of synanthropic vegetation. The syntaxonomic scheme of plant community with participation of *S. ciliatum* in the region is given.

Syntaxonomic scheme of plant community with participation of *Symphotrichum ciliatum* (Lindl.) Nesom in Bukovinian Cis-Carpathian

Синтаксономічна схема рослинних угруповань за участю *Symphotrichum ciliatum* (Lindl.) Nesom у Буковинському Передкарпатті

Cl. *Digitario sanguinalis-Eragrostietea minoris* Mucina, Lososová et Šilc in Mucina et al. 2016

Ord. *Eragrostietalia* J. Tx. ex Poli 1966

All. *Eragrostion cilianensi-minoris* Tx. ex Oberd. 1954

Ass. *Eragrostio-Amaranthesetum albi* Morariu 1943

Cl. *Polygono-Poetea annuae* Rivas-Mart. 1975

Ord. *Polygono arenastri-Poetalia annuae* Tx. in Géhu et al. 1972 corr. Rivas-Mart. et al. 1991

All. *Polygono-Coronopodium* Sissingh 1969

Ass. *Lolio-Plantagnetum majoris* (Linkola 1921) Beger

Ass. *Polygonetum arenastri* Gams 1927

Cl. *Artemisietea vulgaris* Lohmeyer et al. in Tx. ex von Rochow 1951

Ord. *Meliloto-Artemisitalia absinthii* Elias 1979

All. *Dauco-Melilotion albi* Gors em Elias 1980

Ass. *Odontito-Ambrosietum artemisiifoliae* Mucina 1993

Ord. *Agropyretalia intermedio-repentis* T. Müller et Görs 1969

All. *Convolvulo arvensis-Agropyron repentis* Görs 1967

Ass. *Convolvulo-Agropyretum repentis* Felföldy 1943

According to the main climatic indices, such as thermoregulation, ombrorezhime and criorezhime, *S. ciliatum* is characterized by stenotopic amplitude and belongs to submezoterms, subaridophytes and hemicrophytes (Table 2).

Table 2. **Ecological characteristic (climatop) of *Symphotrichum ciliatum* (Lindl.) Nesom in the Bukovinian Cis-Carpathian**

Таблиця 2. **Екологічна характеристика (кліматоп) *Symphotrichum ciliatum* (Lindl.) Nesom у Буковинському Передкарпатті**

| Indices of climatic factors (points), ecological groups, width of ecological amplitude | | | |
|--|--------------------------|----------------------------|-------------------------|
| Tm* | Om | Kn | Cr |
| <u>8.78–9.90</u> | <u>11.07–11.65</u> | <u>8.45–9.11</u> | <u>8.17–8.93</u> |
| 9.26 | 11.38 | 8.79 | 8.56 |
| Stenotopic submezoterm | Stenotopic subaridophyte | Stenotopic hemicontinental | Stenotopic hemicrophyte |

Comment: Tm – thermorezhime, Om – ombrorezhime, Kn – continental climate, Cr – criorezhime; its and in the table 3 indicated: in numerator – value of tolerance amplitude, in denominator – mean value of tolerance amplitude to ecological factor

Примітка: Тм – терморезим, Ом – омброрезим, Кн – континентальність клімату, Сг – криорезим; тут і в табл. 3 – у чисельнику вказується величина амплітуди толерантності, у знаменнику – середнє значення амплітуди толерантності до цього екологічного фактора

In the region the in relation to the water regime of the soil and the variability of moisture *S. ciliatum* is stenotypic mesophyte and hemihydroconstrastophile and grows

in forest-meadow ecotopes with complete blotting root zone and melted water ($W_{np} = 100\text{--}145$ mm) (Table 3). It grows on slightly acidic ($\text{pH} = 5.5\text{--}6.5$), enriched with salts, relatively nitrogen poor soils of mineral nitrogen and avoids carbonate substrates.

Table 3. Ecological characteristic (edaphotop) of *Symphotrichum ciliatum* (Lindl.) Nesom in the Bukovinian Cis-Carpathian

Таблиця 3. Екологічна характеристика (едафотоп) *Symphotrichum ciliatum* (Lindl.) Nesom у Буковинському Передкарпатті

| Indices of edaphic factors (pointsy), ecological groups, wide of ecological amplitude | | | | | | |
|---|---|--------------------------|-------------------------|----------------------------------|----------------------------|--------------------------------|
| Hd* | fH | Rc | Tr (Sl) | Ca | Nt | Ae |
| <u>10.73–11.60</u> | <u>7.04–7.75</u> | <u>7.79–8.05</u> | <u>7.78–8.32</u> | <u>6.22–7.13</u> | <u>5.88–6.82</u> | <u>6.25–7.16</u> |
| 11.27 | 7.38 | 7.94 | 8.15 | 6.82 | 6.45 | 6.84 |
| stenotopic tmezophyte | c stenotopic hemihydro- contrastophyte | stenotopic subacidophile | stenotopic semieuritope | stenotopic hemicarbonatophyle | stenotopic heminitrophylae | Hemistenotopic subaerophobe |

Comment: Hd – humidity of soil, fH – moisture change, Rc – soil acidity, Sl – soil salinity, Ca – content of carbonates in soil, Nt – content of nitrogen in soil, Ae – soil aeration

Примітка: Hd – вологість ґрунту, fH – змінність зволоження, Rc – кислотність ґрунту, Sl – засоленість ґрунту, Ca – вміст у ґрунті карбонатів, Nt – вміст азоту у ґрунті, Ae – аерація ґрунту

As to rate and character of distribution in the Bukovinian Cis-Carpathian *S. ciliatum* is potential invasive species.

According to the classification of habitats on the territory of Ukrainian Carpathians and Transcarpathian Lowlands [4], plant communities with *S. ciliatum* are comprised of the next habitats: “Sand-clay deposits (shifts) of quarries and anthropogenic substrates of a moving type”; “Seliteb area”; “Areas with ruderal xeromesophyte vegetation (beyond the seliteb area)”; “Muddy river banks with *Chenopodium rubri* p. p. and *Bidention* p. p. Vegetation”. Last one is under threat of disappearance and need the protection on European level because of its presence in the Natura 2000, in particular, 3270 “Muddy river banks with *Chenopodium rubri* p. p. and *Bidention* p. p. Vegetation” [19].

CONCLUSION

The *S. ciliatum* in the Bukovinian Cis-carpathian occurs in five associations, four unions, four orders, and three classes of synanthropic vegetation: Ass. *Eragrostio-Amaranthenetum albi* Cl. *Digitario sanguinalis-Eragrostietea minoris*, Ass. *Lolio-Plantaginetum majoris* and *Polygonetum arenastri* Cl. *Polygono-Poetea annuae*, Ass. *Odontito-Ambrosietum artemisiifoliae* and *Convolvulo-Agropyretum repentis* Cl. *Artemisietea vulgaris*.

The species in the region has stenotope ecological amplitude and according to main climatic and edaphic indices is submezoterm, subaridophyte, hemicontinental, hemicriophyte, mesophyte, hemihydroconstrastophile, subacidophile, semievitrophe, gemicarbonatophyte, heminitrophylous and subaerophobe. One can consider this species as potential invasive.

ACKNOWLEDGEMENT

The authors would like to thank the Dr. Olena V. Borysova (M.G. Kholodny Institute of Botany, NAS of Ukraine) for valuable comments and editing of English text.

1. *Anastasiu P., Negrean G., Samoilă C., Memedemin D., Cogălniceanu D.* A comparative analysis of alien plants species along the Romanian Black Sea coastal area. The role of harbours. **J. Coast. Conserv.**, 2011; 15: 595–606.
2. *Broż E., Podgórska M.* *Symphytotrichum ciliatum (Brachyactis ciliata) (Asteraceae)* in Poland. **Fragm. Flor. et Geobot. Polon.**, 2005; 12(2): 291–299.
3. *Budzhak V. V., Chorney I. I., Tokaryuk A. I.* Supplement to the method of flora species mapping (Chernivtsy district example). **Scientific Herald of Chernivtsi University: collection of scientific papers. Biology**, 2009; 455: 168–170. (In Ukrainian).
4. **Catalogue of type habitats of Ukrainian Carpathian and Transcarpathian lowland** / Ed. B. Prots and O. Kagalo. Lviv: Merkator Press, 2012. 294 p. (In Ukrainian).
5. **COSEWIC assessment and update status report on the Gulf of St. Lawrence aster *Symphytotrichum laurentianum* in Canada.** Committee on the Status of Endangered Wildlife in Canada. Ottawa, 2004: 39.
6. *Didukh Ya.P.* Prepararation of edition of “Ecoflora of Ukraine” [as base of phytoindication state of ecosystem]. **Ukr. Phytocoen J.**, 1998; 1(10): 4–17. (In Ukrainian).
7. *Dubyna D.V., Protopopova V.V., Dubovik O.M.* New for the flora of Ukr. SSR species *Brachyactis* Ledeb. **Ukr. Bot. J.**, 1986. 43(2): 51–54. (In Ukrainian).
8. *Geydeman T.S.* **Determinator of Higher plants of Moldavian SSR.** Kishinyov: Shtiinca Press, 1986: 530. (In Russian).
9. *Guzik J.* Hałda Huty im. Sendzimira w Krakowie – miejsciem występowania interesujących obcych gatunków roślin. **Sesja Naukowa: Hałda przemysłowa – obiekt obserwacji procesów biologicznych.** Uniwersytet Śląski w Katowicach, 2002: 7.
10. *Guzik J.* Hałda Huty im. Sendzimira w Krakowie – miejscem występowania interesujących obcych gatunków roślin. **Arch. Ochr. Środ.**, 2003; 29(2): 13–19.
11. *Jarolímek I., Zaliberová M., Mucina L., Mochnacký S.* **Rastlinné spoločenstvá Slovenska 2. Synantropná vegetácia.** Bratislava: Veda Vydavateľstvo Slovenskej Akadémie Vied, 1997: 416 s.
12. *Krylov P.F.* **Flora of Western Siberia.** Vol. 11. Tomsk: Botanical Section of Tomsk Nature Society, 1949. P. 2630–3070. (In Russian).
13. *Kuzyarin A..T.* New for Lviv Region alien species. **Sc. Letters of State Natural History Museum**, 2012; 28: 143–144. (In Ukrainian).
14. *Matuszkiewicz W.* **Przewodnik do oznaczania zbiorowisk roślinnych Polski.** Warszawa: Wydawnictwo naukowe PWN, 2001. 537 s.
15. *Moravec J. a kol.* **Roslinna společenstva České republiky a jejich ochrozeni.** 2 vydání. ČR: Okresni vlastivědné muzeum, 1995. 206 s.
16. *Mucina L., Bültmann H., Dierßen K., Theurillat J.-P., Raus T., Čarni A., Šumberová K., Willner W., Dengler, J., Gavilán García R., Chytrý M., Hájek M., Di Pietro R., Iakushenko D., Pallas J., Daniěls F.J.A., Bergmeier E., Santos Guerra A., Ermakov N., Valachovič M., Schaminée J.H.J., Lysenko T., Didukh Ya.P., Pignatti S., Rodwell J.S., Capelo J., Weber H.E.,*

- Solomeshch A., Dimopoulos P., Aguiar C., Hennekens S.M. & Tichý, L.* Vegetation of Europe: Hierarchical floristic classification system of vascular plant, bryophyte, lichen, and algal communities. **Applied Vegetation Science**, 2016; 19(1): 3–264.
17. *Nesom G.L.* Review of the taxonomy of the *Aster* sensu lato (Asteraceae: Astereae), emphasizing the New World species. **Phytologica**, 1994; 77(3): 141–297.
 18. *Nobis M., Pliszko A.* New localities of *Symphotrichum ciliatum* (Asteraceae) in Poland. **Acta Mus. Siles. Sci. Natur**, 2016; 65: 283–286.
 19. **Habitat conception of conservation of biodiversity: bases documents of EU** / ed. O.O. Kagalo & B.H. Prots. Lviv ZUKTs press, 2012. 278 p. (In Ukrainian).
 20. *Plyuta P.G.* Principles of formation and use of phytoclimatic scales. **Ukr. Phytocoe, J**, 1998; C; 1(10): 17–27. (In Ukrainian).
 21. *Protopopova V.V., Shevera M.V., Mosyakin S.L., Solomakha V.A., Solomakha T.D., Vasylyeva T.V., Petryk S.P.* Invasive species in flora of Northern Cis Black Sea area. Kyiv: Phytosociocentre Press, 2009. 56 p. (In Ukrainian).
 22. *Protopopova V.V., Shevera M.V., Chorney I.I., Tokaryuk V.V., Korzhan K.V.* The transformer species in flora of Bukovinian Cis-Carpathian. **Ukr. Bot. J**, 2010; 67(6): 852–864. (In Ukrainian).
 23. *Sîrbu C., Ferus P., Eliáš P. jun., Samuil C., Oprea A.* *Symphotrichum ciliatum* in Romania: trends of spread and invaded plant communities. **Open Life Sciences**, 2015; 10: 147–164.
 24. *Sîrbu C., Oprea A.* **Plante adventive în flora României**. Iași: Editura "Ion Ionescu de la Brad", 2011: 413–414.
 25. *Sîrbu C., Oprea A., Eliáš P. jun., Ferus P.* New contribution to the study of alien flora in Romania. **J. Plant Develop**, 2011; 18: 121–134.
 26. *Shyshkin B.K.* Genus *Brachyactis* Ledeb. In Flora of USSR. Vol. 25 / Ed. B.K. Shyshkin. Moscow; Leningrad: Institute of Botany, Acad. Sc. USSR, press, 1959: 631 p. (In Russian).
 27. *Solomakha V.A.* Syntaxonomy of vegetation of Ukraine. Third approximation. Kyiv: Phytosociocentre Press, 2008. 296 p. (In Ukrainian).
 28. *Syrenko I.P.* Creation of Databases for Floristic and Phytocoenologic Researches. **Ukr. Fitotsen. Zb**, 1996; A (1): 9–11.
 29. *Tichy L.* JUICE, software for vegetation classification. **J. Veg. Sci**, 2002; 13: 451–453.
 30. *Tokarska-Guzik B., Dajdok Z., Zajac M., Zajac A., Urbisz A., Danielewicz W., Hołdyński C.* Rośliny obcego pochodzenia w Polsce ze szczególnym uwzględnieniem gatunków inwazyjnych. Warszawa, 2012. 198 p.
 31. *Tucharz M., Nobis A., Nobis M.* Chromosome data – a useful tool in taxonomy of *Symphotrichum ciliatum*. **Acta Biol. Cracov**, 2011; Botan; 53(1): 117–119.
 32. *Tzvelev N.N.* Genus *Brachyactis* Ledeb. In: **Flora of European Part of USSR**. Vol. 7 / ed. by N.N. Tzvelev. Sankt-Petersburg: Nauka Press, 1994. 195 p. (In Russian).

**ХОРОЛОГІЧНІ ТА ЕКОЛОГО-ЦЕНОТИЧНІ ОСОБЛИВОСТІ
SYMPHYOTRICHUM CILIATUM (LINDL.) NESOM (ASTERACEAE)
У БУКОВИНСЬКОМУ ПЕРЕДКАРПАТТІ**

А. І. Токарюк¹, І. І. Чорней¹, В. В. Буджак¹, В. В. Протопопова², М. В. Шеєра^{3,2}

¹ Чернівецький національний університет імені Юрія Федьковича
вул. Федьковича, 11, Чернівці 58022, Україна
e-mail: iichorney@ukr.net, vbudzhak@gmail.com

² Закарпатський угорський інститут імені Ференца Ракоці II
пл. Кошута, 6, Берегове, Закарпатська обл. 90200, Україна
e-mail: protopopova.vira@gmail.com

³ Інститут ботаніки ім. М.Г. Холодного НАН України
вул. Терещенківська, 2, Київ 01004, Україна
e-mail: shevera.myroslav@gmail.com

Представлено результати вивчення поширення та еколого-ценотичної приуроченості адвентивного виду *Symphyotrichum ciliatum* (Lindl.) Nesom (*Asteraceae*) у Буковинському Передкарпатті (Україна). Виявлено нові локалітети виду в регіоні, в т.ч. у складі оселищ Natura 2000.

Уперше з'ясовано бальні показники провідних едафічно-кліматичних факторів виду. Встановлено, що вид вирізняється здебільшого стенотопною екологічною амплітудою. Відмічений у складі угруповань асоціації *Eragrostio-Amarantheum albi* Morariu 1943 класу *Digitario sanguinalis-Eragrostietea minoris* Mucina, Lososová et Šilc in Mucina et al. 2016, асоціацій *Lolio-Plantaginetum majoris* (Linkola 1921) Beger та *Polygonetum arenastri* Gams 1927 класу *Polygono-Poetea annuae* Rivas-Mart. 1975, асоціацій *Odontito-Ambrosietum artemisiifoliae* Mucina 1993 та *Convolvulo-Agropyretum repentis* Felföldy 1943 класу *Artemisietea vulgaris* Lohmeyer et al. in Tx. ex von Rochow 1951, які приурочені до мезофітних, гемігідроконтрастотрофільних, субацидотрофільних, семіевтрофних, гемікарбонатотрофільних, гемінітрофільних умов.

Ключові слова: *Symphyotrichum ciliatum*, *Asteraceae*, поширення, еколого-ценотична приуроченість, Буковинське Передкарпаття, Україна

Одержано: 16.08.2017