

8. Технічні умови «Набори продуктів польові» ТУ У 10.8-00034022-153:2016.

9. Русаков В. Коллективные и индивидуальные пайки для военнослужащих ВС США. Зарубежное военное обозрение. 2014. № 3. С. 29-35.

10. Дорошевич В.И., Мошчик К.В., Ширко Д.И. Гигиеническая характеристика организации питания в армиях некоторых зарубежных стран. Военная медицина. 2016. № 1. С. 47-52.

11. Ширко Д.И. Гигиеническая оценка фактического питания военнослужащих. Военная медицина. 2015. № 3. С. 61-62.

REFERENCES

1. *Kosheliev N.F. & Mikhaylov V.B.* Gigiena pitaniia voisk [Nutrition Hygiene for Troops]. Leningrad ; 1993 : 257 p. (in Russian).

2. *Simakhina H.O. & Ukrainets A.I.* Naukovi pratsi Natsionalnoho universytetu kharchovykh tekhnolohii. 2016 ; 22 (1) : 192-200 (in Ukrainian).

3. NATO/RTO Nutrition Science and Food Standards for Military Operations (Nutrition et normes d'alimentation pour les opérations militaires) : Final Report of RTO Task Group RTG-154. NATO ; 2010. 286 p.

4. *Doroshevich V.Y. & Shyrko D.Y.* Voennaia meditsina. 2016 ; 2 : 78-82 (in Russian).

5. Pro vnesennia zmin do norm kharchuvannia viiskovosluzhbovt-siv Zbroinykh Syl ta inshykh viiskovykh formuvan : Postanova KМУ vid 22 lypnia 2016 r. № 470 [On the Introduction of the Changes into the Nutrition Norms of the Servicemen of the Armed Forces and Other Armed Formations: Resolution of the CMU, July 22, 2016 № 470]. Available at : <http://zakon3.rada.gov.ua/laws/show/470-2016-%D0%BF> (in Ukrainian).

6. *Pokrovskii A.A. (ed.)* Khimichieskiy sostav pishchevykh produktov [Chemical Composition of Food Products]. Moscow ; 1977 : 228 p. (in Russian).

7. *Ukrainets A.I., Simakhina H.O. & Naumenko N.V.* Naukovi pratsi Natsionalnoho universytetu kharchovykh tekhnolohii. 2015 ; 21 (3) : 209-215 (in Ukrainian).

8. Tekhnichni umovy «Nabory produktiv poliovi» [Technical Requirements "Field Food Sets"] : ТУ У 10.8-00034022-153:2016 (in Ukrainian).

9. *Rusakov V.* Zarubiezhnoie voiennoie obozreniie. 2014 ; 3 : 29-35 (in Russian).

10. *Doroshevich V.I., Moshchik K.V. & Shirko D.I.* Voennaia meditsina. 2016 ; 1 : 47-52 (in Russian).

11. *Shirko D.I.* Voennaia meditsina. 2015 ; 3 : 61-62 (in Russian).

Надійшло до редакції 18.09.2016

ПОШИРЕНІСТЬ ЗДОРОВОГО СПОСОБУ ЖИТТЯ СЕРЕД УКРАЇНСЬКИХ СТУДЕНТІВ

Якименко І., Цибулін О., Шаповалов Є.

HEALTHY LIFESTYLE BEHAVIORS AMONG UNIVERSITY STUDENTS IN UKRAINE

H

Healthy lifestyle behaviors of young people influence their health during future life and thus affect public health of future generation of adults. Healthy lifestyle characteristics include nutritious diet, sufficient physical activity, non-smoking, alcohol restriction, safe sexual behavior, and behaviors that do not contribute to injuries [1]. Obviously, adherence to healthy lifestyle depends on socioeconomic factors, cultural background, age and gender of the individual. In 2015, the worldwide population of young people aged between 10 and 24 years is estimated to be about 1.8 billion, representing 25% of the global population [2]. The size of this part of population makes youth's health status important not only as a determinant of health of future adult generation, but also for global social

¹YAKYMENKO I.,

²TSYBULIN O.,

¹SHAPOVALOV YE.

¹National University of Food Technologies, Kyiv, Ukraine
²Bila Tserkva National Agrarian University, Bila Tserkva, Ukraine

УДК: 612.014.482

Keywords: health, lifestyle, youth, environment, smoking, alcohol, diet, physical activity.

РАСПРОСТРАНЕННОСТЬ ЗДОРОВОГО ОБРАЗА ЖИЗНИ СРЕДИ УКРАИНСКИХ СТУДЕНТОВ

¹Якименко І., ²Цыбулин А., ¹Шаповалов Е.

¹Национальный университет пищевых технологий

²Белоцерковский национальный аграрный университет

Цель работы. Изучить приверженность студентов украинских университетов здоровому образу жизни.

Материалы и методы. Методом добровольного анонимного анкетирования проведена оценка приверженности студентов двух университетов киевского региона (n=211) здоровому способу жизни.

Результаты. Выявлено, что среди студентов возрастом от 17 до 21 года, излишний вес имеют 7,1% студентов, значительная часть молодежи, особенно парни (74,7%), систематически занимаются физическими упражнениями. Вместе с тем, 25,6% студентов курят и 61,6% употребляют алкоголь (хотя бы иногда). Менее половины опрошенных употребляют овощи и фрукты в достаточном количестве и молочные продукты ежедневно. Более 70% студентов допускают негативное влияние окружающей среды на их здоровье и признают недостаточное материальное обеспечение для лучшей приверженности здоровому способу жизни.

Выводы. В целом студенческая молодежь в Украине имеет большую приверженность к здоровому образу жизни, чем старшее поколение, но курение и потребление алкоголя остаются серьезной угрозой для ее здоровья.

Ключевые слова: здоровье, образ жизни, молодежь, окружающая среда, курение, алкоголь, питание, физические нагрузки.

© Якименко І., Цибулін О., Шаповалов Є. СТАТТЯ, 2017.

**ПОШИРЕНІСТЬ ЗДОРОВОГО СПОСОБУ ЖИТТЯ
СЕРЕД УКРАЇНСЬКИХ СТУДЕНТІВ**

¹Якименко І., ²Цибулін О., ¹Шаповалов Є.

¹Національний університет харчових технологій

²Білоцерківський національний аграрний університет

Мета роботи. Дослідити прихильність студентів українських університетів до здорового способу життя.

Матеріали і методи. Методом добровільного анонімного анкетування проведено оцінку прихильності студентів двох університетів київського регіону (n=211) до здорового способу життя.

Результати. Виявлено, що серед студентів віком 17-21 років надмірну вагу мають 7,1%, значна частина молоді, особливо хлопці (74,7%), систематично займаються фізичними вправами.

Разом з тим, 25,6% студентів палять і 61,6% вживають алкоголь (принаймні іноді). Менше половини опитаних студентів споживають фрукти та овочі у необхідній кількості та молочні продукти щодня. Понад 70% опитаних припускають негативний вплив довкілля на їхнє здоров'я та визнають недостатню матеріальну забезпеченість для кращої прихильності до здорового способу життя.

Висновки. Загалом студентська молодь в Україні має кращу прихильність до здорового способу життя, ніж старше покоління, але паління та вживання алкоголю залишаються серйозною загрозою для її здоров'я.

Ключові слова: здоров'я, спосіб життя, молодь, довкілля, паління, алкоголь, харчування, фізичне навантаження.

and economic development [3]. And analysis of healthy lifestyle behaviors in this cohort of population is closely connected to the issue. It is known, that healthy lifestyle behaviors significantly reduce the level of diseases and as a result decrease all-cause mortality of population [1]. On the other hand, unhealthy lifestyle behaviors such as bad dietary habits and a lack of physical activity are well known risk factors for cardiovascular disease, type 2 diabetes and some cancers [4].

Ukraine has one of the worst public health indices in the region [5]. The prevalence of poor health was previously reported to be much higher for young adults in the former Soviet Union regions than in Western Europe and North America (odds ratio 4.26, 95% CI 1.77–10.24) [6]. Indeed, according to the Parliament hearing in Ukraine, in 2011, 45% of young males and 35% of young females smoked; and 68% of boys and 64% of girls

drunk alcohol [7]. For comparison, in the USA, only 9.6% of adolescents aged 12-17 consume alcohol and 20% smoke, which resulted from rigorous public health efforts [8]. Meanwhile, we hardly can find modern data on health behaviors among Ukrainian university students as the most active and educated part of youth.

Our study aims to survey Ukrainian university students on their adherence to healthy lifestyle behaviors.

Materials and methods.

The survey was carried out during March-June 2016 among volunteer students at two universities of Kyiv region, central part of Ukraine. A printed questionnaire included 24 questions on age, gender, height and weight of individual, smoking, consumption of alcohol, consumption of fruits and vegetables, daily milk/dairy product consumption, having breakfast daily, using a computer and a cell phone, environmental impacts, and economic conditions of the individual.

Totally 237 students were surveyed, and 211 questionnaires (89.4%) were selected as filled in understandable without missing any questions. Thus, they were used for subsequent analysis.

Body mass index (BMI) was derived from weight and height of respondents according to the formula $BMI = \text{Weight} / \text{Height}^2$ (kg/m²). The normal BMI was restricted in our analysis 18 to 25 kg/m² according to the World Health Organization (WHO) recommendations. Smoking was assessed in Yes/No question with additional question on quantity of cigarettes per day for those who smoked. Consumption of alcohol was assessed in Yes/No question with further specification for those who drink: less than 2 bottles of beer or 2 glasses of wine per week; more than that amount of alcohol per week; and more than that amount of alcohol per day. Proper fruit and non-starchy vegetable consumption was assessed as 400 g per day or more, according to the WHO

Table 1

Healthy lifestyle behaviors among Ukrainian university students

Indices	All students (n=211)		Females (n=104)		Males (n=107)		RR Females / Males	CI, 95%
	n	%	n	%	n	%		
Normal BMI	187	88,62	93	89,42	94	87,85	1,01	0,92 - 1,12
Non-smoking	157	74,4	85	81,73	72	67,28	1,21	1,03- 1,42
Non-drinking alcohol	81	38,38	51	49,03	30	28,03	1,74	1,21- 2,51
Rare drinking alcohol	98	46,44	49	47,11	53	49,53	0,96	0,73- 1,28
Proper physical activity	139	65,87	59	56,73	80	74,76	0,75	0,62- 0,92
Proper consumption of fruits and vegetables	100	47,39	64	61,53	37	34,57	1,77	1,31- 2,40
Daily milk/dairy products consumption	96	45,49	50	48,07	47	43,92	1,09	0,81-1,46
Daily breakfast	123	58,29	62	59,61	61	57,01	1,04	0,83-1,31


recommendations. Daily milk and dairy product consumption, and daily breakfast eating were estimated in Yes/No questions. Proper physical activity was recognized when individual spent at least 2.5 hours on physical exercises per week.

There were calculated total indices of surveyed population and assessed differences between gender subgroups, as well as between subgroups with normal BMI and those who overweight. Differences between subgroups were assessed due to relative risk approach (RR, CI 95%).

Results and discussion.

The students surveyed were 17-21 years old, 49.3% of females, mostly freshmen with biological or environmental science background. Most students had normal BMI (table 1), while 7.1% of them were overweight with BMI ranged from 25.1 to 32.1 kg/m², and 4.3% were underweight with BMI ranged from 15.9 to 17.4 kg/m². Overweight male students comprised 10.3% of male respondents and overweight female students comprised 3.8% of female respondents.

Three quarters of students did not smoke, and females smoked significantly less often than males, 19.3% against 32.7% (table 1). About half of females but only 28% of males did not consume alcohol at all. And about half of respondents, both young women and men consumed alcohol in amount less than 2 bottles of beers per week.



ГІГІЕНА ДІТЕЙ ТА ПІДЛІТКІВ

Significantly more male students practiced proper physical activity than females, 74.8% against 56.7%, respectively. Instead, young women had almost twice higher percent of those who consumed proper amount of fruits and vegetables than young men. But totally only 47.5% of students consumed proper amount of fruits and vegetables daily. And only 45.5% of respondents consumed milk or dairy products daily. Daily breakfast had 58.3% of respondents.

As for the possible impact of modern technologies on students health, 45.1% of respondents talked on a cell phone up to 1 hour or more per day, and 58.5% of students used a computer up to 3 hours or more per day.

Among socio-economic aspects of respondents' health behaviors, there were discovered that 33.3% of students did not know about official healthy lifestyle recommendations, including the WHO recommendations. Significant part of respondents (77.1%) recognized that a bad physical environment could affect their

health, and 67.5% of them admitted that the Chernobyl accident / Chernobyl radiation could affect their health. And most students (70.4%) thought that they need better economic status for improving their healthy lifestyle adherence.

The analysis of healthy lifestyle indices in overweight cohort of students against their peers with normal BMI did not reveal statistically significant differences between two subgroups (table 2). Nevertheless, the subgroup of overweight students had twice less amount of individuals who did not drink alcohol at all and 31.5% less individuals with proper consumption of fruits and non-starchy vegetables.

Generally, our analysis demonstrated that Ukrainian university students have a broad variety in adherence to different healthy lifestyle characteristics. The positive indicator was a high level of young people with normal BMI. According to our findings, only 7.1% of respondents were overweight. This result corresponds to data of Tsos et al., 2014 [9], who found 8.7% of

Health behaviors among overweight Ukrainian university students as compared with normal weight students

Table 2

Indices	Overweight (n=15)		Normal weight(n=187)		RR	CI, 95%
	n	%	n	%		
BMI	25,1-32,1		18 - 25			
Non-smoking	11	73,33	139	74,33	0,98	0,71- 1,35
Non-drinking alcohol	3	20	75	40,1	0,49	0,17- 1,39
Rare drinking alcohol	9	60	83	44,38	1,35	0,86- 2,10
Proper physical activity	9	60	128	68,44	0,87	0,57- 1,34
Proper consumption of fruits and vegetables	5	33,33	91	48,66	0,68	0,32- 1,42
Daily milk/dairy products consumption	7	46,66	87	46,52	1	0,57 - 1,760
Daily breakfast	10	66,66	107	57,21	1,16	0,79- 1,70

overweight students in Ukraine in 2013. On the other hand, according to Chagarna and Andreeva, 2014 [10], the risk of overweight increased among older population in Ukraine. For example, in 2000, among those under 40 years old, 16% of men and 12% of women were overweight, while among 40 and more years old already 47% of men and 60% of women were overweight in Ukraine. Thus, the obvious problem here is to promote healthy lifestyle behaviors and control the BMI from young to older ages. Overweight and obesity is a serious health problem for many developed country nowadays. For example, in 2000 only 40% of adult Americans had normal weight [11], and according to 2003-2006's national survey only 15.6% Americans aged 20-39 had normal body fat percentage [12]. While Ukrainian population seems do not bear a pattern of obesity epidemic found in Western countries, the overweight of older people is a significant risk factor for the older population in the country.

Optimistic result of our analysis is a high percentage of male students practicing sufficient level of physical activity systematically, at least 2.5 h per week, while female population was significantly less physically active. These data differ from the findings of the other survey of 2000 [10], which indicated that about 71% of adult population in Ukraine did not practiced any physical activities. It seems that we need the urgent national strategy on promotion of proper physical activity among older people to keep positive trend among student youth.

Improper level of fruit and vegetable consumption among Ukrainian students obviously has both social and economic roots, as every third student did not know about healthy behavior recommendations and most of the respondents recognized their insufficient economic status for proper healthy lifestyle behaviors.

Comparatively low level of daily milk consumption among students combined with often insufficient consumption of fruits and vegetables, especially among young men, can lead to a significant restriction of student diet in important nutrients like vitamins, minerals and phytochemicals. Our findings demonstrate that Ukrainian students 17-21 years old share the problem of inadequate nutrition with European peers. For example, according to the HELENA study, adolescents from the EU cities aged 12.5-17.5 ate a half of the recommended amount of fruits and vegetables and less than two thirds of recommended amount of milk and dairy products [13].

Intensive usage of cell phones and computers by significant part of university students can contribute to their health risks as well. For example, previously we demonstrated that up to 40% of Ukrainian students had physical discomfort and/or pain in the head or ear during long-term cell phone talks [14].

Our analysis confirmed that smoking and drinking alcohol are serious risk factors for Ukrainian population. While surveyed students were only 17-21 years old, quarter of them smoked and almost two thirds consumed alcohol at least sometimes. Moreover, among male students 32.7% smoked and 72% consumed alcohol (at least up to two bottles of beer per week). For comparison, among adult population in Ukraine, 54.3% smoked and 79% consumed vodka (at least rare) in 2000 [10].

Our study has some limitations. As we used self-reported data, it could be some misinterpretations in the answers, eg,

students might overestimate or underestimate their daily portions of fruits and vegetables, or precise level of physical activity. Also, we did not assess the level of food insecurity as a lack of consistent access to healthy safe food among the students. For example, recently there was reported that up to 37% of American college freshmen can be classified as food-insecure [15]. We can't interpret if the low percentage of Ukrainian students with excessive weight was due to proper diet and sufficient physical activity or food insecurity/malnutrition contributed in the issue. These issues will be a next stage of our research.

Conclusions. Ukrainian university students have much better adherence to healthy lifestyle behaviors than adult population in Ukraine. Low percentage of overweight students (7.1%) and rather high level of physically active students, especially among males (74.7%) are optimistic tendencies.

On the other hand, 25.6% of students smoked and 61.6% consumed alcohol (at least rare). Only 47.4% of surveyed students consumed sufficient amount of fruits and vegetables and only 45.5% consumed milk and dairy products daily. Most students recognized negative impacts of the physical environment, including Chernobyl radiation, on their health, and assumed their insufficient economic status for better adherence to healthy lifestyle.

REFERENCES

1. Loef M., Walach H. The combined effects of healthy lifestyle behaviors on all cause mortality: a systematic review and meta-analysis. *Preventive medicine*. 2012. Vol. 55(3). P. 163-170.
2. United Nations, Department of Economic and Social Affairs. *World Population Prospects: The 2015 Revision*. New York, 2015. 66 p.
3. Gore F.M., Bloem P.J., Patton G.C. et al. Global burden of disease in young people aged 10-24 years: a systematic analysis. *The Lancet*. 2011.

Vol. 377(9783). P. 2093-2102.

4. WHO/FAO Diet, nutrition and the prevention of chronic diseases. WHO Technical Report Service No 916. 2003.

5. МОЗ України.

Національна стратегія реформування системи охорони здоров'я в Україні на період 2015 – 2020 років. Київ, 2015. 41 с.

6. Witvliet M.I., Arah O.A., Stronks K., Kunst A.E. Examining self-rated health of young central and eastern Europeans in the context of other world regions. *European Journal of Public Health*. 2014. Vol. 24(2). P. 314-321.

7. Рекомендації парламентських слухань про становище молоді в Україні "Молодь за здоровий спосіб життя". Київ, 2011. Режим доступу : <http://zakon0.rada.gov.ua/laws/show/2992-vi>

8. Center for Behavioral Health Statistics and Quality. Key substance use and mental health indicators in the United States: Results from the 2015 National Survey on Drug Use and Health (HHS Publication No. SMA 16-4984, NSDUH Series H-51). 2016. Retrieved from <http://www.samhsa.gov/data/>

9. Tsos A., Bergier B., Bergier J. Physical activity, physical development and eating habits within the lifestyle of students from Ukraine. *Health Problems of Civilization*. 2014. Vol. 8 (4). P. 46-53.

10. Chagarna N., Andreeva T.I. Lifestyle correlates of overweight and obesity among the population of Ukraine. *Tobacco Control and Public Health in Eastern Europe*. 2014. Vol. 4 (1). P. 15-28.

11. Reeves M.J., Rafferty A.P. Healthy lifestyle characteristics among adults in the United States, 2000. *Archives of Internal Medicine*. 2005. Vol. 165(8). P. 854-857.

12. Loprinzi P.D., Branscum A., Hanks J., Smit E. Healthy lifestyle characteristics and their joint association with cardiovascular disease biomarkers in US adults. *Mayo Clin Proc*. 2016. Vol. 91(4). P. 432-442.

13. Ottevaere C., Huybrechts I., Benser J., De Bourdeaudhuij I., Cuenca-Garcia M., Dallongeville J. et al. Clustering patterns of physical activity, sedentary and dietary behavior among European adolescents: The HELENA study. *BMC Public Health*. 2011. Vol. 11(1). P. 328.

14. Yakymenko I., Mor O., Tsybulin O., Kolesnik Ya., Kyrylenko S., Sidorik E. Subjective symptoms in young cell phone users in Ukraine. *Довкілля та здоров'я*. 2015. №2. С. 40-43.

15. Bruening M., Brennhofer S., van Woerden I., Todd M., Laska M. Factors Related to the High Rates of Food Insecurity among Diverse, Urban College Freshmen. *Journal of the Academy of Nutrition and Dietetics*. 2016. Vol. 116 (9). P. 1450-1457.

REFERENCES

1. *Loef M. & Walach H.* *Preventive Medicine*. 2012 ; 55(3) : 163-170.

2. United Nations ; Department of Economic and Social Affairs. *World Population Prospects: The 2015 Revision*. New York ; 2015 : 66 p.

3. *Gore F.M., Bloem P.J., Patton G.C., Ferguson J., Joseph V., Coffey C., Sawyer S.M. & Mathers CD.* *The Lancet*. 2011 ; 377(9783) : 2093-2102.

4. WHO/FAO Diet, Nutrition and the Prevention of Chronic Diseases. WHO Technical Report Service No 916. 2003.

5. Ministry of Public Health of Ukraine. *Natsionalna stratehiia reformuvannia systemy okhorony zdorovia v Ukraini na period 2015 – 2020 rokiv* [National Strategy of Reforming the Health Care System in Ukraine on the Period 2015 – 2020]. Kyiv; 2015 : 41 p. (in Ukrainian).

6. *Witvliet M.I., Arah O.A., Stronks K. & Kunst A.E.* *European Journal of Public Health*. 2014 ; 24(2) : 314-21.

7. *Rekomendatsii parlamentskikh slukhan pro stanovyshe molodi v Ukraini "Molod za zdorovi sposib zhyttia"* [Parliament Attendant

on the Youth Status in Ukraine "Youth for Healthy Life Style"]. Kyiv; 2011. URL: <http://zakon0.rada.gov.ua/laws/show/2992-vi> (in Ukrainian).

8. Center for Behavioral Health Statistics and Quality. Key Substance Use and Mental Health Indicators in the United States: Results from the 2015 National Survey on Drug Use and Health (HHS Publication No. SMA 16-4984, NSDUH Series H-51). 2016. URL : <http://www.samhsa.gov/data/>

9. *Tsos A., Bergier B. & Bergier J.* *Health Problems of Civilization*. 2014 ; 8(4) : 46-53.

10. *Chagarna N. & Andreeva T.I.* *Tobacco Control and Public Health in Eastern Europe*. 2014 ; 4(1) : 15-28 (in Russian).

11. *Reeves M.J. & Rafferty A.P.* *Archives of Internal Medicine*. 2005 ; 165(8) : 854-857.

12. *Loprinzi P.D., Branscum A., Hanks J. & Smit E.* *Mayo Clinic Proceedings*. 2016 ; 91(4) : 432-442.

13. Ottevaere C., Huybrechts I., Benser J., De Bourdeaudhuij I., Cuenca-Garcia M., Dallongeville J. et al. *BMC Public Health*. 2011 ; 11(1) : P. 328.

14. *Yakymenko I., Mor O., Tsybulin O., Kolesnik Ya., Kyrylenko S. & Sidorik E.* *Dovkillia ta zdorovia* [Environment and Health]. 2015 ; 2 : 40-43.

15. *Bruening M., Brennhofer S., van Woerden I., Todd M. & Laska M.* *Journal of the Academy of Nutrition and Dietetics*. 2016 ; 116 (9) : 1450-1457.

Надійшло до редакції 18.09.2016