

Beata Gavurova
PhD (Economics), Associate Professor,
Technical University of Kosice,
32 Nemcovej Str., Kosice,
040 01, Slovak Republic
beata.gavurova@tuke.sk





Efficiency of day surgery in Slovak regions during the years 2009-2014

Abstract. The paper deals with CCR, BCC and FDH efficiency analyses results of valid available indicators of eight Slovak regions with one day surgery facilities both for adult and paediatric patients in the period of 2009-2014, where the input is the number of beds and the output is the number of patients. In 2014, Banska Bystrica region was efficient from the viewpoint of CCR model of paediatric patients. According to the BCC model, there were four efficient cases: Trnava - in 2009, Banska Bystrica - in 2014 and Presov - two years running (2013 and 2014). In the case of CCR model regarding adult patients, Zilina region was efficient in 2009. On the basis of the BCC DEA model, six efficient cases have been determined: Trnava and Zilina regions - in 2009, Zilina region - in 2010, Banska Bystrica - in 2012 and Bratislava region - in 2012 and 2014. The results of the analysis present a valuable platform for the creators of health and social policies as well as for other stakeholders of health system in Slovakia that will transform into correct stabilization and regulatory mechanisms of health and social policies.

Keywords: Day Surgery; Hospital; Efficiency; Healthcare; Data Envelopment Analysis; Slovakia

JEL Classification: H51; C61

Acknowledgement. The work was supported by the VEGA Project No. 1/0986/15 «Proposal of the dimensional models of the management effectiveness of ICT and information systems in health facilities in Slovakia and the economic-financial quantification of their effects on the health system in Slovakia».

Our thanks go to the National Health Information Center of Slovakia (http://www.nczisk.sk) for providing the Slovak day surgery database, as well as for a long-term support of our research activities within health policy.

DOI: http://dx.doi.org/10.21003/ea.V159-17

Беата Гавурова

кандидат економічних наук, доцент, факультет економіки, Технічний університет у Кошице, Словацька Республіка Самуел Корони

кандидат економічних наук, доцент, факультет економіки, Університет Матея Бела, Банська Бистриця, Словацька Республіка Ефективність відділень денної хірургії в регіонах Словацької Республіки в період 2009–2014 років

Анотація. У статті розглядаються ССЯ, ВСС та FDH моделі аналізу середовища функціонування (АСФ, англ. мовою – Data Envelopment Analysis (DEA)), побудовані на основі використання наявних достовірних показників, що характеризують відділення денної хірургії для дорослих та дітей у восьми регіонах Словацької Республіки в 2009–2014 рр. У якості вхідної змінної був обраний показник «кількість ліжок», а в якості вихідної – «чисельність пацієнтів». З огляду на ССЯ модель, педіатричні відділення регіону Банська Бистриця були ефективними у 2014 році. За результатами застосування ВСС моделі було виділено чотири випадки ефективності відділень: в регіоні Трнава – у 2009 році, у регіоні Банська Бистриця – у 2014 році, а також у регіоні Прешов протягом 2013 та 2014 років. У випадку ССЯ моделі, яка застосовувалася для визначення ефективності відділень для дорослих пацієнтів, ефективними були відділення в регіоні Жиліна в 2009 році. На основі моделі ВСС було отримано шість випадків ефективності відділень: у регіонах Трнава та Жиліна – у 2009 році, у регіоні Жиліна – у 2010 році, у регіоні Банська Бистриця – у 2012 році та в Братиславському регіоні у 2012 та 2014 роках. Результати проведеного аналізу є платформою для розробки політики охорони здоров'я та соціального захисту в Словаччині, формуючи основу для створення ефективних механізмів стабілізації та регулювання у цих сферах.

Ключові слова: відділення денної хірургії; ефективність; охорона здоров'я; аналіз середовища функціонування.

Беата Гавурова

кандидат экономических наук, доцент, факультет экономики, Технический университет в Кошице, Словацкая Республика **Самуел Корони**

кандидат экономических наук, доцент, факультет экономики,

Университет Матея Бела, Банска Быстрица, Словацкая Республика

Эффективность отделений дневной хирургии в регионах Словацкой Республики в период 2009-2014 годов

Аннотация. В статье рассматриваются ССR, ВСС и FDH модели анализа среды функционирования (АСФ, на англ. языке – Data Envelopment Analysis (DEA)), построенные на основе использования имеющихся достоверных показателей, характеризующих отделения дневной хирургии для взрослых и детей в восьми регионах Словакии в период 2009–2014 гг. В качестве входной переменной был выбран показатель «количество коек», а в качестве выходной – «численность пациентов». С точки зрения модели ССR, педиатрические отделения Банскобыстрицкого региона были эффективными в 2014 году. По результатам применения модели ВСС были выделены четыре случая эффективности отделений: в регионе Трнава – в 2009 году, в регионе Банска Быстрица – в 2014 году, а также в регионе Прешов на протяжении 2013 и 2014 годов. В случае с моделью ССR, которая применялась для отделений взрослых пациентов, были эффективны отделения в Жилинском регионе в 2009 году. На основе модели ВСС DEA было определено шесть случаев эффективности отделений: в регионах Трнава и Жилина – в 2009 году, в Жилинском регионе – в 2010 году, в регионе Банска Быстрица – в 2012 году, а также в Братиславском регионе – в 2012 и в 2014 годах. Результаты проведенного анализа представляют собой ценную платформу для разработки политики здравоохранения и социальной защиты в Словакии, формируя основу для создания корректных механизмов стабилизации и регулирования в этих сферах.

Ключевые слова: отделения дневной хирургии; эффективность, здравоохранение; анализ среды функционирования.

1. Introduction & Brief Literature Review

In Slovakia, day surgery has been developing since 2000. Its development is connected with a formation of the Slovak Association of Day Surgery that supported the formation of a legal framework to create day surgery workplaces. However, the ratio of day surgery (approximately 7% out of the total number of realised surgeries in comparison to 40-70% ratio abroad) is very low in spite of the fact that the Ministry of Health of the Slovak Republic supports it significantly (Gavurova & Hyranek, 2013; Gavurova et al., 2014) [1-2]. Day surgery has been operating successfully abroad for decades and it has also been used by older patients as it is interconnected with subsequent nursing care in many countries.

The fundamental characteristic of day surgery operation states that it is an operation, which is relatively patient-friendly with regard to the patient's body and it does not represent any extremely high post-operative burden for the patient's body (Jarrett & Staniszewski, 2006)[3]. It is also necessary to take into consideration the level of operational invasiveness, the relevant complications, the risk of blood loss, the estimated post-operative pain or the time which is necessary to perform day surgery operations (Kroneman, 2001; Soltes et al., 2011a; Soltes & Gavurova, 2014) [4-6]. Threrefore, minimally invasive procedures, particularly in general surgery, represent a viable option (Soltes et al., 2010) [7]. Furthermore, there are also patients' characteristics that may promote or prohibit day surgery preference for the given procedure (Soltes & Radonak, 2014; Popa et al., 2015) [8-9]. Day surgery decreases a risk of nosocomial infections that extend the treatment period and increases treatment costs. Day surgery differs within each country. This may be caused by many reasons of both economic and non-economic character (Soltes, 2011b; Szabo & Sidor, 2014; Szczygiel et al., 2014) [10-12]. Reimbursement mechanisms represent an important aspect in the individual countries (Soltes & Gavurova, 2015; Bem et al., 2014; Bem & Michalski, 2015) [13-15].

In the context of the given facts, this article provides another partial view of health system efficiency by using a chosen method of evaluating the day surgery efficiency in Slovakia. We obtained the data on the basis of contractual cooperation with the National Health Information Center.

2. Purpose

The main purpose of our paper was to do day surgery efficiency analyses in the Slovak regions (NUTS III level) in the period of 2009-2014 (the relevant day surgery data have been collected on the basis of the Ministry of Health of the Slovak Republic Guidelines since 2009). Our main analytical methods were DEA (CCR, BCC and FDH) input oriented models. We assume that the number of patients is already defined and it is not rational to increase it due to different medical, personal and technological requirements of day surgery therapy patients.

3. Data and Research Methods

«The Annual Report of Day Healthcare» is the data source for our DEA analysis of day surgery healthcare efficiency (The Ministry of Health of the Slovak Republic). It was offered to us by the National Health Information Center of Slovakia.

Slovak regions with day surgery facilities are our research objects. We have analysed suitable available indicators of day surgery efficiency during the years 2009-2014. The number of day surgery patients in two age groups (paediatric patients under 18 years old and adult patients of 19 or more years old) was the output. The corresponding number of both paediatric and adult beds was used as the input.

Besides descriptive statistics, we have used three input oriented DEA models (CCR, BCC a FDH) (Cooper et al., 2007) [16]. The CCR model with constant returns to scale was the

first DEA model (Charnes et al., 1978) [17]. The BCC model with variable returns to scale was published later (Banker et al., 1984) [18]. The linear combination of units (in our case the units are Slovak regions in corresponding years) is used to compute optimum values of inputs and outputs in the DEA models. It evokes the question of reality of such «regions» in our case.

DEA is a non-parametric approach for frontier estimation. Basic models are discussed in works of many authors and applied in different areas. In the Slovak Republic, the DEA models have been used to measure the efficiency of financial institutions (Kocisova, 2012) [19] or the efficiency of the agricultural sector (Kocisova, 2015) [20].

A different approach to the problem of efficiency was published in 1984 (Deprins et al., 1984; Cooper et al., 2007) [16-17]. It does not require the basic classic condition for the production models - convexity of production sets. Another important difference in comparison with other DEA models is that evaluated units are compared only to real efficient units, not to their linear combinations. Optimal (dominant) units are on the so-called Free Disposal Hull (FDH) of production set. Its definition is very simple:

$$FDH(x_0, y_0) = \{(x, y) : x \ge x_0; y \le y_0\},$$
 (1)

Production units are compared with optimum units on FDH in such a way that none of the units has a smaller input and a larger output. Then, for all $(x,y) \in FDH(x_0,y_0)$ at least one of the three conditions is valid $x>x_0, y=y_0; x=x_0, y< y_0; x>x_0, y< y_0$. The optimal unit produces the same output by a smaller input or a larger output by the same input or a larger output by a smaller input. Input oriented FDH model is then defined by:

$$\min \theta, X\lambda \leq \theta x_0, Y\lambda \geq y_0, \epsilon\lambda = 1, \lambda \in \{0,1\},$$
 (2)

where θ is the input efficiency, X and Y are input and output matrices and λ is the Boolean variable with values 0 or 1. Computational simplicity is an advantage of the FDH method. It is not necessary to use optimisation programming methods. Only pair-wise comparisons of units are needed. Possible existence of so-called slacks is a disadvantage of the FDH. However, it is also the problem of the classic radial CCR and BCC DEA models. In our case of one input and one output, the existence of possible slacks can be easily controlled visually. For statistical and graphical analyses, we have used the statistical system SPSS version 19. The software DEA solver was used to conduct the DEA analyses (Cooper et al., 2007) [10]. In next part, we briefly present statistical parameters of the analysed variables (day surgery indicators).

4. Results

a) Descriptive statistics results

There are eight self-government regions in Slovakia (NUTS III level). In the paper, we use their abbreviations from National Health Information Center of Slovakia: Banska Bystrica (BC), Bratislava (BL), Kosice (KI), Nitra (NI), Presov (PV), Trnava (TA), Trencin (TC) and Zilina (ZI). The available data cover a period of five years: 2009-2014 (then, for example, the case of Trencin region in 2012 is marked as TC12). Basic statistical parameters of examined available variables (the number of day surgery paediatric patients, the number of day surgery adult patients, the number of day surgery adult beds) are shown in Table 1. We used them in the efficiency analysis of Slovak regions (Mean = arithmetic mean, Std. Dev = standard deviation. The number of cases is 48 (eight Slovak regions in the period of six years).

The mean number of day surgery paediatric patients is 1,463. The minimum number (201) was in Trencin region in 2009, whereas the maximum number (3,377) was in Presov region in 2014.

The number of day surgery adult patients has a mean value of 15,410. The minimum number (4,233) was in Trnava region in 2009, whereas the maximum number (33,049) was in Bratislava region in 2014.

Tab. 1: Statistical parameters of day surgery in Slovak regions during the years 2009-2014

Variable	Mean	Std. Dev.	Minimum	Region	Maximum	Region
No. of paediatric patients	1,463	961.1	201	TC9	3,377	PV14
No. of adult patients	15,410	7,357.8	4,233	TA9	33,049	BL14
No. of paediatric beds	43	30.2	5	TA9	173	NI14
No. of adult beds	172	86.1	30	TA9	439	BL14

Source: Own calculations

The mean number of day surgery paediatric beds is 43 ranging from the minimum value of 5 in Trnava region in 2009 to the maximum value of 173 in Nitra region in 2014.

The mean number of day surgery beds for adults is 172 with their minimum number of 30 in Trnava region in 2009 and their maximum number of 439in Bratislava region in 2014. In the next part of this paper, we present the results of the application of the DEA methods to calculate the available day surgery indicators.

b) DEA analyses results We have a production dependence with one input (the number of day surgery beds) and one output (the number of day surgery patients). It is easy to depict it in the common scatter-plot where we can see relative positions of Slovak regions during the analysed time period in production possibility set between the number of day surgery patients (the vertical axis) and the corresponding number of day surgery beds (the horizontal axis). Figure 1 is shows a production set of paediatric day surgery efficiency problem with the CCR (constant returns to scale) and the BCC (variable returns to scale) production frontiers. The FDH production frontier is not depicted because the relevant software was unavailable we have not got software for it. From the viewpoint of the CCR model of paediatric day surgery, is efficient Banska Bystrica region was efficient in 2014 (BC14). A line from the point (0.0) goes through it and determines the CCR efficient production frontier. Other Slovak regions should shift to the CCR production frontier in the horizontal direction to the left to become input efficient. The BCC production frontier is piecewise linear function which goes through four BCC efficient cases.

An overview of the results of DEA analyses for paediatric day surgery is reflected in Table 2 which contains CCR, BCC and FDH efficiency scores. A reduction of day surgery beds that is needed for Slovak regions (with day surgery facilities) to become efficient is computed easily from efficiency scores, e.g. if efficiency is 0.7, then the relative reduction is 1 - 0.7 = 0.3 (30%). Of course, the efficient units have a 0% reduction. The most conservative is the CCR (with constant returns to scale) model. Usually, only one unit is efficient in its case. Then follows the BCC model (with variable returns to scale) with more than one efficient unit. The FDH model is the least conservative. It has even a larger proportion of efficient units.

Then, three regions follow, namely Zilina in 2014 (ZI14), Banska Bystrica in 2013 (BC13) and Kosice in 2012 (KI12) with almost the same reduction just above 50%. Nitra was relatively the «worst» region in 2014 (NI14) with its necessary bed reduction of large proportion equal to 96% (from 173 beds to only 6 beds). It is also clear from Figure 1. The region is in the lower right corner of production set and falls far behind the other regions. Furthermore, Nitra region has been at the end of the list of rank order through all the years.




Fig. 1: Production set of number of day surgery paediatric patients vs the number of day surgery paediatric beds with corresponding CCR and BCC DEA frontiers

Source: Own construction

efficiency analyses of paediatric day surgery in Slovak regions in the period of 2009-2014

nk Region CCR eff. Region BCC eff. Region FDH e

Tab. 2: Results of input oriented CCR, BCC a FDH DEA

Rank	Region	CCR eff.	Region	BCC eff.	Region	FDH eff.
1	BC14	1	TA9	1	PV14	1
2	ZI14	0.497	BC14	1	BC14	1
3	BC13	0.496	PV13	1	PV13	1
4	KI12	0.495	PV14	1	BC13	1
5	BL10	0.424	BC13	0.651	TA10	1
6	TA11	0.399	BL10	0.565	BC10	1
7	KI11	0.398	TA10	0.562	TA9	1
8	PV13	0.373	TA11	0.556	TA11	0.941
9	BL9	0.373	BC10	0.553	BL10	0.889
10	PV14	0.362	ZI14	0.518	BL11	0.885
11	BC12	0.345	KI12	0.502	BC9	0.800
12	ZI11	0.339	BC12	0.468	TA12	0.800
13	BC11	0.339	TA12	0.461	BL9	0.793
14	ZI12	0.331	BC11	0.456	BL14	0.767
15	TA12	0.325	BL9	0.442	BC12	0.762
16	BC10	0.313	BL11	0.404	TC12	0.727
17	ZI13	0.309	KI11	0.403	BC11	0.727
18	BL11	0.309	TA14	0.400	TA13	0.684
19	TA14	0.304	TC12	0.388	ZI14	0.622
20	BL14	0.301	BL14	0.378	TA14	0.615
21	TA9	0.289	ZI11	0.370	BL13	0.615
22	PV12	0.283	BC9	0.366	PV9	0.593
23	TC14	0.283	TA13	0.352	KI12	0.535
24	KI13	0.265	ZI12	0.339	TC11	0.533
25	TC12	0.259	BL13	0.329	ZI11	0.523
26	PV11	0.252	ZI13	0.329	ZI9	0.500
27	KI10	0.245	TC14	0.312	TC13	0.500
28	KI14	0.223	PV12	0.298	TC10	0.500
29	BL13	0.219	TC13	0.295	NI11	0.481
30	TA10	0.218	PV9	0.294	ZI10	0.469
31	TC13	0.211	TC11	0.290	TC14	0.460
32	BC9	0.211	ZI9	0.281	NI10	0.457
33	KI9	0.200	KI13	0.270	KI11	0.437
34	TC11	0.196	KI10	0.268	ZI13	0.426
35	Z19	0.195	PV11	0.267	BL12	0.411
36	PV9	0.195	TC9	0.267	KI9	0.411
37	TA13	0.182	TC10	0.262	KI10	0.411
38	ZI10	0.174	NI11	0.252	ZI12	0.363
39	BL12		KI9	0.235	PV12	
		0.158				0.371
40	PV10	0.153	KI14	0.228	PV10	0.371
41	TC10	0.139	ZI10	0.223	PV11	0.343
42	NI11	0.133	NI10	0.211	NI9	0.333
43	NI10	0.123	BL12	0.200	KI13	0.295
44	NI12	0.104	PV10	0.189	NI12	0.280
45	TC9	0.074	NI9	0.175	TC9	0.263
46	NI9	0.058	NI12	0.133	KI14	0.253
47	NI13	0.045	NI13	0.070	NI13	0.139
48	NI14	0.036	NI14	0.052	NI14	0.092

Source: Own calculations

Four cases are efficient according to the BCC model: Trnava in 2009 (TA9), Banska Bystrica - in 2014 (BC14) and Presov two years running in 2013 and 2014 (PV13 and PV14). To gain input efficiency it is necessary for inefficient regions to shift horizontally to the BCC production frontier. For example, Bratislava region in 2009 (BL9) should move to the line between points of Trnava region in 2009 (TA9) and of Banska Bystrica region in 2014 (BC14). The first inefficient region after the efficient regions is Banska Bystrica region in 2013 with a 34.9% input reduction. Once again. Nitra region was the last one in 2014 (NI14) with a slightly smaller input reduction 94.8% in comparison with the CCR model required reduction. As we have already mentioned, the FDH model is even more free. In our case seven regions are FDH efficient. In the year 2009, there was one efficient region: Trnava (TA9). Two regions were efficient in 2010: Banska Bystrica (BC10) and Trnava (TA10). In 2013 and 2014 other two regions were FDH efficient: Banska Bystrica (BC13, BC14) and Presov (PV13, PV14). The first inefficient region was Trnava in 2011, which has a reduction of only 5.9%. The last region in 2014 was Nitra with a reduction of 90.8%. The most frequent peer was Banska Bystrica region in 2013 (17 times); in 2014 (21 times) and in 2010 (4 times). Its proportion is 87.5%. If we take the criterion of 50% or more in the necessary reduction of beds as useless surplus of beds. then among the worst regions are: Nitra region (in all the analysed years from 2009 to 2014 it had double the required beds), Kosice - during five years, Presov and Zilina - during three years. In figure 2 are depicted The CCR and BCC production frontiers with Slovak regions in the period of 2009-2014 for adult day surgery are depicted in Figure 2.

In Table 3, there is an overview of corresponding DEA results analyses.

Now, from the viewpoint of the CCR model, Zilina region was efficient in 2009 (ZI9) (see also Figure 2). The second region is Zilina region in 2010 (ZI10) with a small 1.7 % reduction. Thus, Zilina region was almost efficient in 2010. The last region was Presov in 2011 (PV11) with a large necessary input reduction of 80.8 %.

On the basis of the BCC model, there are six efficient cases: Trnava and Zilina regions in 2009 (TA9 and ZI9), Zilina region in 2010 (ZI10), Banska Bystrica in 2012 (BC12) and Bratislava region in 2012 and in 2014 (BL12 and BL14). The first inefficient region was Kosice region in 2013 with a 3.5% input reduction. The last one is again Presov region in 2011 (PV11), with its slightly smaller input reduction (80.1%), was the last one.




Fig. 2: Production set of number of day surgery adult patients vs the number of day surgery adult beds with corresponding CCR and BCC DEA frontiers

Source: Own construction

Fifteen cases are efficient according to FDH model. In the year 2009, there were three efficient regions: Nitra (NI9), Trnava (TA9) and Zilina (ZI9). Two regions were efficient in 2010: Trnava (TA10) and Zilina (ZI10). In 2011, only Bratislava region (BL11) was FDH efficient. Four regions were efficient in 2012: Banska Bystrica (BC12), Bratislava (BL12), Trnava (TA12) and Zilina (ZI12). Three regions were efficient in the year 2013: Banska Bystrica (BC13), Kosice (KI13) and Trnava (TA13). In 2014, two regions, namely Bratislava (BL14) and Trencin (TC14), were efficient. The first inefficient region was Banska Bystrica in 2011 with an input reduction of only 0.8%. The last region was Presov in 2011 (PV11) with a reduction of 75.1%. The most frequent peers are Banska Bystrica in 2012 (11 times) and in 2013 (4 times), Zilina in 2009 (7 times) and Trnava in 2012 (4 times). Their proportion is 54.2%. Surplus beds are not so frequent as in the case of paediatric day surgery. More than 50% reduction is needed only in two regions: Presov (2009 and 2011) and Trencin (2009, 2010 and 2011).

Now we can make some summary evaluations. is An overview of the FDH efficiency results of Slovak regions in the period of six analysed years is shown in Table 4.

Banska Bystrica was relatively the best region in the field of paediatric day surgery. It was efficient three times: in 2010, 2013 and 2014. Two regions were efficient during two years: Presov (in 2013 and 2014) and Trnava (in 2009 and in 2010). All the other regions were not efficient at all during six years interval.

A different situation is in the field of adult day surgery where Trnava, which was FDH efficient during four years (2009, 2010, 2012 and 2013), is relatively the best region. The second place is held by two regions: Bratislava (2011, 2012 and 2014) and Zilina (2009, 2010 and 2012) with three efficient years. Banska Bystrica region follows with two efficient periods (2012 and 2013). Three regions were efficient once: Kosice (2013), Nitra (2009) and Trencin (2014). Presov region was not efficient during 2009-2014.

Tab. 3: Results of input oriented CCR, BCC a FDH DEA efficiency analyses of adult day surgery in Slovak regions in the period of 2009-2014

Rank	Regions	CCR eff.	Regions	BCC eff.	Regions	FDH eff.	
1	Z19	1	TA9	1	TC14	1	
2	ZI10	0.983	ZI9	1 BL14		1	
3	TA12	0.909	ZI10	1 TA13		1	
4	TA9	0.887	BC12	12 1 KI13		1	
5	BC12	0.881	BL12	1	NI9	1	
6	TA10	0.878	BL14	1	BC13	1	
7	NI9	0.788	KI13	0.965	TA9	1	
8	NI13	0.762	TA10	0.944	ZI12	1	
9	BL12	0.761	ZI12	0.933	ZI9	1	
10	TA11	0.757	TA12	0.920	TA12	1	
11	TA13	0.739	KI14	0.916	BL12	1	
12	ZI12	0.735	BL11	0.907	BC12	1	
13	BL11	0.728	ZI14	0.866	BL11	1	
14	BC11	0.683	NI13	0.854	ZI10	1	
15	ZI14	0.671	TC14	0.853	TA10	1	
16	BC13	0.670	NI9	0.826	BC11	0.992	
17	ZI13	0.636	BC13	0.792	KI14	0.981	
18	PV10	0.634	KI12	0.781	ZI13	0.956	
19	BC10	0.601	BL13	0.769	KI12	0.949	
20	TC14	0.599	TA13	0.767	ZI14	0.942	
21	KI13	0.588	TA11	0.760 ZI11		0.921	
22	NI12	0.587	ZI13	0.749	NI13	0.917	
23	BL13	0.587	BC11	0.727	NI14	0.845	
24	ZIII	0.575	BC14	0.683	PV10	0.835	
25	PV12	0.572	KI11	0.671	BL9	0.807	
26	KI14	0.571	ZI11	0.660	TA11	0.779	
27	NI10	0.568	NI12	0.648	BL13	0.777	
28	KI12	0.563	PV12	0.645	NI12	0.752	
29	BC9	0.562	PV10	0.635	BC14	0.751	
30	BL9	0.542	NI14	0.616	NI10	0.746	
31	BC14	0.541	BC10	0.604	KI11	0.740	
32	NI14	0.534	NI10	0.591	PV13	0.738	
33	KI11	0.531	PV14	0.588	BC10	0.724	
34	KI9	0.508	BL9	0.573	BL10	0.699	
35	BL10	0.485	PV13	0.570	KI9	0.691	
36	PV13	0.477	BC9	0.566	PV14	0.671	
37	PV13	0.477	KI9	0.551	PV12	0.665	
38	BL14	0.473	BL10	0.518	TC12	0.623	
39	NI11	0.444	TC13	0.494	BC9	0.596	
40	TC13	0.444	KI10	0.494	TA14	0.571	
41	TC12	0.443	TA14	0.456	TC13	0.550	
42	KI10 TC11	0.432	TC12 NI11	0.453	NI11 KI10	0.542	
44	TA14	0.420	TC11	0.429	TC11	0.460	
45	TC10	0.386		TC10 0.388 PV9		0.438	
46	TC9	0.334	TC9	0.368	TC9	0.427	
47	PV9	0.326	PV9	0.340	TC10	0.402	
48	PV11	0.192	PV11	0.199	PV11	0.249	

Source: Own calculations

Tab. 4: FDH efficient cases of day surgery in Slovak regions in the period of 2009-2014

Group	Paediatric day surgery						
Region/Year	2009	2010	2011	2012	2013	2014	
Banska Bystrica	1000	+	1000		+	+	
Bratislava							
Kosice							
Nitra							
Presov					_+	+	
Trnava	+	+					
Trencin	-						
Zilina							
Group	Adult day surgery						
Region/Year	2009	2010	2011	2012	2013	2014	
Banska Bystrica				+	+		
Bratislava			+	+		+	
Kosice			-	-	+		
Nitra	+			-			
Presov							
Trnava	+-	+		+	4		
Trencin						+	
Zilina	+	+		+			

Source: Own calculations

If we put together the results from both groups of day surgery, then Trnava, which was efficient six times during 12 evaluated cases (regions and years), is relatively the best region. The second place is held by Banska Bystrica which was efficient for five times. Two regions were efficient three times: Bratislava and Zilina.

We have found out the differences in the efficiency of units that were performing day surgery separately for children and adult patients in the individual regions of Slovakia by means of the analysis. It is very complex to form particular reasons of given differences. The deliberate use of own reporting methods in the given health facilities represents one of those reasons. It is caused by the present price strategies of health insurance companies. They influence day surgery availability by concluding the contracts with healthcare providers and setting financial limits. This leads to a deliberate transformation of procedures from day surgery group procedures to a separately reimbursed group of procedures by many healthcare providers (the patient is discharged within 72 hours - a 3-day procedure). This becomes financially more advantageous. If a healthcare provider performs the procedure that is already recorded as a one-day procedure without stating adequate reasons by means of a traditional hospitalization form, he receives a payment in the amount of a day surgery procedure from his/her health insurance company. It is related to deeper system issues that are connected with pricing policies of health insurance companies, short-term period of contractual relationships between healthcare providers and health insurance companies, as well as with other issues (legal, procedural, etc.).

5. Conclusions

The article deals with the issue of day surgery efficiency in Slovakia. It is part of health system and is not very highlighted. Thus, the development of day surgery in Slovakia has been stagnating for a long period of time. This may be proved by a minimum rate of day surgery procedures within the total amount of procedures. We have chosen a methodology that corresponds to the structure of available data, as well as to the given research target to evaluate the level of day surgery efficiency with regard to the individual regions of Slovakia.

We have conducted input oriented CCR, BCC a FDH DEA efficiency analyses of day surgery in Slovak regions (NUTS III level) in the period of 2009-2014 on basis of data offered to us by the National Health Information Center. From the available indicators, we have used the number of beds as the input. The number of day surgery patients was the output.

In case of paediatric day surgery, the most frequent peer was Banska Bystrica region in 2013 (17 times), in 2014 (21 times) and in 2010 (4 times) according to the FDH model. Its proportion is 87.5%. If we take the criterion of 50% or more in the necessary reduction of beds as useless surplus of beds, then among the worst regions are: Nitra region (in all the analysed years from 2009 to 2014 it had double the required beds), Kosice - during five years and Presov and Zilina - during three years.

In the field of adult day surgery, the most frequent peers are: Banska Bystrica in 2012 (11 times) and in 2013 (4 times), Zilina in 2009 (7 times) and Trnava in 2012 (4 times). Their proportion is 54.2%. Surplus beds are not as frequent as in the case of paediatric day surgery. More than a 50% reduction was needed only in two regions: Presov (2009 and 2011) and Trencin (2009, 2010 and 2011).

Significant differences were found on the basis of efficiency analysis of facilities that provide day surgery for child and adult patients in individual regions. It is an important result for those who will form health and social policies in Slovakia and for those who will form a strategic framework as well as for health legislation experts. The submitted analysis contains valuable information related to the specialisation of day surgery clinics. Day surgery will provide many options to increase health system efficiency, which is connected with bed reduction, cutting the hospitalisation time, elimination of waiting list periods for individual procedures, as well as making efficient a complex planning system. Its procedural and system consistency, together with the allocation of the present healthcare providers' network, will support the efficiency processes in Slovakia, as well as the contentment on the side of patients, healthcare providers, health insurance companies, the Ministry of Health of the Slovak Republic and other stakeholders.

References

- 1. Gavurova, B., & Hyranek, E. (2013). Determinants of Day Health Care Development in Slovakia. Ekonomicky casopis (Economic Annals), 61(2), 134-154
- (in Slovak). Retrieved from https://www.sav.sk/journals/uploads/0621142302%2013%20Gavurova-Hyranek-RS.pdf
 2. Gavurova, B., Stefko, R., & Bacik, R. (2014). The Analysis of Social Services' Structure in a Specific Region and its Significance for Health and Social Policy in Slovakia. *Polish Journal of Management Studies*, 10(2), 43-53. Retrieved from http://www.pjms.zim.pcz.pl/files/the-analysis-of-social-services--structure-in-a-
- 3. Jarrett, P., & Staniszewski, A. (2006). The Development of Ambulatory Surgery and Future Challenges. In: Lemos, P. Jarrett, P. and Philip, B. (eds): Day Surgery Development and Practice. London, UK: International Association for Ambulatory Surgery (IAAS), 21-34.

 4. Kroneman, M. W., et al. (2001). International variations in availability and diffusion of alternatives to in-patient care in Europe: the case of day surgery.

- 4. Krolenian, M. W., et al. (2001). International variations in availability administratives to in-patient care in Europe. the case of day surgery. 4. Krolenian, M. W., et al. (2001). 147-154. doi: http://dx.doi.org/10.1016/s0966-6532(01)00120-2
 5. Soltes, M., Pazinka, P., & Radonak, J. (2011a). Thermal injuries in laparoscopic surgery (in Slovak). Endoskopie (Endoscopy), 20(1), 14-16.
 6. Soltes, V., & Gavurova, B. (2014). Analysis of Selected Demographic Aspects of Day Surgery in Slovak Health Policy. Journal of Applied Economic Sciences, 29(3), 476-486. Retrieved from http://cesmaa.eu/journals/jaes/files/JAES_2014-Fall.pdf
 7. Soltes, M., Pazinka, P., & Radonak, J. (2010). Laparoscopic hernioplasty TAPP in treatment of groin hernia (in Slovak). Rozhledy v chirurgii (Complications in Surgery), 89(6), 384-389. Retrieved from http://www.prolekare.cz/rozhledy-v-chirurgii-clanek/laparoskopicka-hernioplastika-tapp-v-liecbe-slabinovej-prietrze
- 8. Soltes, M., & Radonak, J. (2014). A risk score to predict difficulty of elective laparoscopic cholecystectomy. Videosurgery and Other Miniinvasive Techniques, 9(46), 608-612. doi: http://dx.doi.org/10.5114/wiitm.2014.47642
- 9. Popa, D., Soltes, M., Uranues, S., & Fingerhut, A. (2015). Are there specific indications for laparoscopic appendectomy? A review and critical appraisal of the
- literature. Journal of Laparoendoscopic & Advanced Surgical Techniques, 25(11), 897-902. doi: http://dx.doi.org/10.1089/lap.2014.0624

 10. Soltes, M. (2011b). Economic aspects of laparoscopic surgery (in Slovak). Miniinvazivna chirurgia a endoskopia (Miniinvasive Surgery and Endoscopy), 15(3), 7-10.

 11. Szabo, S., & Sidor, J. (2014). The Performance Measurement System Potentials and Barriers for its Implementation in Healthcare Facilities, Journal of
- 11. Szabo, S., & Sidor, J. (2014). The Performance Measurement System Potentials and Barriers for its Implementation in Healthcare Facilities, *Journal of Applied Economic Sciences*, 30(4), 728-735. Retrieved from http://cesmaa.eu/journals/jaes/files/JAES_2014_Winter_short%20version.pdf
 12. Szczygiel, N., Rutkowska-Podolska, M., & Michalski, G. (2014). Information and Communication Technologies in Healthcare: Still Innovative and Entrepreneurial Value creating Approach in Healthcare Management. 5th Central European Conference in Regional Science CERS, 2014, 1020-1029.
 13. Soltes, M., & Gavurova, B. (2015). Quantification and Comparison of Avoidable Mortality Causal Relations and Modification of Concepts. Technological and Economic Development of Economy, 21(6), 917-938. doi: http://dx.doi.org/10.3846/20294913.2015.1106421
 14. Bem, A., Predkiewicz, K., Predkiewicz, P., & Ucieklak-Jez, P. (2014). Determinants of Hospital's Financial Liquidity. Procedia Economics and Finance, 12, 27-36. doi: http://dx.doi.org/10.1016/S2212-5671(14)00317-7
- 15. Bem, A., & Michalski, G. (2015). Hospital profitability vs. selected healthcare system indicators, CEFE 2015 Central European Conference in Finance and
- Economics, Technical University of Kosice, Kosice, 52-61.

 16. Cooper, W. W., Seiford, L. M., & Tone, K. (2007). Data Envelopment Analysis. A comprehensive text with models, applications, references and DEA-solver software. Springer. doi: http://dx.doi.org/10.1016/s0377-2217(02)00304-1
- 17. Charnes, A., Cooper, W. W., & Rhodes, E. (1978). Measuring the efficiency of decision-making units. *European Journal of Operational Research*, 2(6), 429-444. doi: http://dx.doi.org/10.1016/0377-2217(78)90138-8

 18. Banker R. D., Charnes, A., & Cooper, W. W. (1984). Some models for estimating technical and scale inefficiencies in Data Envelopment Analysis. *Management*
- Science, 30(9), 1078-1092. doi: http://dx.doi.org/10.1287/mnsc.30.9.1078

 19. Kocisova, K. (2012). Application of DEA Models at the Analysis of Bank Branches Technical Efficiency. Ekonomicky casopis (Economic Annals), 60(2), 169-186. Retrieved from https://www.sav.sk/journals/uploads/0622125302%2012%20Ko%C4%8Di%C5%A1ov%C3%A1%20RS.pdf
- 20. Kocisova, K. (2015). Application of the DEA on the measurement of efficiency in the EU countries. *Agricultural Economics*, 61(2), 51-62. doi: http://dx.doi.org/10.17221/107/2014-AGRICECON

 21. Deprins, D., Simar, L., & Tulkens, H. (2006). Measuring Labor Efficiency in Post Offices. In: Chander, P. Dreze, J. Lovell, C. K. Mintz, J. (Eds). *Public Goods, Environmental Externalities and Fiscal Competition*, pp. 285-309. doi: http://dx.doi.org/10.1007/978-0-387-25534-7_16

Received 23.03.2016