

KARPOV P. A.[✉], DEMCHUK O. M., OZHEREDOV S. P., SPIVAK S. I., YEMETS A. I., BLUME Ya. B.

*Institute of Food Biotechnology and Genomics of Natl. Acad. Sci. of Ukraine,
Ukraine, 04123, Kyiv, Osypovs'koho str., 2A*

[✉] *karpov@nas.gov.ua, (050) 544-69-03*

CONSERVATION OF DINITROANILINE/PHOSPHOROTHIOAMIDATE SITE OF α -TUBULIN IN *PLASMODIUM* SPECIES DISTRIBUTED IN INDIA

Aim. To evaluate polymorphism of dinitroaniline/phosphorothioamidate binding site in α -tubulin molecules from *Plasmodium* species (*P. falciparum*, *P. vivax*, *P. ovale*, and *P. malariae*) and strains discovered in India. **Methods.** Literature and database search. Bioinformatical comparison of protein sequences and structures. Multiple sequence alignment, phylogenetic profiling, protein structure modeling, etc. **Results.** 14 complete sequences of α -tubulin from four *Plasmodium* species were selected from UniProtKB. Despite certain differences in complete sequences of α -tubulin molecules from different *Plasmodium* species and strains, sites of their interaction with dinitroaniline and phosphorothioamidate compounds were shown to be identical.

Conclusions. Complete identity of dinitroaniline/phosphorothioamidate binding sites in all studied isotypes of α -tubulin molecules from *P. falciparum*, *P. vivax*, *P. ovale*, and *P. malariae* was confirmed. This suggests identity of the ligand-protein interaction mechanism and similar destabilizing effect of dinitroaniline and phosphorothioamidate compounds on microtubules of all *Plasmodium* species officially registered in India.

Keywords: malaria, *Plasmodium*, α -tubulin, dinitroanilines, phosphorothioamidates, binding site.

Plasmodium parasites are the main cause of global malaria infections with 300–500 million clinical cases and 1–3 million deaths per year [1]. It is known that malaria is caused at least by a few species of *Plasmodium* protozoan parasites: *P. falciparum*, *P. vivax*, *P. ovale*, and *P. malariae* [2]. The National Vector Borne Disease Control Program of India reported ~1.6 million cases and ~1100 malaria deaths in 2009 [3]. Historically, malaria in India was predominantly caused by *P. vivax*, accounting for 53% of the estimated cases. After the spread of drug-resistant *P. falciparum* in the 1990s, the prevalence of the two species re-

mained equivalent. The burden of malaria in India is complex because of the highly variable malaria eco-epidemiological profiles, transmission factors, and the presence of multiple *Plasmodium* species and *Anopheles* vectors [4]. Now two major human malaria species in India are *P. falciparum* and *P. vivax*. *P. malariae* has been reported in the eastern India state of Orissa [5], while *P. ovale* appears to be extremely rare if not absent [3]. *P. falciparum* is most severe strain of the malaria species correlated with almost every malarial death (CDC - www.cdc.gov/malaria/about/disease.html). *P. falciparum* is strongly associated with severe disease syndrome known as cerebral malaria, which is associated with high mortality [2].

The control of malaria infection is hampered by many factors, including emerging drug resistance. It is a fact that many of existing now malaria therapeutics are increasingly ineffective and it is an urgent need in development of principally new therapeutic strategies and agents. During last ten years there has been a new growth of interest in tubulin as an important antiprotozoan target [6, 7]. Several classes of microtubule (MT) inhibitors have demonstrate potent activity against malarial parasites in *in vivo*: vinblastine [8–10], dolastatin 10 [11], auristatins [12] and taxoids [13, 14]. Most of these agents have been demonstrated to disrupt or stabilize normal microtubular structures. Unfortunately, most all these compounds show toxicity to mammalian cells [15, 16] due to the interspecies conservation of tubulin [16]. *P. falciparum* and human α -tubulins share ~83% identity and β -tubulins ~87%. However it was found that human antimalarial drugs (e.g. sulfadiazine, sulfadoxine, pyrimethamine, cycloguanyl) were lethal for the model plant *Arabidopsis thaliana* at similar concentrations to market herbicides (glufosinate and glyphosate) [17]. Although MT inhibitors from anti-cancer programs have proved useful in probing MT function in parasites, such non-selective agents

[©] **KARPOV P. A., DEMCHUK O. M., OZHEREDOV S. P., SPIVAK S. I., YEMETS A. I., BLUME Ya. B.**

have no prospects as antimalarial drugs [15, 16]. Later it was found that dinitroanilines (oryzalin, trifluralin, chloralim, pendimethalin) and phosphorothioamides (aminprophos-methyl) destabilize MT and have overlapping sites in plant tubulin. It was proved that dinitroanilines and phosphorothioamides bind plant α -tubulin with high affinity, inhibit tubulin assembly and disrupt MTs [18, 19]. Affected cells do not form a cell plate and may contain restitution and polyploidy nuclei [16].

It is very important that dinitroaniline/phosphorothioamide site is absent in human tubulins [16, 20]. However, this site was found in some *Protozoa* [16]. This explains why some dinitroaniline derivatives with selectivity for parasite MTs are more than 800-fold effective on the parasite than on various human cell lines [21]. As a result of this relationship, studies have demonstrated that dinitroanilines and phosphorothioamides in most cases active against plants are also active against *P. falciparum* and thus could act as antimalarial drug leads [17]. α -Tubulin appears to be an outstanding target for such parasitic diseases as leishmaniasis and trypanosomiasis, as the respective parasites also display altered or absent checkpoints [22, 23]. Phosphorothioamides have similar effects on plants as the dinitroanilines and bind to the same molecular site. As phosphorothioamides have a more than 100-fold higher solubility in aqueous solutions than dinitroanilines, these compounds are more promising candidates for modification than dinitroanilines, where biological studies indicated that maintaining high sufficient drug concentrations is difficult [24]. Therefore dinitroanilines and phosphorothioamides can be considered as a priority group for the search of antimalarial agents.

The purpose of current study was to compare dinitroaniline/phosphorothioamide binding site in α -tubulin molecules from *Plasmodium* species (*P. falciparum*, *P. vivax*, *P. ovale*, and *P. malariae*) and strains, native for Indian region, and draw conclusions regarding the level of its polymorphism.

Materials and methods

Complete sequences of 14 α -tubulins from *P. falciparum* (PLAFA|PLAFK|PLAF7: A0A1D3KY74_PLAFA, A0A2Y9URV4_PLAFA, Q25790_PLAFA, TBA_PLAFK, Q6ZLZ9_PLAF7, Q8IFP3_PLAF7), *P. vivax* (A0A1G4GUY4_PLAVI, A0A1G4H9I2_PLAVI), *P. ovale* (A0A1C3KQ56_9APIC, A0A1A8VN48_9APIC, A0A1A8VPR2_9APIC, A0A1A8YLP3_9APIC), and *P. malariae* (A0A1A8VRI3_PLAMA, A0A1A8VSS4_PLA

MA) were selected from UniProtKB [25].

The information on amino acid residues of dinitroaniline/phosphorothioamide binding site in α -tubulin molecule and residues of known mutations upset ligand binding were based on the analysis of previously published results [20, 26-30].

Structural model of α -tubulin molecule from *P. falciparum* (TBA_PLAFK, UniProtKB: P14642) was built using protein structure homology-modelling server Swiss-Model [31] based on template RCSB Protein Data Bank (www.rcsb.org) [32] structures: 5UBQ (A) – α -tubulin from cilia of *Tetrahymena thermophila* (Cryo-EM structure) [33] and 2.5 E X-RAY structure 5KX5 (A) of α -tubulin from *Ovis aries* [34]. All visualizations and analysis of PDB-structures and constructed model of *Plasmodium* α -tubulin were performed using PyMOL v.1.5.0.5 software (www.pymol.org).

Multiple alignments of the complete amino acid sequences of α -tubulin molecules from different *Plasmodium* species and strains were performed with the ClustalX (V.2.0.10) (www.clustal.org) software using BLOSSUM weight matrices [35]. In the case of the comparison of small polypeptide fragments of the sites of interaction with dinitroaniline and phosphorothioamide compounds, we applied direct comparison of the sequences in ClustalX with restriction of gaps [35]. All clustering was based on similarity of complete amino acid sequences and their fragments. Visualizations and analysis of dendograms were performed with the MEGA7 software package (www.megasoftware.net/) [36].

Results and discussions

Initial analysis of information presented in the UniProtKB database revealed 17 α -tubulin sequences of four *Plasmodium* species distributed in India (*P. falciparum*, *P. vivax*, *P. ovale*, and *P. malariae*). At the same time, only 14 of them are presented by complete sequences covering binding region of dinitroaniline and phosphorothioamide compounds. From above mentioned sequences six of them belong to *P. falciparum* (isolates PLAFA, PLAFK/K1 and PLAF7/3D7), 2 – *P. vivax* (PLAVI), 4 – *P. ovale* (9APIC), and 2 – *P. malariae* (PLAMA). The subsequent alignment and reconstruction of the phylogenetic tree revealed variations in the complete α -tubulin sequences from different *Plasmodium* species, isolates and strains (Fig. 1). Thus, it was a probability of certain differences of dinitroaniline/phosphorothioamide binding sites in α -tubulin molecules from different species and ecotypes.




Fig. 1. Phylogenetic clustering of complete sequences of α -tubulin molecules from *Plasmodium* species (*P. falciparum*, *P. vivax*, *P. ovale*, and *P. malariae*) and strains officially registered in India.

To reveal similarity of dinitroaniline/phosphorothioamidate binding site in α -tubulin molecules from different *Plasmodium* species, literature analysis was carried out and information on the amino acid composition and structure of this site among *Protozoa* (*P. falciparum* and *Toxoplasma gondii*) and higher plant *Eleusine indica* was summarized [20, 26-30]. Selecting functionally important amino acids, we used such criteria as direct participation in the binding of ligands, involvement

in the structure of site, as well as amino acids, mutations on which cause resistance to dinitroaniline and phosphorothioamidate compounds (Table). Additionally, the amino acids described in the literature as related to the dinitroaniline / phosphorothioamidate binding site, were verified based on their structural position using 3D-model of α -tubulin from *P. falciparum* (TBA_PLAFK, UniProtKB: P14642) as control (Fig. 2).

Table. Amino acid residues of dinitroaniline/phosphorothioamidate binding site in α -tubulin and residues of known mutations upset ligand binding

Species	Deposition in UniProtKB	Amino acids of binding site	Source of information
<i>Plasmodium falciparum</i>	TBA_PLAFK (P14642)	Arg2, Gln133, Arg243, Asn249, Val250, Asp251, Val252, Thr253, Glu254	[27, 28]
<i>Toxoplasma gondii</i>	TBA_TOXGO (P10873)	Arg2, His28, Gln133, Thr239, Ala240, Arg243	[30]
		Arg2, Glu3, Val4, Trp21, Phe24, His28, Asp47, Arg64, Cys65, Thr239, Arg243, Phe244	[20]
		Val4, Ser6, Phe24, His28, Leu136, Ile235, Thr239, Arg243 (mutated in resistant parasites)	[29]
<i>Eleusine indica</i>	TBA1_ELEIN (O22347)	Arg2, Gln133, Arg243, Asn249, Val250, Asp251, Val252, Asn253, Glu254	[26, 27, 28]

Note. Italics mark amino acid residues, mutation of which causes resistance to dinitroaniline and phosphorothioamidate compounds.

Comparing known data, the results of structural modeling and alignment of 14 α -tubulin sequences from *P. falciparum*, *P. vivax*, *P. ovale* and *P. malariae*, the residues of dinitroaniline/phosphorothioamidate binding site were identified. Based on this information, the associated fragments were extracted from complete sequences. In addition to the above-mentioned amino acid residues, amino acids adjacent to them along the N- and C-directions of the polypeptide chain were also selected. The alignment of selected fragments reveals their complete identity, what was also confirmed by results of clustering (Fig. 3).

Thus, despite differences in complete amino acid sequences of α -tubulin from *Plasmodium* species (*P. falciparum*, *P. vivax*, *P. ovale*, and *P. malariae*) and strains, distributed in India, their sites of interaction with dinitroanilines and phosphorothioamidates are completely identical. Considering inherent tubulins strong similarity of spatial structures [27, 37-39], it can be said with confidence that dinitroaniline and phosphorothioamidate compounds will have the similar effect on the MTs of all studied species and strains.




Fig. 2. Structural topology (a) of dinitroaniline and phosphorothioamidate binding site and its amino acids (b) in *P. falciparum* α -tubulin: Arg2, Glu3, Val4, Ser6, Trp21, Phe24, His28, Asp47, Arg64, Cys65, Gln133, Leu136, Ile235, Thr239, Ala240, Arg243, Phe244, Asn249, Val250, Asp251, Val252, Thr253 and Glu254. Red color (dark grey in monochrome representation) sticks mark amino acid residues, mutations of which cause resistance to dinitroanilines and phosphorothioamidates: Leu136, Ile235, Thr239 and Arg243.

A0A1D3KY74_PLAFA_GN=PocGH01_05	MREVISI-CWELFC-EHG-DDA-PRCV-LQGFLM-VISSLTASLRFD-LNVDVTEF
A0A2Y9URV4_PLAFA_GN=PocGH01_07	MREVISI-CWELFC-EHG-DDA-PRCV-LQGFLM-VISSLTASLRFD-LNVDVTEF
Q25790_PLAFA_GN=AAA29498.1	MREVISI-CWELFC-EHG-DDA-PRCV-LQGFLM-VISSLTASLRFD-LNVDVTEF
TBA_PLAFK_P14642_isolate_K1_Th	MREVISI-CWELFC-EHG-DDA-PRCV-LQGFLM-VISSLTASLRFD-LNVDVTEF
Q6ZLZ9_PLAF7_isolate_3D7_GN=PF	MREVISI-CWELFC-EHG-DDA-PRCV-LQGFLM-VISSLTASLRFD-LNVDVTEF
Q8IFP3_PLAF7_isolate_3D7_GN=PF	MREVISI-CWELFC-EHG-DDA-PRCV-LQGFLM-VISSLTASLRFD-LNVDVTEF
AOA1G4GUY4_PLAVI_PVC01_0700069	MREVISI-CWELFC-EHG-DDA-PRCV-LQGFLM-VISSLTASLRFD-LNVDVTEF
AOA1G4H9I2_PLAVI_GN=PVC01_0500	MREVISI-CWELFC-EHG-DDA-PRCV-LQGFLM-VISSLTASLRFD-LNVDVTEF
AOA1A8VRI3_PLAMA_GN=PmUG01_050	MREVISI-CWELFC-EHG-DDA-PRCV-LQGFLM-VISSLTASLRFD-LNVDVTEF
AOA1A8VSS4_PLAMA_Tubulin_GN=Pm	MREVISI-CWELFC-EHG-DDA-PRCV-LQGFLM-VISSLTASLRFD-LNVDVTEF
AOA1C3KQ56_9APIC_GN=PowCR01_07	MREVISI-CWELFC-EHG-DDA-PRCV-LQGFLM-VISSLTASLRFD-LNVDVTEF
AOA1A8VN48_9APIC_GN=POVCU1_011	MREVISI-CWELFC-EHG-DDA-PRCV-LQGFLM-VISSLTASLRFD-LNVDVTEF
AOA1A8YLP3_9APIC_GN=POVWA1_012	MREVISI-CWELFC-EHG-DDA-PRCV-LQGFLM-VISSLTASLRFD-LNVDVTEF
AOA1A8VPR2_9APIC_GN=POVCU1_009	MREVISI-CWELFC-EHG-DDA-PRCV-LQGFLM-VISSLTASLRFD-LNVDVTEF
..	*****

Fig. 3. Results of clustering (at the left) and alignment (on the right) of fragments, associated with dinitroaniline and phosphorothioamidate binding site in sequences of α -tubulins of studied strains of *P. falciparum*, *P. vivax*, *P. ovale*, and *P. malariae*, registered in India.

Conclusions

Complete identity of dinitroaniline/phosphorothioamidate binding sites in all studied isotypes of α -tubulin molecules from *P. falciparum*, *P. vivax*, *P. ovale*, and *P. malariae* was confirmed. Considering inherent tubulin molecules strong similarity of spatial structures, it can be concluded that individual dinitroaniline and phosphorothioamidate compounds will have the same effect

on MTs of all *Plasmodium* species officially registered in India. Such conservatism makes it possible to simplify subsequent protocol of virtual screening and perform it against only one molecular target.

This research was supported by grant of the Joint Ukraine-Indian Republic R&D Projects in 2019–2021 initiated by the Ministry of Education and Science of Ukraine and The Department of Science & Technology of Ministry of Science & Technology, India.

References

- Schantz-Dunn J., Nour N.M. Malaria and pregnancy: a global health perspective. *Rev. Obstet. Gynecol.* 2009. Vol. 2 (3). P. 186–192.
- Molina-Cruz A., DeJong R.J., Ortega C., Haile A., Abban E., Rodrigues J., Jaramillo-Gutierrez G., Barillas-Mury C. Some strains of *Plasmodium falciparum*, a human malaria parasite, evade the complement-like system of *Anopheles gambiae* mosquitoes. *Proc. Natl. Acad. Sci. USA.* 2012. Vol. 109 (28). E1957–1962.
- Das A., Anvikar A.R., Cator L.J., Dhiman R.C., Eapen A., Mishra N., Nagpal B.N., Nanda N., Raghavendra K., Read A.F., Sharma S.K., Singh O.P., Singh V., Sinnis P., Srivastava H.C., Sullivan S.A., Sutton P.L., Thomas M.B., Carlton J.M., Valecha N. Malaria in India: The center for the study of complex malaria in India. *Acta Tropica.* 2012. Vol. 121 (3). P. 267–273.
- Anvikar A.R., Shah N., Dhariwal A.C., Sonal G.S., Pradhan M.M., Ghosh S.K., Valecha N. Epidemiology of *Plasmodium vivax* malaria in India. *Am. J. Trop. Med. Hyg.* 2016. Vol. 95 (6S). P. 108–120.
- Sharma S.K., Tyagi P.K., Padhan K., Upadhyay A.K., Haque M.A., Nanda N., Joshi H., Biswas S., Adak T., Das B.S., Chauhan V.S., Chitnis C.E., Subbarao S.K. Epidemiology of malaria transmission in forest and plain ecotype villages in Sundargarh District, Orissa, India. *Trans. R. Soc. Trop. Med. Hyg.* 2006. Vol. 100 (10). P. 917–925.
- Britsun V.M., Yemets A.I., Lozinskii M.O., Blume Ya.B. 2,6-Dinitroanilines: synthesis, herbicidal and antiprotozoan properties. *Ukr. Bioorg. Acta.* 2009. Vol. 7 (1). P. 16–27.
- Ota S., Tomioka S., Sogawa H., Satou R., Fujimori M., Karpov P., Shulga S., Blume Ya., Kurita N. Binding properties between curcumin and malarial tubulin: molecular docking and ab initio fragment molecular orbital calculations. *Chem-Bio Informatics J.* 2018. Vol. 18. P. 44–57.
- Usanga E.A., O'Brien E., Luzzato L. Mitotic inhibitors arrest the growth of *Plasmodium falciparum*. *FEBS Lett.* 1986. Vol. 209. P. 23–27.
- Bell A., Wernli B., Franklin R.M. Effects of microtubule inhibitors on protein synthesis in *Plasmodium falciparum*. *Parasitol. Res.* 1993. Vol. 79. P. 146–152.
- Dieckmann-Schuppert A., Franklin R.M. Compounds binding to cytoskeletal proteins are active against *Plasmodium falciparum* in vitro. *Cell Biol. Int.* 1989. Vol. 13. P. 411–418.
- Nath J., Schneider I. Anti-malarial effects of the anti-tubulin herbicide trifluralin: studies in *Plasmodium falciparum*. *Clin. Res.* 1992. Vol. 40. #331A.
- Fennell B.J., Carolan S., Pettit G.R., Bell A. Effects of the antimitotic natural product dolastatin 10, and related peptides, on the human malarial parasite *Plasmodium falciparum*. *J. Antimicrob. Chemother.* 2003. Vol. 51. P. 833–841.
- Schrevel J., Sinou V., Grellier P., Frappier F., Guénard D., Potier P. Interactions between docetaxel (Taxotere) and *Plasmodium falciparum* – infected erythrocytes. *Proc. Natl. Acad. Sci. USA.* 1994. Vol. 91. P. 8472–8476.
- Pouvelle B., Farley P.J., Long C.A., Taraschi T.F. Taxol arrests the development of blood-stage *Plasmodium falciparum* in vitro and *Plasmodium chabaudi adami* in malaria-infected mice. *J. Clin. Invest.* 1994. Vol. 94. P. 413–417.
- Bell A. Microtubule inhibitors as potential antimalarial agents. *Parasitol. Today.* 1998. Vol. 14. P. 234–240.
- Fennell B.J., Naughton J.A., Dempsey E., Bell A. Cellular and molecular actions of dinitroaniline and phosphorothioamidate herbicides on *Plasmodium falciparum*: tubulin as a specific antimalarial target. *Mol. Biochem. Parasitol.* 2006. Vol. 145 (2). P. 226–238.
- Corral M.G., Leroux J., Stubbs K.A., Mylne J.S. Herbicidal properties of antimalarial drugs. *Sci. Repts.* 2017. Vol. 7. # 45871.
- Lyons-Abbott S., Sackett D.L., Wloga D., Gaertig J., Morgan R.E., Werbovetz K.A., Morissette N.S. 6-Tubulin mutations alter oryzalin affinity and microtubule assembly properties to confer dinitroaniline resistance. *Eukar. Cell.* 2010. Vol. 9 (12). P. 1825–1834.
- Yemets A.I., Blume Y.B. Antimitotic drugs for microprotoplast-mediated chromosome transfer in plant genomics, cell engineering and breeding. In: Blume Y.B., Baird W.V., Yemets A.I., Breviaro D. (eds) *The Plant Cytoskeleton: a Key Tool for Agro-Biotechnology*. Springer, Dordrecht, 2008. P. 419–434.
- Morissette N.S., Mitra A., Sept D., Sibley L.D. Dinitroanilines bind α -tubulin to disrupt microtubules. *Mol. Biol. Cell.* 2004. Vol. 15 (4). P. 1960–1968.
- Morgan R.E., Ahn S., Nzimiro S., Fotie J., Phelps M.A., Cotrill J., Yakovich A.J., Sackett D.L., Dalton J.T., Werbovetz K.A. Inhibitors of tubulin assembly identified through screening a compound library. *Chem. Biol. Drug Design.* 2008. Vol. 72 (6). P. 513–524.
- Robinson D.R., Sherwin T., Ploubidou A., Byard E.H., Gull K. Microtubule polarity and dynamics in the control of organelle positioning, segregation, and cytokinesis in the trypanosome cell cycle. *J. Cell Biol.* 1995. Vol. 128. P. 1163–1172.
- Werbovetz K.A. Tubulin as an antiprotozoal drug target. *Mini Rev. Med. Chem.* 2002. Vol. 2. P. 519–529.

24. Dhooghe E., Van L.K., Eeckhaut T., Leus L., Van H.J. Mitotic chromosome doubling of plant tissues *in vitro*. *Plant Cell Tiss. Org. Cult.* 2011. Vol. 104. P. 359–373.
25. UniProt Consortium T. UniProt: the universal protein knowledgebase. *Nucl. Acids Res.* 2018. Vol. 46 (5). P. 2699.
26. Nyporko A.Yu., Yemets A.I., Klimkina L.A., Blume Ya.B. Sensitivity of *Eleusine indica* callus to trifluralin and amiprophen-methyl in correlation with the binding of these compounds to tubulin. *Russ. J. Plant Physiol.* 2002. Vol. 49 (3). P. 413–418.
27. Nyporko A.Y., Yemets A.I., Brytsun V.N., Lozinsky M.O., Blume Y.B. Structural and biological characterization of the tubulin interaction with dinitroanilines. *Cytol. Genet.* 2009. Vol. 43. P. 267–282.
28. Chu Z., Chen J., Nyporko A., Han H., Yu Q., Powles S. Novel α -tubulin mutations conferring resistance to dinitroaniline herbicides in *Lolium rigidum*. *Front. Plant Sci.* 2018. Vol. 9. P. 97. doi.org/10.3389/fpls.2018.00097.
29. Ma C., Li C., Ganesan L., Oak J., Tsai S., Sept D., Morrisette N.S. Mutations in alpha-tubulin confer dinitroaniline resistance at a cost to microtubule function. *Mol. Biol. Cell.* 2007. Vol. 18 (12). P. 4711–4720.
30. Ma C., Tran J., Gu F., Ochoa R., Li C., Sept D., Werbovetz K., Morrisette N. Dinitroaniline activity in *Toxoplasma gondii* expressing wild-type or mutant alpha-tubulin. *Antimicrob. Agents Chemother.* 2010. Vol. 54 (4). P. 1453–1460.
31. Waterhouse A., Bertoni M., Bienert S., Studer G., Tauriello G., Gumienny R., Heer F.T., de Beer T.A.P., Rempfer C., Bordoli L., Lepore R., Schwede T. SWISS-MODEL: homology modelling of protein structures and complexes. *Nucl. Acids Res.* 2018. Vol. 46 (W1). W296–W303.
32. Berman H.M., Westbrook J., Feng Z., Gilliland G., Bhat T.N., Weissig H., Shindyalov I.N., Bourne P.E. The Protein Data Bank. *Nucl. Acids Res.* 2000. Vol. 28 (1). P. 235–242.
33. Ichikawa M., Liu D., Kastritis P.L., Basu K., Hsu T.C., Yang S., Bui K.H. Subnanometre-resolution structure of the doublet microtubule reveals new classes of microtubule-associated proteins. *Nat. Commun.* 2017. Vol. 8. P. 15035. doi: 10.1038/ncomms15035.
34. Leverett C.A., Sukuru S.C., Vetelino B.C., Musto S., Parris K., Pandit J., Loganzo F., Varghese A.H., Bai G., Liu B., Liu D., Hudson S., Doppalapudi V.R., Stock J., O'Donnell C.J., Subramanyam C. Design, synthesis, and cytotoxic evaluation of novel tubulysin analogues as ADC payloads. *ACS Med. Chem. Lett.* 2016. Vol. 7. P. 999–1004.
35. Larkin M.A., Blackshields G., Brown N.P., Chenna R., McGettigan P.A., McWilliam H., Valentin F., Wallace I.M., Wilm A., Lopez R., Thompson J.D., Gibson T.J., Higgins D.G. ClustalW and ClustalX version 2.0. *Bioinformatics*. 2007. Vol. 23 (21). P. 2947–2948.
36. Kumar S., Stecher G., Tamura K. MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol. Biol. Evol.* 2016. Vol. 33 (7). P. 1870–1874.
37. Nogales E., Wolf S.G., Downing K.H. Structure of the alpha beta tubulin dimer by electron crystallography. *Nature*. 1998. Vol. 391 (6663). P. 199–203.
38. Blume Ya.B., Nyporko A.Yu., Yemets A.I., Baird W.V. Structural modeling of the interaction of plant α -tubulin with dinitroaniline and phosphoroamide herbicides. *Cell Biol. Int.* 2003. Vol. 27 (3). P. 171–174.
39. Tuszyński J.A., Carpenter E.J., Huzil J.T., Malinski W., Luchko T., Luduena R.F. The evolution of the structure of tubulin and its potential consequences for the role and function of microtubules in cells and embryos. *Int. J. Dev. Biol.* 2006. Vol. 50 (2–3). P. 341–358.

КАРПОВ П. А., ДЕМЧУК О. М., ОЖЕРСДОВ С. П., СПІВАК С. І., ЄМЕЦЬ А. І., БЛЮМ Я. Б.

Державна установа «Інститут харчової біотехнології та геноміки НАН України»,
Україна, 04123, м. Київ, вул. Осиповського, 2а

КОНСЕРВАТИВНІСТЬ САЙТУ ЗВ'ЯЗУВАННЯ ДИНІТРОАНІЛІНІВ І ФОСФОРТОІОАМІДІВ МОЛЕКУЛАМИ α -ТУБУЛІНУ У РОЗПОВСЮДЖЕНИХ В ІНДІЇ ПРЕДСТАВНИКІВ РОДУ *PLASMODIUM*

Мета. З'ясувати ступінь поліморфізму сайту зв'язування динітроаніліну/фосфортоіоаміду на поверхні молекул α -тубуліну у різних штамів представників роду *Plasmodium* (*P. falciparum*, *P. vivax*, *P. ovale*, *P. malariae*), що зустрічаються на території Індії. **Методи.** Аналіз літератури та баз даних. Біоінформатичне порівняння білкових послідовностей і структур. Множинні вирівнювання послідовностей, філогенетичне профілювання, моделювання просторової структури білків тощо. **Результатами.** З бази даних UniProtKB було відібрано 14 повних послідовностей α -тубуліну чотирьох видів роду *Plasmodium*. Встановлено, що незважаючи на певні відмінності повних послідовностей молекул α -тубуліну у різних видів і штамів *Plasmodium*, сайти їх взаємодії з похідними динітроаніліну і фосфортоіоаміду повністю ідентичні. **Висновки.** Підтверджено повну ідентичність сайтів зв'язування динітроаніліну/фосфортоіоаміду у всіх дослідженіх ізотипів молекул α -тубуліну з *P. falciparum*, *P. vivax*, *P. ovale* та *P. malariae*. Це свідчить про ідентичність механізму ліганд-білкової взаємодії та схожий дестабілізуючий ефект похідних динітроаніліну і фосфортоіоаміду на мікротрубочки вищезазначених видів *Plasmodium*, що офіційно зареєстровані на території Індії.

Ключові слова: малярія, *Plasmodium*, α -tubulin, похідні динітроаніліну, похідні фосфортоіоаміду, сайт зв'язування.