- 2. In looking for EQ precursors (supposing be aperiodic) annual variation should be well studied and reduced.
- 3. Causes of annual variation may be a) the change of electrical conductivity in the Earth's in-

terior, forming induction vectors; b) seasonal variation of the external source parameters leading to deflection from T—C model. c) superposition of seasonally variable parts of the last terms in equation (3).

## References

Fujita S. Monitoring of Time Changes of Conductivity Anomaly Transfer Functions at Japanese Magnetic Observatory Network // Memoirs of the Kakioka magnetic observatory. — 1990. — 23, № 2. — P. 53—87.

Korepanov V. E., Tregubenko V. I. Features of construction and trends of modern instruments development for magnetotelluric and magnetovariational soundings // Geophys. J. — 2009. — 31, № 4. — P. 181—190 (in Russian).

Kuboki T. The Anomaly of Geomagnetic Variation in Japan (Part V) // Memoirs of the Kakioka magnetic observatory. — 1972. — 15, № 1. — P. 63—80 (in Japanese).

Moroz Yu. F., Moroz T. A. Magnetovariational studies

in the Baykal lake region // Fizika Zemly. — 2006. — N 11. — P. 93—98 (in Russian).

Moroz Yu. F., Smirnov S. E., Moroz T. A. The results of monitoring of geomagnetic field variations in Kamchatka // Fizika Zemly. — 2006. — № 3. — P. 49—56 (in Russian).

Shiraki M., Yanagihara K. Transfer Functions at Kakioka (Part I) // Memoirs of the Kakioka magnetic observatory — 1975. — 16, № 2. — P. 143—155 (in Japanese).

Shiraki M., Yanagihara K. Transfer function at Kakioka (Part II). Reevaluation of Their Secular Changes // Memoirs of the Kakioka magnetic observatory. — 1977. — 17, № 1. — P. 19—25 (in Japanese).

## **High Rayleigh Number Mantle Convectionon GPU**

© D. Sanchez<sup>1</sup>, D. Yuen<sup>1</sup>, G. Wright<sup>2</sup>, G. Barnett Jr.<sup>3</sup>, 2010

<sup>1</sup>Minnesota Supercomputing Institute, University of Minnesota, USA <sup>2</sup>Department of Mathematics, Boise State University, USA <sup>3</sup>Department of Applied Mathematics, University of Colorado, Boulder, USA

The performance, potential growth, availability, and affordability of GPUs has made them attractive to scientists for many years. Although historically cumbersome and difficult to use for scientific software, the introduction and refinement of high-level development tools, such as CUDA, have made GPU computing accessible. With the advent of architectures, such as NVIDIA's Fermi, which explicitly cater to scientists by enabling more memory, faster access to that memory, and better double-precision support than ever before, members of the computational world are finding GPU difficult to ignore.

Using finite-difference methods with second-order accuracy in space and third-order accuracy in time, we investigate 2D and 3D thermal convection at the infinite Prandtl number limit, at resolutions on the order of  $1000\times2000~2D$  and  $400\times400\times200~3D$  grid points. Our CUDA code makes extensive use of




Fig. 1. Performance scaling with grid size on Tesla C1060.



Fig. 2. 2D mantle convection at a Rayleigh number of 109.

highly-optimized CUBLAS routines, allowing us to unlock a significant fraction of GPU's performance. This performance has enabled us to study the behavior of high Rayleigh number simulations, on the order of 10<sup>9</sup>, in 2D and 10<sup>7</sup> in 3D over sufficient time scales to see evidence of flow-reversal (Fig. 1).

We compare our CUDA code's performance with Jacket-accelerated Matlab code and CPU-only Matlab code across the Tesla C1060, Tesla C2050, and GTX 480 (Fig. 2).

## Observation and analysis of ULF data associated with $M_b$ =4.5 Koyna-Warna (India) earthquake

© A. Sharma, A. Patil, 2010

Space and Earth Sciences, Department of Physics, Shivaji University Kolhapur, India sharma\_ashokkumar@yahoo.com

In recent years ULF (Ultra Low Frequency) emission is detected during the seismic activity and this emission is recognized as one of the most promising candidate for earthquake prediction. In this paper, we present measurements of magnetic field anomalies detected for moderate earthquake that occurred on 29 July, 2008 having magnitude (M<sub>b</sub>) 4.5. The ULF observation system uses three-component induction coil magnetometer and it has been installed at Shivaji University Kolhapur (16.40°N, 74.15°E), India.

Data of moderate earthquake have been analyzed using spectral density and polarization methods [Hattori et al., 2002a, b]. Long term data analysis shows that, two anomalous enhancements in

intensity of magnetic field were observed about two and one months before the earthquake. Short term data analysis shows that, maximum enhancement in intensity of magnetic field was prominent within  $\pm 1$  hours around the main shock time and it started to be observe four to five days before the earthquake. The enhancement in intensity of magnetic field is examined in terms of space magnetic pulsation and ULF emission associated with earthquake by using polarization parameter [Hayakawa et al., 2007; Sharma et al., 2008] and planetary index  $(K_p)$ . It is concluded that there is no relation between enhancements in intensity of magnetic field and geomagnetic activity.

## References

Hattori K., Akinaga Y., Hayakawa M., Yumoto K., Nagao T., Uyeda S. ULF Magnetic Anomaly Preceding the 1997 Kagoshima Earthquakes // Seismo-Electromagnetics: Lithosphere-Atmosphere-Ionosphere Coupling / Eds. M. Hayakawa, O. A. Molchanov. — 2002. — P. 19—28.

Hayakawa M., Hattori K., Ohta K. Monitoring of ULF (ultra-low-Frequency) Geomagnetic Variations Associated with Earthquakes // Sensors. — 2007. — 7. — P. 1108—1122.

Hayakawa M., Ohta K. M., Nickolaenko A. P., Ando Y. Anomalous Effect in Schumann Resonance Phenomena Observed in Japan Possibly Associated with the Chi-Chi Earthquake in Taiwan // Annales Geophys. —- 2005. — 23. — P. 1335—1346.

Sharma A. K., Patil A. V., Haridas R. N., Bhosale R. V. ULF emissions associated with seismic activity recorded at Kolhapur Station // Current Sci. — 2008. — 94, № 8. — P. 25.