

заболевания на психоэмоциональное состояние пациентов.

Ключевые слова: акне, психоэмоциональное состояние, CADI, наследственная предрасположенность.

O.I. Litus, A.V. Petrenko

Comparison of the Psycho-Emotional State of Patients with Acne with the Severity of Hereditary Factors

Shupyk National Medical Academy of Postgraduate Education, Kyiv, Ukraine

Abstract. Acne is one of the most common skin disease in young adults and occurs in 85% of boys and girls. The disease is characterized by progressive course of the formation of psycho-emotional disorders, mainly depression row in about half of patients. Installed heredity is almost 80% of first-degree relatives of patients with acne. In patients with positive family history of acne the disease occurs earlier and course harder.

The aim of the study – is to determine the impact of acne on psycho-emotional status in patients.

Materials and methods. We examined 78 patients with severe

and moderate acne. All patients were questioned by questionnaire CADI (Cardiff Acne Disability Index).

Results. Among the surveyed men with moderate acne average value was 6, median – 6, mode – 6; in men with severe acne course average value was 8.6, median – 9.5, mode – 10. Among women surveyed the results of the form were somewhat different from data that were shown by male patients. Women with moderate acne had average value 7 points, the median – 7.5 mode – 8; among women who had severe acne course the average value was – 8, median – 9.

Conclusions. As can be seen from the results of the questionnaire, respondents in all groups were shown a significant effect of disease on the psycho-emotional state of patients.

Keywords: *acne, psycho-emotional state, CADI, a genetic predisposition.*

Надійшла 01.06.2017

Завершено рецензування 12.06.2017

Прийнято до друку 15.06.2017

DOI: 10.21802/gmj.2017.2.4

I.M. Luchko, T.V. Guranych, N.M. Voronych-Semchenko, O.O. Shlyakhovenko, L.S. Storozhuk

Some Structural and Chemical Changes in Endocardial Endothelium of Rats in Emotional and Pain Stress Complicated by Hypercholesterolemia

Department of Physiology

Ivano-Frankivsk National Medical University, Ivano-Frankivsk, Ukraine

E-mail: fisiology@ifnmu.edu.ua

Abstract. The objective of the research was to study the content of some neutral lipids of endocardial endothelium in rats in relation to structural changes occurring in it, in the co-existence of emotional and pain stress, as well as alimentary hypercholesterolemia.

Materials and methods. The electric-impulse model was used for stress modeling. Alimentary hypercholesterolemia was modeled feeding animals an atherogenic diet. The concentration of triacylglycerols, free and esterified cholesterol were examined using the method of thin-layer chromatography performed on silica gel. The concentration of free fatty acids was determined using the radiochemical method. The state of endocardial endothelium was studied with the help of light microscopy; the impression smears obtained from macro preparations of ventricle were analyzed.

Results. In co-existence of stress and hypercholesterolemia, significant increase in free cholesterol as well as free fatty acid concentration was noticed. This essentially exceeded the analogical indices under the action of stress only. Structural changes in the endocardium followed by desquamation of separate endotheliocytes were the result of stress reaction. In the action of both pathogenic factors, this process was intensified; layer-by-layer exfoliation of endotheliocytes was observed.

Conclusions. In acute emotional and pain stress, changes in lipid spectrum of membrane structures of endocardial endotheliocytes the main manifestation of which is the accumulation of free cholesterol in cells and increase in the levels of free fatty acids take place. The increase in the number of desquamated endothelial cells is the result of stress action as well. Alimentary hypercholesterolemia significantly increases such pathological changes.

Keywords: *endothelium; endocardium; stress; hypercholesterolemia; lipids*

Problem statement and analysis of the recent research

Among various etiological factors possessing a significant damaged influence on the endothelium of the heart and vessels, stress situations of different genesis as well as increased free cholesterol levels are considered as one of the most important [2, 6, 7, 11]. These factors cause the development of endothelial dysfunction, which is the key link in the pathogenesis and clinical manifestations of most cardiovascular diseases [4, 5, 12]. Despite numerous experimental and clinical studies, some questions of the pathogenic effect of stress, and hypercholesterolemia (HCE) on morphology and function of cardiac endothelial lining, endothelial cell of the endocardium in particular, are not sufficiently studied. Damaging action of any stress factor on the heart is proven to be followed by activation of lipolysis, elevated levels of free fatty acids (FFA), degradation of membrane phospholipids with further accumulation of their isoforms, as well as intracellular accumulation of double-stranded fatty acids and acetyl coenzyme A [1, 6]. Since lipids are the basis of all biological membranes, the aforementioned disorders can obviously lead to significant morphological changes in different structures of cardiac endotheliocytes and vessels of mammals. In HCE, the excessive accumulation of cholesterol (C) in

membranes of these cells is possible; as a result, the relationship between different classes of lipids is disturbed [2, 3]. It can significantly change the reaction of endotheliocytes to the action of neurohumoral factors during stress. At the same time, the literature data concerning the action of stress factors on lipid spectrum and morphological reconstruction of cardiac endothelial lining in HCE are quite limited.

The objective of the research was to study the content of some neutral lipids of endocardial endothelium in rats in relation to structural changes occurring in it, in the co-existence of emotional and pain stress (EPS), as well as alimentary hypercholesterolemia.

Materials and methods

The research was carried out on 22 nonlinear white male rats weighting 180-240 g, that were divided into three groups: Group I – the control group, Group II – rats, that were under the action of EPS for three hours, Group III – animals that were under the action of EPS on the background of HCE. The electric-impulse model of Desiderato O. was used for modelling EPS [8]. Alimentary HCE was modeled by daily administration of 250 mg of cholesterol in 1 ml of lard with obligatory adding of bile acids in the oral cavity using a probe. Rats were fed atherogenic diet for 60 days.

Animals were kept and manipulated in accordance with “European Convention for the Protection of Vertebrate Animals used for Experimental and Other Scientific Purposes” (Strasbourg, 1986), and “General Ethical Principles of Experiments on Animals”, approved by the First National Congress of Bioethics (Kyiv, 2001). Euthanasia was carried out under nembutal anesthesia at a dose of 40 mg/kg of animal's weight. Lipids were extracted from the endocardial and myocardial tissues using a mixture of chloroform and methanol in the ratio of 2:1. The eluates were evaporated in an atmosphere of nitrogen and the method of thin-layer chromatography performed on silica gel (Woelm) was used. Free fatty acids, triacylglycerols (TAG) (Serva, Germany), cholesterol (Reanal, Hungary), cholesterol oleate (Amersham, England) were used as standards. The quantitative determination of fractions was made with the help of test-kits “Hemapol” (Czech Republic). The concentration of free fatty acids was determined applying the radiochemical method. For determination of cholesterol in blood plasma, there was used the qualitative reaction of Lieberman-Burchard which consists in adding a few drops of acetic anhydride and concentrated sulphuric acid to the chloroform solution of the sterol-formation of a green colour after a few minutes is positive.

To study the state of endocardial endothelium, the impression smears obtained from macro preparations of the left and right ventricles were prepared. Ventricular walls of the endocardium were previously air-dried. Micro preparations were stained using a Romanovsky method. The degree of damage to endocardial endothelium was determined with the help of light microscope ($E \times 630$), according to the number of desquamated cells per 1 mm² of smear square, that equals to 40 fields of view. The obtained data were statistically processed with the help of mathematical programs Microsoft Excel, and Statistica 7.0.

Results and discussion

The results of the research and their analysis showed, that acute three-hour EPS was followed by significant changes in the concentration of different classes of neutral lipids in endocardial endothelium (Table 1) being one of the signs of sympathoadrenal system activation, as well as the elevation of catecholamine levels in the blood [1].

Catecholamines are known to play a key role, since they have a strong lipomobilizing effect which is enhanced by glucocorticoids, adrenocorticotrophic hormone and glucagon [1, 2, 10]. In addition, catecholamines increase the intensity of processes disintegrating membranes and promote lipid peroxidation. In stress, in rat endocardium, the level of non-esterified C and FFA increased by 13.9% ($p < 0.01$) and 36.8%,

Table 1. Lipid concentration in endocardial endotheliocytes of rats in emotional and pain stress and emotional and pain stress in case of hypercholesterolemia ($M \pm m$)

Research object	№	Experimental conditions	Indices			
			Free C, mkg/g of dry tissue	Esterified C, mkg/g of dry tissue	FFA, mkg/g of dry tissue	TAG, mkg/g of dry tissue
Endo-cardium	I	Control group, n=6	37.4±0.78	25.4±0.57	1.9±0.13	17.0±0.41
	II	EPS, n=8	42.6±0.89 $p_{1,2} < 0.01$	20.9±0.73 $p_{1,2} < 0.001$	2.6±0.08 $p_{1,2} < 0.001$	15.1±0.62 $p_{1,2} < 0.05$
	III	EPS on the background of HCE, n=8	49.5±1.16 $p_{1,3} < 0.001$ $p_{2,3} < 0.001$	33.6±0.82 $p_{1,3} < 0.001$ $p_{2,3} < 0.001$	3.0±0.14 $p_{1,3} < 0.001$ $p_{2,3} < 0.05$	16.4±0.62 $p_{1,3} > 0.05$ $p_{2,3} < 0.2$

respectively ($p < 0.001$), while the level of etherified C and TAG decreased by 17.7% ($p < 0.001$) and 11.2% ($p < 0.05$). Such changes are obviously associated with the intensification of C and TAG ester hydrolysis under the influence of cholesterolases and triacylglycerollipases, the activity of which increases in stress reaction [1].

The results of the determination of the indices under the action of stress in animals being previously fed an atherogenic diet for 2 months were slightly different (Table 1). It is worth mentioning that in rats being fed an atherogenic diet, the level of total C in blood plasma increased by 1.8 time ($p < 0.001$) as compared to the control group and the level of high-density lipoprotein C decreased by 27.4% ($p < 0.001$), being a sign of elevation of plasma atherogenicity.

In the co-existence of stress and HCE, in the endocardium, there was observed a significant increase in the level of free C (by 32.4%, $p < 0.001$) exceeding the analogical index during the action of EPS only by 13.9% ($p < 0.001$). Thus, stress on the background of atherogenic diet considerably increases the accumulation of free C in membrane structures of endotheliocytes of rats. The increase in the levels of esterified C by 32.3 % ($p < 0.001$) was observed as well. Moreover, in this group of animals, a significant accumulation of FFA in the examined structure took place. In the endocardium, their concentration increased by 57.9% ($p < 0.001$) while under the action of EPS only, their level increased by 36.8% ($p < 0.001$). The endogenous source of FFA in this case, according to some authors, can be phospholipids, TAG and esters of C [10].

Summarizing the obtained data, we can conclude that stress on the background of alimentary HCE is very dangerous, since it is accompanied by the accumulation of free C and FFA in plasma membranes of endocardial endothelial cells which is one of the molecular mechanisms of their damage and can lead to formation of endothelial bodies (cytoplasts) and desquamation. Some other mechanisms including calcium, electrolyte-osmotic and acidic ones being activated by neurohumoral component of stress are involved in endotheliocyte damage [1, 4]. Stressful situations result in the occurrence of endogenous HCE, which together with exogenous HCE deepens the damaging effect [2, 9].

The analysis of the impression smears obtained from ventricular endocardium confirmed the increase in endotheliocyte damage through endothelial desquamation under the action of EPS on the background of HCE (Table 2).

After acute three-hour stress, the number of desquamated endotheliocytes in the impression smears obtained from the endocardium of the right ventricle and the left ventricle increased by 4.3 times ($p < 0.001$), while on the background of HCE, such factor led to the increase of this index by 6.4-6.5 times ($p < 0.001$) as compared to the control group of animals. Clusters of 20 and more desquamated endotheliocytes could be seen in the

Table 2. Number of endotheliocytes in impression smears (cells/mmI) obtained from ventricular endocardium in emotional and pain stress and emotional and pain stress in case of hypercholesterolemia (M±m)

№	Groups of animals and experimental conditions	Research object	
		Endocardium of the right ventricle	Endocardium of the left ventricle
I	Control group, n=6	22±1.0	24±1.7
II	EPS, n=8	94±3.5 $p_{1-2}<0.001$	102±5.6 $p_{1-2}<0.001$
III	EPS on the background of HCE, n=8	140±5.8 $p_{1-3}<0.001$ $p_{2-3}<0.05$	156±8.2 $p_{1-3}<0.001$ $p_{2-3}<0.001$

microscopes field of view (Fig. 1) indicating their possible layer-by-layer exfoliation.

Thus, alimentary HCE considerably potentiates the damaging effect of stressor on the endothelial layer of ventricular endocardium which once more confirms the danger of such factor combination.

Conclusions

1. In acute emotional and pain stress, changes in lipid spectrum of membrane structures of endocardial endotheliocytes the main manifestation of which is the accumulation of free C in cells and increase in the levels of FFA take place.

2. Along with changes in lipid composition, the effect of EPS, and, especially its combination with HCE, is accompanied by structural reconstruction of endothelial cells, the extreme manifestation of which is desquamation of the latter.

Prospects for further research

The obtained results, as well as the data of another experimental studied, allow us to talk about the systemic character of damage to cardiac endothelium and vessels of different caliber [3, 7, 9] in response to the action of one of the atherogenic factors – exogenous cholesterol. Therefore, in our opinion, future study of the effects of stressor and hypercholesterolemia on the structure and function of vascular endothelium of various organs will promote the discovery of new links in the pathogenesis of endothelial dysfunction in case of different diseases.

References

1. Baraboy VA. Stress: priroda, biologicheskaya rol, mekhanizmy, iskhody. Kyiv: Fitotsenter; c2006. 424p.
2. Bratus VV, Shumakov VA, Talaeva TV. Ateroskleroz, ishemicheskaya bolezn serdtsa, ostryy koronarnyy sindrom: patogenez, diagnostika, klinika, lecheniye. Kyiv: Chetverta khvylia; c2004. 575 p.
3. Vatsaba MO. Kliniko-diahnostychnye znachennia endoteliinoii dysfunktsii ta arteriinoii zhorstkosti u khvorykh na arteriinu hipertenziiu u poiednanni iz ozhyrinniam ta podahroi. Lvivskiy medychniy chasopys. 2014;20(1):48-52.
4. Vozna KhI, Moskaliuk VD. Endothelium: functional properties and dysfunction. Infektsiini khvoroby. 2015;1:66-71. [published in Ukrainian]
5. Zaprovalna OYe. Vplyv endotelialnoii dysfunktsii na osnovni chynnyky ishemichnoii khvoroby sertsia u patsientiv riznykh vikovykh hrup. Krovoobih ta hemostas. 2015;1:86-87.
6. Ilenko M. Dyferentsiina rol stresiv u vynykenni khvorobyvykh syndromiv. Osvita rehionu. Politolohiia. Psykholohiia. Komunikatsiia. 2011;1:272-276.
7. Kreminska IB, Levytskyi VA. Strukturno-metabolichni aspekty systemnoii reaktsii sudynnoho endoteliu pry eksperymentalnii




Fig. 1. Endotheliocytes in impression smears obtained from the endocardium of the left ventricle of rats:

A – the control group;

B – after three-hour stress;

C – after the action of three-hour EPS on the background of HCE Mg: × 630.

Notes: 1 – separate endotheliocytes, 2 – layer of endotheliocytes

hiperkholesternemii, fizychnykh navantazhenniakh riznoii intensyvnosti ta yikh poiednanni. Arkhiv klin. medytsyny. 2007;1(11):22-25.

8. Levytskyi VA, Luchko IM. Biokhimichni ta morfolohichni proiavy endotelialnoi dysfunksii endokarda i sudyn miokarda pry emotsiino-boliovomu stresi u poiednanni z hiperkholesternemiieiu. Fiziolohichniy zhurnal. 2008;54(5):75-80.

9. Prsyazhna OD, Sahach VF. Rol mitokhondrialnykh bilkiv-roziednuvachiv u porushenni endoteliizaleznykh skorochuvalnykh reaktsii sertsia ta sudyn za umov eksperymentalnoho tsukrovoho diabetu. Fiziolohichniy zhurnal. 2008;54(1):10-17.

10. Sadyak OV. Endothelium as regulator immune, inflammatory, and free radicals processes mediated L-arginini system – nitric oxide.

Medychna khimii. 2013;15(4):96-99. [published in Ukrainian]

11. Khorosh MV, Golovanova IA. Modernization of lifestyle as a way of primary prevention of cardiovascular diseases. Zdorovia natsii. 2016;1:212-217. [published in Ukrainian]

12. Abdelhamed AI, Reis SE, Sane DC. No effect of an L-arginine-enriched medical food (HeartBars) on endothelial function and platelet aggregation in subjects with hypercholesterolemia. Anj. American Heart Journal. 2008;145(3):492. doi: 10.1067/mhj.2003.160

Received: 16.04.2017

Revised: 22.05.2017

Accepted: 31.05.2017

DOI: 10.21802/gmj.2017.2.5

Z.R. Ozhohan, A.V. Biben

Clinical Effectiveness of Using Aesthetic Fixed Prosthetic Appliances with Combined Occlusal Surface

Department of Prosthetic Dentistry

Ivano-Frankivsk National Medical University, Ivano-Frankivsk, Ukraine

E-mail: Andrijko07@gmail.com

Abstract. The objective of the research was to evaluate the clinical effectiveness of using aesthetic fixed prosthetic appliances with combined occlusal surface.

Materials and methods. The study included 30 patients who were divided into 2 groups: Group I included 20 patients with combined occlusal surface of the crowns; Group II included 22 patients with ceramic occlusal surface of the crowns. The patients were observed 3, 6 and 12 months after prosthetic repair.

Results. 6 months after prosthetic repair, the occlusal contact surface area in Group I increased by 0.79%, while in Group II, it increased by 1.4%. 12 months after prosthetic repair, the occlusal contact surface area in Group I and Group II increased by 2.37% and 5%, respectively. The aforementioned data indicated that the occlusal contact surface area increased as the period of ceramic fixed prosthetic appliance wearing increased. When combining the occlusal surface in dental prostheses, the occlusal contact surface area increased approximately 2 times slower. The study demonstrated convincingly that crowns burnt on a zirconium dioxide frame were a better choice as compared to cast framework made of nickel-cobalt-chrome alloy. In case of cast frameworks, the indicators of crown margin discrepancy were worse than in case of milled frameworks - 15% versus 5%, respectively. In addition, one cast framework crown (5%) in Group II was remodeled and one crown (5%) was recemented.

Conclusions. Aesthetic fixed prosthetic appliances with combined occlusal surface demonstrated high functional and aesthetic characteristics. The use of the USHPS system showed a decisive advantage of milled frameworks and combined occlusal surface over traditional cast ceramic frameworks. The combination of high mechanical, strength and tribological properties of zirconium dioxide and high biological as well as aesthetic properties of ceramic materials helped reveal high clinical characteristics of aesthetic appliances with combined occlusal surface.

Keywords: prosthetic appliances; combined occlusal surface; zirconium dioxide; ceramics

Problem statement and analysis of the recent research

Despite the many advantages of using ceramic materials, there are some disadvantages including their propensity for chipping

as well as a high degree of the abrasion of the antagonistic teeth [1].

In prosthetic dentistry, zirconium dioxide has been used for manufacturing dental prostheses relatively recently. Zirconium dioxide has an obvious advantage over ceramic materials in its biological, physical and tribological properties. It has weak spots as well; it is mainly due to its low aesthetic characteristics.

However, when veneering zirconium dioxide framework with ceramics, aesthetic characteristics of such prosthetic appliances are even higher compared to traditional porcelain-fused-to-metal restorations [2], since metal-free ceramics shows the optical properties as those of natural teeth; translucency and opalescence are the major ones [3]. There is no doubt that metal-free ceramics is the choice of the future.

When veneering the occlusal surface of prosthetic appliances with ceramics - its disadvantages are observed – high abrasion of the antagonistic teeth as well as the probability of chipping. It leads to the increase in the occlusal surface area resulting in periodontal tissue overload, functional overloading of the masticatory muscles and their parafunction.

Therefore, we have proposed the design of a fixed prosthesis where the framework and interocclusal spaces between the antagonistic teeth were made of zirconium dioxide and all the other surfaces were veneered with ceramics. It provides a combination of positive characteristics of both construction materials. High aesthetic qualities of ceramics as well as its biological inertness combine with strength and tribological properties of zirconium dioxide.

The objective of the research was to evaluate the clinical effectiveness of using aesthetic fixed prosthetic appliances with combined occlusal surface.

Materials and methods

The study included 30 patients who were divided into 2 groups. Group I included 20 patients with combined occlusal surface of the