

## КОРОТКІ ПОВІДОМЛЕННЯ

DOI: 10.21802/gmj.2019.2.11

*Arda Isik, Eray Kurnaz, Nurten Arslan Isik*

### **Intermammary Pilonidal Disease**

Erzincan University, Turkey

kararda@yahoo.com

Pilonidal disease is an abnormal skin growth that contains hairs. It occurs mainly in the sacrococcygeal area; however, it may rarely occur in the axilla, groin, interdigital web space or umbilicus. In the literature, there were only 13 cases pilonidal

chronic discharging sinus in the intermammary area for six month. The examination revealed that the patient was hirsute and had multiple sinuses in the intermmarian region (Fig. 1).

Serous discharge was present in the patient; however, there was no abcess formation. The patient was treated with antibiotics and curettage of the sinus epithelium. After one week, the discharge stopped. During the 6-month follow-up, there was no recurrence.



**Fig. 1. Multiple sinuses in the intermmarian region**

### **References**

1. Seerwan Hama Shareef, Tahir A Hawrami, Abdulwahid M Salih et al. Intermammary pilonidal sinus: The first case series. *Int J Surg Case Rep.* 2017;41:265-268. doi: 10.1016/j.ijscr.2017.10.021
2. Anil Sunkara, DD Wagh, Sameer Harode. Intermammary Pilonidal Sinus. *Int J Trichology.* 2010;2:116-118. doi: 10.4103/0974-7753.77526

disease in the intermammary region [1, 2].

Here we present a case of a 22-year-old female. She had

Received: 21.05.2019

Revised: 13.06.2019

Accepted: 14.06.2019