

## ABSTRACT&amp;REFERENCES

DOI: 10.15587/2519-4852.2018.140756

## DEVELOPMENT OF THE METHODOLOGICAL APPROACH OF OBTAINING PREPARATIONS BASED ON SOLID DISPERSIONS

p. 4-8

**Inna Kovalevska**, PhD, Associate Professor, Department of Industrial Technology of Drugs, National University of Pharmacy, Pushkinska str., 53, Kharkiv, Ukraine, 61002

E-mail: inga.kovalevska@gmail.com

ORCID: <http://orcid.org/0000-0001-5610-8334>

**Olena Ruban**, Doctor of Pharmaceutical Sciences, Professor, Head of Department, Department of Industrial Technology of Drugs, National University of Pharmacy, Pushkinska str., 53, Kharkiv, Ukraine, 61002

E-mail: ruban\_elen@ukr.net

ORCID: <http://orcid.org/0000-0002-2456-8210>

*Nowadays, the innovative direction of pharmaceutical development is the application of methods to increase the biopharmaceutical efficiency and safety of the use of already existing APIs, including method of incorporating them into solid dispersions in which the substance molecules are included in the supramolecular formations by forming non-covalent bonds by excipients. Today, the preparation of solid dispersions is considered as the most effective way of increasing oral bioavailability, in which dissolution of the carrier facilitates the release of API with its rapid solubilization. According to the literature, it is established that today there is only an empirical approach to obtaining solid dispersions.*

*The aim of the work was the formulation of the main approaches to the development of preparations on the basis of solid dispersions, taking into account the current trends in the development of the composition and technology of preparations with sparingly soluble active pharmaceutical ingredients.*

**Materials and methods.** To develop a methodological approach, external situational content analysis of applied methods for increasing the bioavailability of active pharmaceutical ingredients used in the manufacture of solid dosage forms was used.

**Results.** In the course of accomplishing this goal, a classification of the types of solid dispersions was proposed depending on the structure and method of production, an algorithm was developed for choosing a method for obtaining a solid dispersion, taking into account the physico-chemical properties of the active pharmaceutical ingredients, which can be used in the technological process to increase their bioavailability. Critical components were also found in the development of solid dispersions.

**Conclusions.** Taking into account the physicochemical properties of the active pharmaceutical ingredients, was chosen an algorithm for selecting a method for preparing solid dispersions that can be used in the process to improve their bioavailability. Based on the results of the analysis of the literature data and our own research of the physico-chemical characteristics of the active pharmaceutical ingredients and excipients, type of the solid dispersion structure, the type of solvent and carrier, a methodological approach to the preparation of solid dispersions is proposed, the use of which will optimize the development of solid dosage forms

**Keywords:** solid dispersions, bioavailability, method of obtaining, structure, methodological approach, development stages

## References

1. Chaudhary, A., Nagaich, U., Gulati, N., Sharma, V. K., Khosa, R. L. (2012). Enhancement of solubilization and bioavail-

ability of poorly soluble drugs by physical and chemical modifications: A recent review. *Journal of Advanced Pharmacy Education & Research*, 2 (1), 32–67. Available at: <https://pdfs.semanticscholar.org/1e5b/91e6a3964fa84e49b901051d5ca4bccc620d.pdf> Last accessed: 06.03.2018

2. Krasnyuk, I. I., Stepanova, O. I., Beliatskaya, A. V., Krasnyuk, I. I., Korol, L. A. (2015). Perspektivy polucheniya lekarstvennykh form na osnove tverdykh dispersiy furatsilina. *Razrabotka i registratsiya lekarstvennykh sredstv*, 11. Available at: <https://pharmjournal.ru/articles/stati/perspektivy-polucheniya-lekarstvennyh-form-na-osnove-tvyordyh-dispersij-furacilina-n11-maj-2015> Last accessed: 30.03.2017

3. Grass, M. (2017). Selecting In Vitro Dissolution Tests for Bioavailability Enhancing Oral Formulations. Available at: <https://www.americanpharmaceuticalreview.com/Featured-Articles/343542-Selecting-em-In-Vitro-em-Dissolution-Tests-for-Bioavailability-Enhancing-Oral-Formulations/> Last accessed: 11.03.2018

4. Huang, Y., Dai, W.-G. (2014). Fundamental aspects of solid dispersion technology for poorly soluble drugs. *Acta Pharmaceutica Sinica B*, 4 (1), 18–25. doi: <http://doi.org/10.1016/j.apsb.2013.11.001>

5. Mir, K. B., Khan, N. A. (2017). Solid dispersion: overview of the technology. *International Journal Of Pharmaceutical Sciences And Research*, 8 (6), 2378–2387. doi: [http://doi.org/10.13040/ijpsr.0975-8232.8\(6\).2378-87](http://doi.org/10.13040/ijpsr.0975-8232.8(6).2378-87)

6. Singh, S., Singh Baghe, R., Yadav, L. (2011). A review on solid dispersion // *International Journal of Pharmacy & Life Sciences*, 2 (9), 1078–1095. Available at: <http://www.ijplsjournal.com/issues%20PDF%20files/sep2011/11.pdf> Last accessed: 02.03.2018

7. Harris, R. (2016). Solid Dispersions. A universal formulation strategy for poorly soluble drugs? *Contract Pharma*. Available at: [https://www.contractpharma.com/issues/2016-04-01/view\\_features/solid-dispersions-a-universal-formulation-strategy-for-poorly-soluble-drugs](https://www.contractpharma.com/issues/2016-04-01/view_features/solid-dispersions-a-universal-formulation-strategy-for-poorly-soluble-drugs) Last accessed: 14.04.2018

8. Dhirendra, K., Lewis, S., Udupa, N., Atin, K. (2009). Solid Dispersions: A Review. *Pakistan journal of pharmaceutical sciences*, 22 (2), 234–246. Available at: [https://www.researchgate.net/publication/24250932\\_Solid\\_Dispersions\\_A\\_Review](https://www.researchgate.net/publication/24250932_Solid_Dispersions_A_Review) Last accessed: 02.03.2018

9. Kovalevska, I. V., Ruban, O. A., Hrudko, V. A. (2015). Doslidzhennia vyvlnennia kvartetnyu z tverdykh dyspersii vysokomolekuliarnykh rehovyn. *Zbirnyk naukovykh prats spivrobotnykiv NMAPO im. P. L. Shupyka*, 24 (5), 318–322. Available at: [http://nbuv.gov.ua/UJRN/Znpsnmapo\\_2015\\_24%285%29\\_62](http://nbuv.gov.ua/UJRN/Znpsnmapo_2015_24%285%29_62)

10. Kovalevska, I. V., Grudko, V. A. (2014). Doslidzhennia tysku na rozchynnist kvartetnyu. *Ukrainskyi visnyk psykhonevrolohii*, 22 (2 (79)), 275–277.

11. Shmyreva, Yu., Shmal'ts, D., Bryuneman, E. (2015). Povyshenie rastvorimosti metodom tverdoy dispersii s ispol'zovaniem proizvodnykh tsellyulozy: kontseptsiya, protsessy i obezpechenie kachestva. *Farmatsevticheskaya otrasl*, 6 (53), 70–73. Available at: <http://promoboz.com/uploads/articles/372.pdf>

12. Alexander, A., Tiwle, R. A., Giri, T. K., Tripathi, D. K., Jain, V. (2012). An Exhaustive Review on Solubility Enhancement for Hydrophobic Compounds by Possible Applications of Novel Techniques. *Trends in Applied Sciences Research*, 7 (8), 596–619. doi: <http://doi.org/10.3923/tasr.2012.596.619>

13. Kovalevska, I. V., Ruban, E. A., Kutsenko, S. A., Kutova, O. V., Kovalenko, S. M. (2017). Study of physical and chemical properties of solid dispersions of quercetin. *Asian Journal of Pharmaceutics*, 11 (4), 805–809.

DOI: 10.15587/2519-4798.2018.135132

**BIO-BASED SUCCINIC ACID SAMPLE PREPARATION AND DERIVATIZATION PROCEDURE OPTIMISATION FOR GAS CHROMATOGRAPHY-MASS SPECTROMETRY ANALYSIS**

p. 9-13

**Laurynas Jarukas**, PhD, Department of Analytical and Toxicological Chemistry, Medical Academy of Lithuanian University of Health Sciences, Eivenių g., 4, Kaunas, Lithuania, LT-50161

E-mail: laurynas.jarukas@lsmuni.lt

**Justina Kamarauskaitė**, Department of Pharmacognosy, Medical Academy of Lithuanian University of Health Sciences, Eivenių g., 4, Kaunas, Lithuania, LT-50161

E-mail: justina.kamarauskaite@fc.lsmuni.lt

**Mindaugas Marksa**, PhD, Department of Analytical and Toxicological Chemistry, Medical Academy of Lithuanian University of Health Sciences, Eivenių g., 4, Kaunas, Lithuania, LT-50161

E-mail: mindaugas.marksa@lsmuni.lt

**Sonata Trumbeckaitė**, PhD, Associate Professor, Department of Pharmacognosy, Medical Academy of Lithuanian University of Health Sciences, Eivenių g., 4, Kaunas, Lithuania, LT-50161

E-mail: sonata.trumbeckaite@lsmuni.lt

**Rasa Baniienė**, PhD, Associate Professor, Neuroscience Institute, Lithuanian University of Health Sciences, Eivenių g., 4, Kaunas, Lithuania, LT-50161

E-mail: rasa.baniene@lsmuni.lt

**Liudas Ivanauskas**, PhD, Associate Professor, Department of Analytical and Toxicological Chemistry, Medical Academy of Lithuanian University of Health Sciences, Eivenių g., 4, Kaunas, Lithuania, LT-50161

E-mail: liudas.ivanaukas@lsmuni.lt

*This study focused on bio-based succinic acid sample preparation and derivatization conditions optimization using GC-MS analytical method. Succinic acid, the precursor of a wide range bio-compounds, especially it is important in accumulation of mitochondrial metabolite succinate (citric acid cycle) and during ischemia controls reperfusion injury through mitochondrial reactive oxygen production. Accurate determination of analytes is the key in metabolomics to use as low molecular biomarkers in case to improve diagnostic methods.*

**Methods.** Gas chromatography-mass spectrometry (GC-MS) method. For the quantitative determination of the succinic acid applied derivatization process by silylation using -bis- (trimethylsilyl) -trifluoroacetamide (BSTFA).

**Results.** The derivatization agent BSTFA, the derivatization time of 3-4 hours and derivatization temperature at 70 °C were selected as the optimal derivatization condition for quantification of succinic acid by GC/MS in biological samples. The results show that GC-MS SIM method with evaporation was the most effective to quantify succinate in biological samples after ischemia/reperfusion injury. Selected ion monitoring (SIM) allowed to monitor a subset of fragments with their related mass values in a certain retention time (RT) range for a set of targets.

**Conclusions.** DC – MS has several advantages for measurements of succinate concentration in small kidney tissue samples (lyophilized mitochondria). The method can be applied in small pieces of tissue – biopsy samples, tissues from various organs

**Keywords:** succinic acid, gass chromatography-mass spectrometry, derivatization, BSTFA, metabolomics, GC-MS

**References**

- Orata, F. (2012). Derivatization Reactions and Reagents for Gas Chromatography Analysis. *Advanced Gas Chromatography – Progress in Agricultural, Biomedical and Industrial Applications*. InTech. doi: <http://doi.org/10.5772/33098>
- Lynch, T. P., Grosser, A. P. K. (2000). Inlet Derivatization for the GC Analysis of Organic Acid Mixtures. *Chromatography and Separation Technology*, 13, 12–15.
- Schummer, C., Delhomme, O., Appenzeller, B., Wennig, R., Millet, M. (2009). Comparison of MTBSTFA and BSTFA in derivatization reactions of polar compounds prior to GC/MS analysis. *Talanta*, 77 (4), 1473–1482. doi: <http://doi.org/10.1016/j.talanta.2008.09.043>
- Dunn, W. B., Hankemeier, T. (2013). Mass spectrometry and metabolomics: past, present and future. *Metabolomics*, 9 (1), 1–3. doi: <http://doi.org/10.1007/s11306-013-0507-z>
- Jiye, A., Trygg, J., Gullberg, J., Johansson, A. I., Jonsson, P., Antti, H. et. al. (2005). Extraction and GC/MS Analysis of the Human Blood Plasma Metabolome. *Analytical Chemistry*, 77 (24), 8086–8094. doi: <http://doi.org/10.1021/ac051211v>
- Yip, L. Y., Yong Chan, E. C. (2013). Gas Chromatography/Mass Spectrometry-Based Metabonomics. *Proteomic and Metabolomic Approaches to Biomarker Discovery*. Elsevier, 131–144. doi: <http://doi.org/10.1016/b978-0-12-394446-7.00008-x>
- Fico, D., Margapoti, E., Pennetta, A., De Benedetto, G. E. (2018). An Enhanced GC/MS Procedure for the Identification of Proteins in Paint Microsamples. *Journal of Analytical Methods in Chemistry*, 2018, 1–8. doi: <http://doi.org/10.1155/2018/6032084>
- Peng, J., Tang, F., Zhou, R., Xie, X., Li, S., Xie, F. et. al. (2016). New techniques of on-line biological sample processing and their application in the field of biopharmaceutical analysis. *Acta Pharmaceutica Sinica B*, 6 (6), 540–551. doi: <http://doi.org/10.1016/j.apsb.2016.05.016>
- Ding, W.-H., Chiang, C.-C. (2002). Derivatization procedures for the detection of estrogenic chemicals by gas chromatography/mass spectrometry. *Rapid Communications in Mass Spectrometry*, 17 (1), 56–63. doi: <http://doi.org/10.1002/rcm.819>
- Villas-Bôas, S. G., Smart, K. F., Sivakumaran, S., Lane, G. A. (2011). Alkylation or Silylation for Analysis of Amino and Non-Amino Organic Acids by GC-MS? *Metabolites*, 1 (1), 3–20. doi: <http://doi.org/10.3390/metabo1010003>
- Chouchani, E. T., Pell, V. R., Gaude, E., Aksentijevic, D., Sundier, S. Y., Robb, E. L. et. al. (2014). Ischaemic accumulation of succinate controls reperfusion injury through mitochondrial ROS. *Nature*, 515 (7527), 431–435. doi: <http://doi.org/10.1038/nature13909>
- Ozpinar, A., Weiner, G. M., Ducruet, A. F. (2015). Succinate: A Promising Therapeutic Target for Reperfusion Injury. *Neurosurgery*, 77 (6), 13–14. doi: <http://doi.org/10.1227/01.neu.0000473807.30361.29>
- Rousova, J., Ondrusova, K., Karlova, P., Kubatova, A. (2014). Determination of Impurities in Bioproduced Succinic Acid. *Journal of Chromatography & Separation Techniques*, 6 (2). doi: <http://doi.org/10.4172/2157-7064.1000264>
- Tretter, L., Patocs, A., Chinopoulos, C. (2016). Succinate, an intermediate in metabolism, signal transduction, ROS, hypoxia, and tumorigenesis. *Biochimica et Biophysica Acta (BBA) – Bioenergetics*, 1857 (8), 1086–1101. doi: <http://doi.org/10.1016/j.bbabi.2016.03.012>

DOI: 10.15587/2519-4852.2018.141393

## SCIENTIFIC AND PRACTICAL SUBSTANTIATION OF DIRECTIONS OF MUTUAL COOPERATION OF HIGHER EDUCATIONAL INSTITUTIONS WITH EMPLOYERS OF PHARMACEUTICAL SECTOR OF HEALTH CARE INDUSTRY

p. 14-19

**Oksana Barkovska**, Assistant, Department of Organization and Economy of Pharmacy, National University of Pharmacy, Pushkinska str., 53, Kharkiv, Ukraine, 61002

E-mail: barroksa@gmail.com

ORCID: <http://orcid.org/0000-0003-2673-1226>

**Svetlana Ogar**<sup>1</sup>, PhD, Associate Professor, Department of General Pharmacy and Safety of drug, Institute of qualification improvement for pharmacists, National University of Pharmacy, Pushkinska str., 53, Kharkiv, Ukraine, 61002

E-mail: ogarsvetlana89@gmail.com

ORCID: <http://orcid.org/0000-0003-1744-951X>

**Olha Rohulia**, PhD, Associate Professor, Department of Pharmaceutical Marketing and Management, National University of Pharmacy, Pushkinska str., 53, Kharkiv, Ukraine, 61002

E-mail: rogulyaolga@gmail.com

ORCID: <http://orcid.org/0000-0001-5065-4545>

*The aim is to study, summarize and develop the directions of mutual cooperation between higher education institutions with employers of the pharmaceutical sector of the healthcare industry, which will allow making the preparation of competitive competent specialists in accordance with the requirements of the labor market.*

*Methods: to achieve the aim, methods of scientific analysis were used, in particular generalization, comparison, system and logical methods. Graphic analysis is used to visualize the generalized results of the expert survey.*

*Results: scientific and practical approaches to determining directions of mutual cooperation between higher education institutions and employers of the pharmaceutical sector of the healthcare industry are substantiated. Modern directions of cooperation between higher education institutions and employers are presented, which contribute to the improvement of the organizational and methodological foundations of interaction, the optimization of the content of education and the educational process, the improvement of the quality of practical training and employment of graduates, the development of research and innovation infrastructure. The forms of cooperation of the National University of Pharmacy (NUPh) with pharmacy institutions and pharmaceutical enterprises are systematized, the introduction of which will improve the quality of practical training of pharmacists.*

*Conclusions: based on the results of the expert survey of pharmacy specialists, it is determined that the most significant motivations for the cooperation of the National University of Pharmacy with the employers of the pharmaceutical sector of the healthcare sector are the training of specialists in accordance with the labor market needs and access to skilled labor resources, and effective forms of cooperation - mentoring and training. Based on the results of the research on the specifics and forms of cooperation, the authors carried out their systematization and defined the areas of mutual cooperation between higher education institutions and employers, using the example of the NUPh, the implementation of which contributes to improving the quality of practical training for future pharmacists and the formation of competitive graduates in accordance with labor market requirements.*

**Keywords:** pharmacy, education, employers, forms of cooperation, practical training, employment, labor market.

## References

1. Yakisna vyshcha osvita: rol partnerstv. Available at: [http://csr-ua.info/csr-ukraine/wp-content/uploads/2014/04/Research\\_Forum\\_Business-and-Universities.pdf](http://csr-ua.info/csr-ukraine/wp-content/uploads/2014/04/Research_Forum_Business-and-Universities.pdf)
2. Nazustrich VIII Natsionalnomu zizdu farmatsevtiv Ukrainy. Robotodavtsi ta universytety, yednaitesia! Available at: <https://www.apteka.ua/article/359304>
3. Ohar, S. V., Barkovska, O. Ya. (2018). Naukovo-teoretychni zasady formuvannia sotsialnoho partnerstva zakladu vyshchoi osvity z robotodavtsiamy. Farmatsevychna nauka ta praktyka: problemy, dosiahnennia, perspektyvy rozvytku: materialy II nauk.-prakt. internet-konf. z mizhnar. uchastiu. Kharkiv: NFAU, 415.
4. The Global University Employability Ranking 2016. Available at: <https://www.timeshighereducation.com/features/global-university-employability-ranking-2016>
5. Šehu, E., Dobrić, D. (2014). University-Employer Cooperation. Beijing Law Review, 05 (04), 272–282. doi: <https://doi.org/10.4236/blr.2014.54026>
6. Rakovska, N., Pavlin, S., Melink, M. (Eds.) (2012). Assessment of cooperation between higher education institutions and employers in Europe. EMCOSU, 72.
7. Pavlin, S. (2015). Considering University-Business Cooperation Modes from the Perspective of Enterprises. European Journal of Education, 51 (1), 25–39. doi: <https://doi.org/10.1111/ejed.12163>
8. Pollard, E. et. al. (2015). Understanding employers' graduate recruitment and selection practices. Institute for employment studies, 248.
9. Closing the skills gap: companies and colleges collaborating for change (2014). A report from The Economist Intelligence Unit, 28. Available at: [https://www.luminafoundation.org/files/publications/Closing\\_the\\_skills\\_gap.pdf](https://www.luminafoundation.org/files/publications/Closing_the_skills_gap.pdf)
10. Thematic University-Business forum: Forum Report. Available at: <http://www.ubforum-basquecountry.eu/>
11. Horodetska, V. I., Lebedynets, V. O., Kovalenko, S. M. (2013). Vyznachennia ta analiz vymoh robotodavtsiv do fakhivtsiv z upravlinnia yakistiu u farmatsevychnomu sektori. Upravlinnia, ekonomika ta zabezpechennia yakosti v farmatsiyi, 5, 25–31.
12. Posylkina, O. V., Kotliarova, V. H. (2017). Stvorennia praktychno-orientovanykh form realizatsiyi vyshchoi farmatsevychnoi osvity. Aktualni pytannia praktychnoi pidhotovky studentiv NFAU v Ukraini ta za kordonom: materialy nauk.-prakt. konf. z praktyky stud. NFAU ta Koledzhu NFAU. Kharkiv: NFAU, 61–63.
13. Partnerstvo biznesu, derzhavy ta universytetiv yak stratehichniy resurs innovatsiinoho rozvytku Ukrainy. Available at: <http://www.edu-trends.info/higher-edu-partnership>
14. Tarasenko, S. I., Demchenko, M. Ye. (2017). Partnerstvo universytetiv ta biznesu: formy ta perspektyvy rozvytku v umovakh pidvyshchennia innovatsiynosti ekonomiky. Ekonomika i suspilstvo, 13, 302–308. Available at: [http://www.economyandsociety.in.ua/journal/13\\_ukr/49.pdf](http://www.economyandsociety.in.ua/journal/13_ukr/49.pdf)
15. Polozhennia pro Radu robotodavtsiv Vchenoi rady fakultetiv zi spetsialnosti «Farmatsiya» u Natsionalnomu farmatsevychnomu universyteti, POL A 2.7-32-185, 2016 r.
16. Partnerstvo z robotodavtsiamy yak stratehichniy resurs rozvytku NFAU. Available at: <https://www.apteka.ua/article/409162>
17. Havrysh, N. B., Barkovska, O. Ya. (2017). Doslidzhenia form spivpratsi Natsionalnoho farmatsevychnoho universytetu z aptechnymy zakladamy ta farmatsevychnymy pidpriemstvamy Ukrainy. Relevant issues of modern medicine: the experience of

Poland and Ukraine: international research and practice conference. Lublin, 119–122.

DOI: 10.15587/2519-4852.2018.141304

### THE INFLUENCE OF INDOLINOREN ON KIDNEY FUNCTION IN CONDITIONS OF WATER AND SALT LOAD

p. 20-23

**Anna Markina**, Senior Laboratory, Department of Clinical Pharmacology, Institute of Pharmacy Professionals Qualification Improvement, National University of Pharmacy, Zahysnykyiv Ukrainy sq., 17, Kharkiv, Ukraine, 61001

E-mail: Kolomanya7@gmail.com

ORCID: <http://orcid.org/0000-0001-5784-0201>

**Oksana Mishchenko**, Doctor of Pharmaceutical Sciences, Professor, Head of Department, Department of Clinical Pharmacology, Institute of Pharmacy Professionals Qualification Improvement, National University of Pharmacy, Zahysnykyiv Ukrainy sq., 17, Kharkiv, Ukraine, 61001

E-mail: clinpharmacol\_ipksph@nuph.edu.ua

ORCID: <http://orcid.org/0000-0001-5862-4543>

*Diuretics are widely used to correct kidney disorders. In turn, the ionic composition of food, mostly sodium ions, which directly affect the excretory function of the kidneys, can act as modulators of the action of diuretics.*

*The aim of the study was to investigate the effects of the new 2-oxoindoline derivative with the conventional name "Indolinoren" on the state of the excretory function of the kidneys under the conditions of water and salt load.*

*Materials and methods. Studies were performed on white non-linear rats. The water load was modelled by introducing distilled water (5 ml per 100 g of body weight of the animal); salt load - intragastric administration of 0.45 % sodium chloride solution in an amount of 3 % of body weight. Indolinoren and comparison drug furosemide were administered intragastrically at a dose of 29.5 mg/kg and 5 mg/kg, respectively.*

*Results and discussion. It has been established that indolinoren has a saluretic effect in conditions of water load, accompanied by an increase in sodium excretion by 132 % and potassium by 2.4 %. Against the background of the introduction of indolinoren, a significant increase in the sodium-potassium coefficient of urine was established 2.3-fold ( $p < 0.05$ ), which indicates a more pronounced natriuresis than kaliuresis. Under conditions of salt load, indolinoren promotes a significant increase in urine output by 381 % ( $p < 0.05$ ), increases sodium excretion by 127 % ( $p < 0.05$ ), potassium by 7 %. There were no significant differences in creatinine excretion. The expression of diuretic activity in conditions of salt load indolinoren exceeds furosemide and does not have a significant difference under conditions of water load.*

**Conclusion:** *The increase in natriuresis and, to a lesser extent, the kaliuresis, as well as the absence of significant changes in the excretion of creatinine, a glomerular filtration marker, on the background of the introduction of indolinoren indicates that its diuretic effect is realized due to oppression of tubular reabsorption. In the mechanism of indolinoren action involved inhibition of mineralocorticoid control of the excretory function of the kidney, as evidenced by the increase in the sodium-potassium coefficient of urine. The obtained data justify the need for further in-depth study of indolinoren as a perspective diuretic*

**Keywords:** *diuretic activity, water load, salt load, indolinoren, 2-oxoindoline derivative, furosemide*

### References

- Bobrov, V. O., Davydova, I. V. (2007). Diuretyky: klasyfikatsiia, farmakokinytyka ta farmakodynamika. Kyiv: Medknyha, 100.
- Kumanyika, S. (1991). Behavioral aspects of intervention strategies to reduce dietary sodium. Hypertension, 17 (1), 190. doi: [http://doi.org/10.1161/01.hyp.17.1\\_suppl.190](http://doi.org/10.1161/01.hyp.17.1_suppl.190)
- Shtrygol', S. Yu. (2007). Modulyatsiia farmakologicheskikh effektov pri razlichnykh solevykh rezhimakh. Kyiv: Avista-VLT, 360.
- Ivanov, D. D., Korzh, O. M. (2014). Nefrolohiia v praktytsi simeinoho likaria. Kyiv: Yzdatelskyi dom "Zaslavskyi", 464.
- Zaika, M. N., Bytsia, Yu. V. (Eds.) (2008). Patolohichna fiziolohiia. Kyiv: Medytsyna, 373–384.
- Mukhin, N. A. (Ed.) (2010). Nefrologiya: neotlozhnye sostoyaniya. Moscow: Eksmo, 288.
- Babini, R., Larose, P., Leacutecrivain, A., Du Souich, P. (1991). Furosemide Dynamics: Influence of Dietary Sodium and of Saralasin. Pharmacology, 43 (5), 282–292. doi: <http://doi.org/10.1159/000138856>
- Markina, A. Yu., Mishchenko, O. Ya. (2017). Vychennia diuretychnoi aktyvnosti novykh atsylovanykh pokhidnykh 2-oksoindolinu. VHO "Asotsiatsiia farmakolohiv Ukrainy". Zaporizhzhia, 88.
- Markina, A., Astapova, N., Tyupka, T. (2013). Effects of indolinoren on renal excretory function in rats with acute renal failure. Actual questions of development of new drugs. Kharkiv, 167.
- Pro zakhyst tvaryn vid zhorstokoho povodzhennia (2006). Zakon Ukrainy No. 3447-IV. 21.02.2006. Available at: <http://zakon4.rada.gov.ua/laws/show/3447-15>
- European Convention for the protection of vertebrate animals used for experimental and other scientific purposes (1986). ETS No. 123. Strasbourg. Available at: <http://conventions.coe.int/treaty/en/treaties/html/123.htm>
- Berkhin, E. B. Ivanov, Yu. I. (1972). Metody eksperimental'nogo issledovaniya pochek i vodno-solevogo obmena. Barnaul: Altaiskie knizhn. izdatel'stvo, 199.
- Shevtsov, I. I. (2006). Pokhidni 2-oksoindolin-3-hlioksylovoi kysloty-potentsiini rehuliatory sechovydilnoi funktsii nyrok. Kharkiv, 20.
- Tymoshenko, O. P., Voronina, L. M., Kravchenko, V. M. et. al. (Eds.) (2005). Klinichni biokhimiia. Kharkiv: Vyd-vo "Professional", 288.
- Kamyshnikov, V. S. (2000). Spravochnik po kliniko-biokhimicheskoy laboratornoy diagnostike. Vol. 1. Minsk: Belarus, 495.
- Lapach, S. N., Chubenko, A. V., Babich, P. N. (2001). Statisticheskie metody v mediko-biologicheskikh issledovaniyakh s ispol'zovaniem EXEL. Kyiv: MORION, 408.
- Toma, C.-C., Olah, N.-K., Vlase, L., Mogoşan, C., Mocan, A. (2015). Comparative Studies on Polyphenolic Composition, Antioxidant and Diuretic Effects of Nigella sativa L. (Black Cumin) and Nigella damascena L. (Lady-in-a-Mist) Seeds. Molecules, 20 (6), 9560–9574. doi: <http://doi.org/10.3390/molecules20069560>

DOI: 10.15587/2519-4852.2018.141397

### DEVELOPMENT AND VALIDATION OF SPECTROPHOTOMETRIC METHOD FOR THE DETERMINATION OF MELDONIUM DIHYDRATE IN DOSAGE FORMS

p. 23-27

**Anastasiia Donchenko**, Postgraduate Student, Department of Analytical Chemistry, Zaporizhzhia State Medical University, Maiaikovskoho ave., 26, Ukraine, Zaporizhzhia, 69035

E-mail: donchenko130791@gmail.com

ORCID: <http://orcid.org/0000-0002-3511-1339>

**Natalia Nahorna**, PhD, Senior Lecturer, Department of Analytical Chemistry, Zaporizhzhia State Medical University, Maiakovskoho ave., 26, Ukraine, Zaporizhzhia, 69035

**E-mail:** natalija.nagornaja@gmail.com

**ORCID:** <http://orcid.org/0000-0001-8312-4356>

**Svitlana Vasyuk**, Doctor of Pharmaceutical Sciences, Professor, Head of Department, Department of Analytical Chemistry, Zaporizhzhia State Medical University, Maiakovskoho ave., 26, Ukraine, Zaporizhzhia, 69035

**E-mail:** svitlanavasyuk@gmail.com

**Aim.** Spectrophotometry is one of the most widely used methods in the pharmaceutical analysis. The main advantages of this absorption method are the highly sensitive, cost-effective and available to quality control laboratories for dosage forms. However, there is a need to find new analytical reagents. Therefore, the aim of the present work was investigation and development of spectrophotometric method based on reaction with *p*-chloranil for the determination of meldonium dihydrate in dosage forms.

**Methods.** In the study were used working standard of meldonium dihydrate, *p*-chloranil, DMF, the sample of finished dosage forms. Absorption of the reaction products was measured using spectrophotometer Specord 200.

**Results.** The optimum conditions of the spectrophotometric analysis has been established during the process of development this procedure. The influence of various parameters such as nature of the solvent, concentration of reagent, temperature, time of heating were investigated. It was experimentally established that meldonium dihydrate reacts with *p*-chloranil in DMF medium to form the coloured reaction product with absorption maximum at 556 nm. The proposed method was subjected to validation tests. The method was validated for the parameters like linearity, precision, accuracy, robustness and range of application. Beer's law was performed at the concentration range of 8.00–20.00 mg/100 ml with correlation coefficient 0.9995. The linearity ranges were calculated with the help of regression analysis by means of least squares. The proposed procedure meets the requirements of State Pharmacopoeia of Ukraine.

**Conclusions.** The spectrophotometric method for the determination of meldonium dihydrate was developed and validated. This procedure is successfully applied for dosage forms analysis. Results of the study showed that the procedure is accurate, simple and relevant for application at the quality control laboratories for dosage forms

**Keywords:** spectrophotometry, derivatives of quinone, *p*-chloranil, meldonium dihydrate, analysis, quantitative determination, validation

## References

1. Prykhodko, Yu. V., Kashkovskiy, D. O., Prykhodko, V. M. (2015). Metabolichna terapiia v praktytysi simeinoho likaria. Znachennia metabolichnykh preparativ v heriatrychnii klinitsi [Metabolic therapy in medical general practice. Metabolic drugs in geriatrics]. Semeynaya meditsina, 1 (57), 7–16.
2. Gorbunova, A. A., Kireev, S. Yu., Rashevskaya, I. V. (2017). Mel'doniy: svyaz' stroeniya, struktury i svoystv [Meldonium: connection of structure, structure and properties]. Vestnik Penzenskogo gosudarstvennogo universiteta, 2 (17), 92–99.
3. Dzerve, V., Kalvinsh, I. (2013). Mildronat v kardiologii. Obzor issledovaniy [Mildronate in cardiology. Research Overview]. Riga, 76.
4. MEL'DONIY (MELDONIUM). Opisanie aktivnykh veshhestv. Spravochnik Kompndium. Available at: <https://compendium.com.ua/akt/77/2984/meldonium/>
5. Gosudarstvennaya farmakopeya Rossiyskoy Federatsii. XII-izd. [State Pharmacopoeia of Russian Federation. 12th ed.]

(2008). Moscow: Izdatel'stvo «Nauchnyy tsentr ekspertizy sredstv meditsinskogo primeneniya», 704.

6. European Pharmacopoeia. 8.3th ed. (2015). Strasbourg: European Department for the Quality of Medicines, 4378.

7. Peng, Y., Yang, J., Wang, Z., Wang, J., Liu, Y., Luo, Z., Wen, A. (2010). Determination of mildronate by LC-MS/MS and its application to a pharmacokinetic study in healthy Chinese volunteers. Journal of Chromatography B, 878 (5-6), 551–556. doi: <http://doi.org/10.1016/j.jchromb.2009.12.030>

8. Pidpruzhnykov, Y. V., Sabko, V. E., Iurchenko, V. V., Zupanets, I. A. (2011). UPLC-MS/MS method for bioequivalence study of oral drugs of meldonium. Biomedical Chromatography, 26 (5), 599–605. doi: <http://doi.org/10.1002/bmc.1703>

9. Gavrilin, M. V., Mudretsova, Yu. V., Senchenko, S. P., Rozhnova, S. A. (2012). Razrabotka metodiki kolichestvennogo opredeleniya kholina al'fostserata i mel'doniya metodom kapillyarnogo elektroforeza [Development of quantification method for determination of choline alphascerate and meldonium by capillary electrophoresis]. Voprosy biologicheskoy, meditsinskoy i farmatsevticheskoy khimii, 4, 12–17.

10. Derzhavna Farmakopeia Ukrainy. Vol. 1 [State Pharmacopoeia of Ukraine. Vol. 1] (2015). Kharkiv: Derzhavne pidpriemstvo «Naukovo-ekspertnyi farmakopeinyi tsentr», 1128.

11. Ermer, J. (2015). Method Validation in Pharmaceutical Analysis: A Guide to Best Practice. Weinheim: Wiley-VCH., 440.

12. Grizodub, A. I. (2016). Standartizovannyye protsedury validatsii metodik kontrolya kachestva lekarstvennykh sredstv [Standardized procedures for the validation of drug quality control methods]. Kharkiv: Gosudarstvennoe predpriyatye «Ukrainskiy nauchnyy farmakopeynyy tsentr kachestva lekarstvennykh sredstv», 396.

DOI: 10.15587/2519-4852.2018.141411

## RESEARCH OF 1,3-OXAZOLE-4-IL-PHOSPHONIC ACID DERIVATIVE ON THE CONTENT OF FATTY ACIDS OF LIPIDS IN RATS WITH ARTERIAL HYPERTENSION

p. 28-31

**Kateryna Matskevych**, Assistant, Department of Pharmaceutical, Biological and Toxicological Chemistry, Bogomolets National Medical University, Shevchenka blvd., 13, Kyiv, Ukraine, 01601

**E-mail:** kateryna.sedko@nmu.ua

**ORCID:** <http://orcid.org/0000-0003-0776-4989>

The number of patients with arterial hypertension (AH) continues to increase. Significant negative effects of arterial hypertension are structural, metabolic and functional disorders in the tissues of the myocardium, vessels and other organs, in particular, changes in the content of fatty acids (FA) and their correlation. The purpose of the study is to investigate the change in fatty acid composition of lipids in blood serum and tissues of rats with arterial hypertension under the influence of a new original compound, 1,3-oxazole-4-yl-phosphonic acid derivative (abbreviated name – oxazole derivative).

**Materials and methods.** The studies were conducted on white, sexually mature rats. Arterial hypertension was modeled by salt load – salt drink (1 % solution of sodium chloride) with free access to it for 21 days. Animals from the 14th day received oxazole derivative at a dose of 25 mg / kg intraperitoneally, once daily, for 7 days. The content of fatty acids of centrifuged blood serum and homogenized in 0.9 % saline NaCl tissue was determined by gas chromatographic analysis.

**Results and discussion.** The administration of oxazole derivative in the background of increased blood pressure in rats did not significantly affect the amount of SFA and USFA in serum in contrast to

the group of rats with hypertension due to the tendency to restore the stearic acid content, but the changed content of linoleic and arachidonic acids practically did not differ from the values in the blank group. There was a restoration of the content of palmitic, stearic, linoleic and arachidonic acids in aorta. In heart, the change in the content of linoleic and arachidonic acids in the reverse direction compared with the blank group was established.

**Conclusions.** The administration of 25 mg/kg ( $ED_{50}$ ) of oxazole derivative intravenous intraperitoneally once daily for 7 days with simultaneous simulation of arterial hypertension by salt load did not cause any adverse changes and led to the restoration of lipid parameters of SFA, USFA and PUFA

**Keywords:** arterial hypertension, fatty acids, 1,3-oxazole-4-yl-phosphonic acid derivative, oxazole derivative, rats

## References

1. World Health Organization. Global status report on non-communicable diseases 2010 (2010). Geneva: WHO.
2. Klinicheskie ispytaniya lekarstvennykh sredstv i farmakonadzor v Ukraine (2001). Mater. nauchno-prak. seminaru 1-2 iyunya 2001 g. Kyiv: Aviacenna, 128.
3. World Health Organization. A global brief on hypertension 2013 (2013). Geneva: WHO.
4. Unification of the protocol of extrinsic medical assistance. Hypertonic crisis. Available at: <http://moz.gov.ua/nakazi-moz>
5. Arutyunov, G. P., Nedogoda, S. V., Gilyarevskiy, S. R., Baranova, E. I. et. al. (2015). Slozhnye voprosy lecheniya arterial'noy gipertenzii: vliyaniye povyshennoy chastoty serdechnykh sokrashcheniy i soputstvuyushchih zabolovaniy na vybor antigipertenzivnoy terapii v praktike kardiologa i terapevta [Complex questions of treatment of arterial hypertension: the effect of increased heart rate and concomitant diseases on the choice of antihypertensive therapy in the practice of a cardiologist and therapist]. *Ration Pharmacother Cardiol*, 11 (1), 63–67.
6. Popov, V. V., Bulanov, N. A., Ivanov, G. G. (2012). Current target of antihypertensive therapy. Data from clinical trials. Part 1. *Rational Pharmacotherapy in Cardiology*, 8 (1), 88–94. doi: <https://doi.org/10.20996/1819-6446-2012-8-1-88-94>
7. Zahorodniy, M. I., Briuzghina, T. S., Svintsytskyi, A. S. (2008). Zminy zhymnokyslotnoho spektru lipidiv u shchuriv zi spontannoiu arterialnoiu hipertenzieiu [Changes in the fatty acid spectrum of lipids in spontaneous arterial hypertension rats]. *Sertse i sudyny*, 3, 80–84.
8. Talati, M., Hemnes, A. (2015). Fatty Acid Metabolism in Pulmonary Arterial Hypertension: Role in Right Ventricular Dysfunction and Hypertrophy. *Pulmonary Circulation*, 5 (2), 269–278. doi: <https://doi.org/10.1086/681227>
9. Dovhan, R. S. (2014). Zminy vmistu zhyrnykh kyslot v miokardi ta plazmi krovi shchuriv z arterialnoiu hipertenzieiu pry zastosuvanni antyhipertenzivnykh zasobiv [Changes in the content of fatty acids in the myocardium and blood plasma of rats with arterial hypertension in the use of antihypertensive drugs]. *Visnyk problem biolohiyi i medytsyny*, 2 (3), 130–134.
10. Nizhenkovska, I. V., Sedko, V. K., Golovchenko, O. V., Golovchenko, O. I. (2018). The study of acute toxicity of 1,3-oxazole-4-yl-phosphonic acid derivative in intraperitoneal administration. *News of Pharmacy*, 1 (93), 43–48. doi: <https://doi.org/10.24959/nphj.18.2186>
11. Nizhenkovskaya, I. V., Zaychenko, A. V., Sed'ko, E. V., Golovchenko, A. V. (2018). Issledovanie vliyaniya proizvodnogo 1,3-oksazol-4-il-fosfonovoy kisloty na arterial'noe davlenie i serdechnyy ritm u krolikov [Investigation of the effect of the 1,3-oxazole-4-yl-phosphonic acid derivative on blood pressure and heart rate in rabbits]. *Recept*, 21 (1), 75–83.
12. European Convention for the Protection of Vertebrate Animals used for Experimental and Other Scientific Purposes. Available at: <http://conventions.coe.int/treaty/en/treaties/html/123.htm>
13. Pro zakhyst tvaryn vid zhorstokoho povodzhennia. Zakon vid 21.02.2006 No. 3447-IV. Verkhovna Rada Ukrainy. Available at: <http://zakon4.rada.gov.ua/laws/show/3447-15>
14. Badyal, D. K., Lata, H., Dadhich, A. P. (2003). Animal models of hypertension and effect of drugs. *Indian J. Pharmacol.*, 35 (6), 349–362.
15. Puzyrenko, A. M., Chekman, I. S., Briuzghina, T. S., Horchakova, N. O. (2013). Vplyv antyhipertenzivnykh ta metabolitnykh preparativ na zhymnokyslotnyi sklad lipidiv kardiomiotsytiv u shchuriv zi spontannoiu arterialnoiu hipertenzieiu [Influence of antihypertensive and metabolic drugs on fatty acids content of lipids in cardiomyocytes of rats with spontaneous hypertension]. *Ukrainskyi biokhimichnyi zhurnal*, 85 (4), 67–74.
16. Dovhan, R. S., Zahorodniy, M. I., Briuzghina, T. S., Horchakova, N. O. (2016). Vychennia diyi metabolitnykh preparativ na zhymnokyslotnyi sklad lipidiv tkanyn shchuriv zi spontannoiu arterialnoiu hipertenzieiu [Investigations of metabolite drugs action on fatty acid composition of lipids in tissues of rats with spontaneous hypertension]. *Visnyk VDNZU «Ukrainska medychna stomatolohichna akademiya»*, 16 (1), 202–205.
17. Nizhenkovska, I. V., Narokha, V. P., Kuznetsova, O. V., Briuzghina, T. S., Seifullina, I. Y., Martsynko, O. E., Chebanenko, O. A. (2015). Vplyv nikotynovoi kisloty ta kompleksu hermaniu z nikotynovoiu kyslotoiu (MIHU1) na zhymnokyslotnyi sklad lipidiv kardiomiotsytiv i hepatotsytiv shchuriv z eksperymentalnoiu khronichnoiu sertsevoiu nedostatnistiu [Effects of nicotinic acid and complex of germanium with nicotinic acid (MIGU-1) on lipid fatty acid composition of cardiomyocytes and hepatocytes in rats with experimental chronic heart failure]. *Farmakolohiya ta likarska toksykolohiya*, 1, 68–75.
18. Cohen, M., Sztokalo, J., Hinsch, E. (1973). The antihypertensive action of arachidonic acid in the spontaneous hypertensive rat and its antagonism by anti-inflammatory agents. *Life Sciences*, 13 (4), 317–325. doi: [https://doi.org/10.1016/0024-3205\(73\)90223-3](https://doi.org/10.1016/0024-3205(73)90223-3)
19. Renke, M., Knap, N., Tylicki, L., Rutkowski, P., Lizakowski, S., Woźniak, M., Rutkowski, B. (2013). Isoprostanes – important marker of the oxidative stress estimation in patients with chronic kidney disease. *Pol. Merkur. Lekarski*, 34 (199), 14–17.

DOI: 10.15587/2519-4852.2018.141412

THE RESEARCH OF FREE AMINOACIDS OF WATER-SOLUBLE PROTEIN-POLYSACCHARIDE COMPLEX OF OYSTER MUSHROOM PLEUROTUS OSTREATUS

p. 32-37

**Boris Varinsky**, PhD, Associate Professor, Department of Physical and Colloidal Chemistry, Zaporizhzhia State Medical University, Maiakovskoho ave., 26, Zaporizhzhia, Ukraine, 69035  
E-mail: [borisav70@yahoo.com](mailto:borisav70@yahoo.com)

**Nataliia Kucherenko**, PhD, Associate Professor, Department of Drug Technology, Organization and Economy of Pharmacy, SI "Luhansk State Medical University", Budivelnikiv str., 32, Rubizhne, Ukraine, 93012

E-mail: [tekhonology.kucherenko@gmail.com](mailto:tekhonology.kucherenko@gmail.com)

ORCID: <http://orcid.org/0000-0001-6462-9295>

**Olga Kolpakova**, Assistant, Department of Chemical and Pharmaceutical Technologies, Institute of Chemical Technologies of the East Ukrainian National University, V. Dal, Volodymyrska str.,

31, Rubizhne, Ukraine, 93009, Postgraduate Student, Department of Drug Technology, Organization and Economy of Pharmacy, SI "Luhansk State Medical University", Budivelnykiv str., 32, Rubizhne, Ukraine, 93012  
**E-mail:** okolpakova91@gmail.com

*Recently, the share of medicinal products of natural origin has increased in the pharmaceutical market of Ukraine. This is due to the fact that the pharmacological effect of natural substances provides a complex of biologically active substances, which has a high therapeutic effect and the minimum number of adverse reactions. Therefore, the study of chemical composition and standardization of natural substances is an important stage in pharmaceutical research. A promising source of active ingredients is the mushroom Pleurotus ostreatus. The chemical composition of this fungus is represented by a large number of biologically active substances, which ensures its many-sided use in folk medicine.*

**Aim.** The determination of qualitative and quantitative composition of free amino acids in the composition of water-soluble protein-polysaccharide complex (WPPSC) of oyster mushroom *Pleurotus ostreatus* was the aim of research.

**Methods.** Physicochemical methods are the most commonly used to study active substances of natural origin. This is due to the fact that they provide high informative, accurate, efficient and reproducible. The method of high performance liquid chromatography (HPLC) was used to analyze the amino acid composition of the WPPSC of oyster mushroom *Pleurotus ostreatus*. This method is based on the separation of individual components due to different adsorption capacity.

**Results.** Research of the amino acid composition of the WPPSC of oyster mushroom *Pleurotus ostreatus* showed that the complex contains 10 (7 %) free amino acids, among which 5 are essential. The detected amino acids are aliphatic, heterocyclic and aromatic compounds. They are presented by glycine, alanine, leucine, isoleucine, threonine, selenum, lysine, arginine, histidine and phenylalanine.

**Conclusion.** The qualitative and quantitative composition of the free amino acids of the WPPSC of oyster mushroom *Pleurotus ostreatus* was established. The method of determination and separation of free amino acids by the method of high-performance liquid chromatography is proposed. The obtained results are proposed for standardization of the initial substance in the process of production of medicines based on WPPSC of oyster mushroom *Pleurotus ostreatus*

**Keywords:** *Pleurotus ostreatus*, biologically active substances, amino acids, high performance liquid chromatography, standardization

#### References

1. Solodovnychenko, N. M., Zhuravlov, M. S., Kovalov, V. M. (2001). *Likarska roslynna syrovyna ta fitopreparaty*. Kharkiv: Vydvo NFAU: Zoloti storinky, 408.
2. Georgievskiy, V. P. (Ed.) (2001). *Analiticheskaya himiya v sozdani, standartizacii i kontrole kachestva lekarstvennyh sredstv*. Vol. 2. Kharkiv: «NTMT», 474.
3. Molitorus, H. P. (1994). Mushrooms in medicine. *Folia Microbiol*, 39 (2), 91–98.
4. Mariappan, S., Vinayagam, S., Durai, M. (2015). Phytochemical screening of bioactive compounds from *Pleurotus ostreatus* (jacq.fr) kumm-an wild edible mushroom. *World Journal of Pharmaceutical Research*, 4 (5), 1603–1618.
5. Mowsurni, F., Chowdhury, M. (2013). Oyster Mushroom: Biochemical and Medicinal Prospects. *Bangladesh Journal of Medical Biochemistry*, 3 (1), 23–28. doi: <https://doi.org/10.3329/bjmb.v3i1.13804>
6. TU U 10.8-02010675-001:2017 *Kompleks bilkovo-polisakharydnyi*.

7. Kucherenko, N. V., Martynov, A. V., Demianenko, V. H. (2008). Rozrobka metodyk standartyzatsiyi vodorozchynnoho bilkovo-polisakharydnoho kompleksu, otrymano z hryba *Pleurotus Ostreatus*. *Farmatsevtichnyi zhurnal*, 1, 92–95.

8. Minaieva, V. O. (2013). *Khromatohrafichnyi analiz*. Cherkasy, 284.

9. Neda, I., Vlazan, P., Oana, R., Sfarloaga, P., Grozescu, I., Segneanu, A.-E. (2012). Peptide and amino acids separation and identification from natural products. *Analytical Chemistry*, 6, 135–146. doi: <https://doi.org/10.5772/51619>

10. Konovalova, E. Yu., Stazhyla, E. N., Lebeda, A. Ph. (2010). Amino acid research of family elaeagnaceae juss. *Plants' leaves. Fitoterapiya*, 2, 60–64.

11. Moran-Palacio, E., Tortoledo-Ortiz, O., Yañez-Farías, G., Zamora-Álvarez, L., Stephens-Camacho, N., Soñanez-Organis, J. et al. (2014). Determination of amino acids in medicinal plants from Southern Sonora, Mexico. *Tropical Journal of Pharmaceutical Research*, 13 (4), 601–606. doi: <https://doi.org/10.4314/tjpr.v13i4.17>

12. Rudakov, O. B., Vostrov, I. A., Fedorov, S. V. et. al. (2004). *Sputnik hromatografista. Metody zhidkostnoy hromatografii*. Voronezh: Vodoley, 528.

13. Minazova, G. I. (2010). Highly effective liquid chromatography in the analysis of natural raw material. *Bashkirskiy himicheskii zhurnal*, 17 (4), 134–136.

14. Syrovaya, A. O., Shapoval, L. G., Makarov, V. A. et. al. (2014). *Aminokisloty glazami himikov, farmacevtov, biologov*. Vol. 1. Kharkiv: Shchedra sadiba plyus, 228.

15. Davies, J. S. (2006). *Amino Acids, Peptides and Proteins*. Cambridge: The Royal Society of Chemistry, 472.

DOI: 10.15587/2519-4852.2018.141423

#### IMPACT OF DRY EXTRACT OF GINGER ON MORPHOLOGICAL STATE OF PANCREAS OF SYRIAN GOLDEN HAMSTERS ON THE BACKGROUND OF HYPERCALORIUM DIET

p. 38-44

**Nadiia Kononenko**, Doctor of Medical Sciences, Professor, Head of Department, Department of Pathological Physiology, National University of Pharmacy, Pushkinska str., 53, Kharkiv, Ukraine, 61002  
**E-mail:** kononenkonn76@gmail.com  
**ORCID:** <http://orcid.org/0000-0002-3850-6942>

**Mariia Sorokina**, Postgraduate student, Department of Pathological Physiology, National University of Pharmacy, Pushkinska str., 53, Kharkiv, Ukraine, 61002  
**E-mail:** pathology@nuph.edu.ua  
**ORCID:** <http://orcid.org/0000-0003-4924-2409>

**Julia Larianovska**, PhD, Senior Researcher, Central Research Laboratory, National University of Pharmacy, Pushkinska str., 53, Kharkiv, Ukraine, 61002  
**E-mail:** cndl@nuph.edu.ua  
**ORCID:** <http://orcid.org/0000-0003-4164-9415>

**Valentina Chikitkina**, PhD, Associate Professor, Department of Pathological Physiology, National University of Pharmacy, Pushkinska str., 53, Kharkiv, Ukraine, 61002  
**E-mail:** valentina.chikitkina@gmail.com  
**ORCID:** <http://orcid.org/0000-0002-8277-0388>

*The aim is to study the effect of dry extract of ginger on the morphological state of the pancreas of Syrian golden hamsters against*

the background of a metabolic syndrome induced by a hypercaloric diet.

**Materials and methods.** The metabolic syndrome was modeled with the help of golden male hamsters, 20 weeks old, that had a hypercaloric diet enriched with energy sources (including 29 % fat – predominantly saturated lipids) and fructose (1 g per 100 g body weight) for 6 weeks. Dry extract of ginger in a dose of 80 mg/kg, species “Arphasetin” at a dose of 16 ml/kg and metformin tablets at a dose of 60 mg/kg were administered intragastrically once a day, starting at 4 weeks of experiment for 14 days.

On the histological sections of the pancreas, the total number of pancreatic islets in the micropreparation was determined, their area was measured, the islets were divided into small, medium and large, and the percentage of each category of pancreatic islets was determined.

**Results.** Prolonged consumption of food rich in fat and carbohydrates by hamsters leads to the development of the state of prediabetes, which is characterized by the morphological inhibition of the insular apparatus: an increase in the relative proportion of small and a decrease in the proportion of medium pancreatic islets, a decrease in availability. According to the morphological characteristics, the introduction of a dry extract of ginger at a dose of 80 mg/kg to hamsters with a metabolic syndrome restores the state of pancreatic islets, their area,  $\beta$ - and  $\alpha$ -cells of the pancreas to the level of intact animals. The pharmacological effect of the dry extract of ginger is most likely due to the phenolic compounds that make up its composition - gingerol and shogaol, which may modulate the release of insulin due to the antioxidant effect.

**Conclusions.** According to the severity of the protective action, the dry extract of ginger exceeds the reference preparations – metformin at a dose of 60 mg/kg and the species of “Arphasetin” in a dose of 16 ml/kg. The obtained results testify to the prospects of further experimental and clinical study of the pharmacological properties of dry extract of ginger with the aim of creating an effective antidiabetic phytopreparation.

**Keywords:** metabolic syndrome, high-calorie diet, dry ginger extract, pancreas, histological examination

## References

1. International diabetes federation Diabetes Atlas. 8<sup>th</sup> ed. Available at: <http://www.diabetesatlas.org>
2. Litvinova, L. S., Kiriienkova, E. V., Mazunin, I. O., Vasilenko, M. A., Fattakhov, N. S. (2015). Patogenez insulinorezistentnosti pri metabolicheskom ozhirenii. *Biomeditsinskaya khimiya*, 61 (1), 70–82.
3. O'Neill, S., O'Driscoll, L. (2014). Metabolic syndrome: a closer look at the growing epidemic and its associated pathologies. *Obesity Reviews*, 16 (1), 1–12. doi: <http://doi.org/10.1111/obr.12229>
4. Shifrin, O. S., Sokolina, I. A., Leonovich, A. E., Ashitko, M. A., Lemina, T. L., Korolev, A. V. (2013). Metabolicheskiy sindrom i porazheniya podzheludochnoy zhelezy. *Rossiyskiy Zhurnal Gastroenterologii, Gepatologii, Koloproktologii*, 2, 17–22.
5. Zvenigorodskaya, L. A., Samsonova, N. G. (2015). Kliniko-dagnosticheskie osobennosti steatoza podzheludochnoy zhelezy u patsientov s metabolicheskim sindromom. *Vestnik semeynoy meditsiny*, 1-2, 32–35.
6. Okovityy, S. V., Napalkova, S. M., Ivkin, D. Yu., Buko, V. U., Nadol'nik, L. I. (2017). Razrabotka vysokoeffektivnykh i bezopasnykh farmatsevticheskikh substantsiy na osnove individual'nykh prirodnykh soedineniy dlya korrektsii narusheniy zhirovoogo i uglevodnogo obmenov. *Metabolicheskiy sindrom: eksperiment, klinika, terapiya*. Grodno: YUrSaPrint, 165–171.
7. Kalmykov, S., Kalmykova, Yu. (2016). Kharakteristika lekarstvennykh rasteniy, primenyaemykh v fitoterapii sakharnogo diabeta 2-go tipa. *Slobozhans'kiy naukovо-sportivniy visnik*, 3 (53), 53–58.
8. Gupta, P. P., Haider, J., Yadav, R. P., Pal, U. (2016). Preclinical evaluation of antidiabetic activity of poly herbal plant extract in streptozotocin induced diabetic rats. *The Journal of Phytopharmacology*, 5 (2), 45–49.
9. Sankar, V., Shoba, F. G. (2017). Antioxidant and antidiabetic activity of Phoenix pusilla Gaertn. unripe fruit extract in streptozotocin-induced sprague dawley rats. *The Journal of Phytopharmacology*, 6 (2), 66–72.
10. Gerber, P. A., Rutter, G. A. (2017). The Role of Oxidative Stress and Hypoxia in Pancreatic Beta-Cell Dysfunction in Diabetes Mellitus. *Antioxidants & Redox Signaling*, 26 (10), 501–518. doi: <http://doi.org/10.1089/ars.2016.6755>
11. Kononenko, N. M., Chikitkina, V. V., Sorokina, M. V., Alkhalaf, M. V. (2018). Eksperymentalne obgruntuvannia vyboru hipohlikemichnoi dozy sukhoho ekstraktu imbyru na normohlike-michnykh shchurakh. *Farmatsevtichnyi zhurnal*, 1-2, 68–77.
12. Kononenko, N. M., Chikitkina, V. V., Sorokina, M. V. (2017). Vyvchennia antyhiperhlikemichnykh vlastyvostei ekstraktu imbyru na eksperymentalnii modeli tsukrovoho diabeta 2 typu, vyklykanoho deksametazonom. *Ukrainskyi biofarmatsevtichnyi zhurnal*, 5 (52), 26–30.
13. Kononenko, N. M., Chikitkina, V. V., Sorokina, M. V., Ostapets, M. O. (2018). The effect of the Ginger dry extract on the indicators of the carbohydrate metabolism under conditions of the experimental metabolic syndrome in Syrian golden hamsters. *Visnik Farmacii*, 1 (93), 49–53. doi: <http://doi.org/10.24959/nphj.18.2200>
14. Zagayko, A., Kravchenko, G., Strelchenko, K., Shkapo, A., Briukhanova, T. (2015). Sex and Age Differences in Lipoprotein Metabolism Proatherogenic Changes under the Experimental Metabolic Syndrome in Hamsters. *Lipoproteins – From Bench to Bedside*, 18–58. doi: <http://doi.org/10.5772/60759>
15. Zapadnyuk, I. P., Zapadnyuk, V. I., Zakhariya, E. A. (1974). *Laboratorye zhyvotnye. Razvedenie, sodержanie, ispol'zovanie v eksperimente*. Kyiv: Vishha shkola, 304.
16. Merkulov, G. A. (1969). *Kurs patologogistologicheskoy tekhniki*. Moscow: Meditsina, Leningr. otd-nie, 424.
17. Bonashevskaya, T. I., Belyaeva, N. N., Kumpan, N. B., Panasyuk, L. V. (1984). *Morfofunktsional'nye issledovaniya v gi-giene*. Moscow, 160.
18. Khalafyan, A. A. (2007). *STATISTICA 6.0 Statisticheskii analiz dannykh*. Moscow: OOO «Binom-Press», 512.
19. Nesterok, Yu. A. (2013). *Morfologicheskoe yssledovanye tryadi orhanov – pechen, podzheludochnaia zheleza y dvednadsatypierstnaia kyshka – pry eksperymentalnom opystorkhoze*. Moscow, 24.
20. Ilkhanizadeh, B., Shirpoor, A., Khadem Ansari, M. hasan, Nemati, S., Rasmi, Y. (2016). Protective Effects of Ginger (*Zingiber officinale*) Extract against Diabetes-Induced Heart Abnormality in Rats. *Diabetes & Metabolism Journal*, 40 (1), 46–53. doi: <http://doi.org/10.4093/dmj.2016.40.1.46>